



UNIVERSIDADE FEDERAL RURAL DO SEMI-ÁRIDO
PRÓ-REITORIA DE PESQUISA E PÓS-GRADUAÇÃO
PROGRAMA DE PÓS-GRADUAÇÃO EM CIÊNCIA ANIMAL
DOUTORADO EM CIÊNCIA ANIMAL

ALANA AZEVEDO BORGES

**ESTABELECIMENTO DE CARIOPLASTOS E CITOPLASTOS DE CATETOS,
Pecari tajacu (LINNAEUS, 1758), VISANDO A CLONAGEM POR TRANSFERÊNCIA
NUCLEAR DE CÉLULA SOMÁTICA EM TAIASSUÍDEOS**

MOSSORÓ

2020

ALANA AZEVEDO BORGES

**ESTABELECIMENTO DE CARIOPLASTOS E CITOPLASTOS DE CATETOS,
Pecari tajacu (LINNAEUS, 1758), VISANDO A CLONAGEM POR TRANSFERÊNCIA
NUCLEAR DE CÉLULA SOMÁTICA EM TAIASSUÍDEOS.**

Tese apresentada ao Doutorado em Ciência Animal do Programa de Pós-Graduação em Ciência Animal da Universidade Federal Rural do Semi-Árido como requisito para obtenção do título de Doutora em Ciência Animal.

Linha de Pesquisa: Morfofisiologia e Biotecnologia Animal

Orientadora: Profa. Dra. Alexsandra Fernandes Pereira.

Co-orientador: Prof. Dr. Alexandre Rodrigues Silva.

MOSSORÓ

2020

© Todos os direitos estão reservados a Universidade Federal Rural do Semi-Árido. O conteúdo desta obra é de inteira responsabilidade do (a) autor (a), sendo o mesmo, passível de sanções administrativas ou penais, caso sejam infringidas as leis que regulamentam a Propriedade Intelectual, respectivamente, Patentes: Lei nº 9.279/1996 e Direitos Autorais: Lei nº 9.610/1998. O conteúdo desta obra tomar-se-á de domínio público após a data de defesa e homologação da sua respectiva ata. A mesma poderá servir de base literária para novas pesquisas, desde que a obra e seu (a) respectivo (a) autor (a) sejam devidamente citados e mencionados os seus créditos bibliográficos.

B732e Borges, Alana Azevedo.
 ESTABELECIMENTO DE CARIOPLASTOS E CITOPLASTOS
 DE CATETOS, Pecari tajacu (LINNAEUS, 1758),
 VISANDO A CLONAGEM POR TRANSFERÊNCIA NUCLEAR DE
 CÉLULA SOMÁTICA EM TAIASSUÍDEOS / Alana Azevedo
 Borges. - 2020.
 263 f. : il.

 Orientadora: Alexsandra Fernandes Pereira.
 Coorientador: Alexandre Rodrigues Silva.
 Tese (Doutorado) - Universidade Federal Rural
 do Semi-árido, Programa de Pós-graduação em
 Ciência Animal, 2020.

 1. Vida selvagem. 2. Clonagem. 3. Biobancos.
 4. Ciclo celular. 5. Ativação oocitária . I.
 Pereira, Alexsandra Fernandes , orient. II.
 Silva, Alexandre Rodrigues , co-orient. III.
 Título.

O serviço de Geração Automática de Ficha Catalográfica para Trabalhos de Conclusão de Curso (TCC's) foi desenvolvido pelo Instituto de Ciências Matemáticas e de Computação da Universidade de São Paulo (USP) e gentilmente cedido para o Sistema de Bibliotecas da Universidade Federal Rural do Semi-Árido (SISBI-UFERSA), sendo customizado pela Superintendência de Tecnologia da Informação e Comunicação (SUTIC) sob orientação dos bibliotecários da instituição para ser adaptado às necessidades dos alunos dos Cursos de Graduação e Programas de Pós-Graduação da Universidade.

ALANA AZEVEDO BORGES

**ESTABELECIMENTO DE CARIOPLASTOS E CITOPLASTOS DE CATETOS,
Pecari tajacu (LINNAEUS, 1758), VISANDO A CLONAGEM POR TRANSFERÊNCIA
NUCLEAR DE CÉLULA SOMÁTICA EM TAIASSUÍDEOS.**

Tese apresentada ao Doutorado em Ciência Animal do Programa de Pós-Graduação em Ciência Animal da Universidade Federal Rural do Semi-Árido como requisito para obtenção do título de Doutora em Ciência Animal.

Linha de Pesquisa: Morfofisiologia e Biotecnologia Animal.

Defendida em: 28/07/2020.

BANCA EXAMINADORA




Prof. Dra. Alessandra Fernandes Pereira (UFERSA)
Presidente e Orientadora




Prof. Dr. Carlos Eduardo Ambrósio (USP)
Membro Examinador



Prof. Dr. Marcelo Bertolini (UFRGS)
Membro Examinador



Prof. Dr. Moacir Franco de Oliveira (UFERSA)
Membro Examinador



Prof. Dr. Alexandre Rodrigues Silva (UFERSA)
Membro Examinador

*Dedico este grau mais alto da minha educação à minha mãe, Elizabeth de Azevedo Silva,
minha maior incentivadora que sempre acreditou e lutou por mim.*

AGRADECIMENTOS

Ao meu amor, Maxson Ariton Sabino da Mota, por estar ao meu lado em todos os momentos desta caminhada, por me acompanhar em cada etapa da construção deste trabalho, por me manter calma em momentos não tão fáceis, por aguardar os momentos de furacão, voltar para emendar os pedaços e tranquilamente me tornar mais forte. Entender distâncias e ausências neste percurso sem exigências tornou tudo menos difícil. Muito obrigada por todo amor e companheirismo.

À minha irmã Amanda Azevedo Borges pela cumplicidade e paciência de me socorrer em todos os momentos sem questionar. Obrigada por apenas entender a importância de cada etapa para mim. Poder ter você como parte fora e diferente de mim, mas que surge de dentro da mesma semente é simplesmente me significar que nunca estarei só.

À minha família, primeiro à minha avó, Francisca de Assis Azevedo, maior alegria e meu melhor sentimento. Mesmo em momentos sem lucidez se alegra em escutar que vou para faculdade, que sou professora e serei doutora. Terei para sempre em minha memória a alegria que a senhora exala. Meus compadres Rosineide Bezerra dos Reis e Antônio Marcos Bezerra de Almeida, por manter nosso laço tão forte com João Firmino Bezerra dos Reis, meu afilhado, que não consigo nem entender de onde ele tira tanto amor por esta madrinha tão ausente. À Marleuza Sabino da Silva que sempre me acolhe com todo carinho, respeito e com um afago de alegria. Muito obrigada por cuidar de mim. À Rosalva Alves Nunes que sempre esteve disposta em facilitar e me proporcionar melhores condições de estudo. Ao meu pai, Armando Borges, pela base educacional, ensinamentos sobre caráter e amor.

À minha mãe Elizabeth de Azevedo Silva, por ser a fonte da força que não se tem como explicar, por acender todas as velas que traduziam a fé que eu sempre vou conseguir, por estar em orações tão perto quando não podia estar fisicamente, mas que torcia para que tudo corresse bem. Quando estando tão perto em seus braços sempre me envolve de amor e paz. Obrigada por toda paciência e tantos sacrifícios para me dar o melhor que podia, por desde cedo insistir em minha educação e me ensinar a paixão por essa vida. A senhora sempre será minha fonte de inspiração, meu porto seguro e minha maior motivação. Não existiria essa Alana sem Elizabeth, mãe, amiga, mulher e amor.

À minha orientadora Alexsandra Fernandes Pereira, não sei ainda como é traduzir a confiança que você tem em mim, não sei como você me enxergou e me confiou este trabalho que é um passo tão bonito na sua história. Eu sou muito grata em me possibilitar de coração aberto tanto aprendizado. Tenho cada lição tatuada em minha alma, sou muito honrada em ser um capítulo desta história, e aspiro ser uma mulher tão forte, determinada e comprometida como és. Nessa caminhada de alguns anos, construímos uma linda amizade, que tenho certeza de que terei sempre comigo. Obrigada por se dispor a escutar tantas dúvidas e anseios, por ser sempre tão gentil e calma em me fazer acreditar e me recuperar em cada desafio que aparecia. Você ensina até o querer, que muitas vezes você acha que não pode ensinar, mas você pode inspirar a acharmos. Seu coração alegre e cheio de amor fez com que tudo fosse muito leve, porque se segundo a Frida Kahlo que nos inspiramos “Donde no puedes amar, no te demores”, junto com você e com o amor que trata tudo... nos quedamos siempre. Juntas vencemos muito mais do que imaginamos. É difícil descrever toda a gratidão, por um

aprendizado de uma vida. Assim, lutarei todos os dias para ser uma Doutora do jeito que você me mostra como é ser uma DOUTORA.

Aos meus parceiros de caminhada, que primeiro era para eu “ensinar” e eles brilhantemente me fizeram aprender muito mais do que eu poderia fazer por eles, sem a colaboração de vocês este trabalho não seria tão brilhante. À Matheus Barbosa do Nascimento, pela disponibilidade e compreensão, por seu meu anjo da guarda, estando sempre presente para ajudar no que fosse possível. À Lucas Emanuel Nascimento por estar desde o mestrado, juntos desvendando os mistérios que só o tempo dirá, agradeço toda sua paciência e dedicação. À Gabriela Pereira de Oliveira Lira, por estar junto tão junto que nem sei mais o tanto de você que tenho em mim, obrigada por lutar sempre para me dar o seu melhor, por aguentar a pressão de ser meu terceiro olho. Ganhei uma amiga que me desconsertou e me trouxe alegria de me ver como sou e não como estava sendo. Este trabalho é fruto de nosso esforço e dedicação. Muito obrigada por toda confiança e amor que construímos durante esse trabalho, vocês são o meu florescer!

Aos meus companheiros de Laboratório, minha casa desde 2013 onde era apenas um quadrado e virou o grandioso Laboratório de Biotecnologia Animal (LBA). Nessa casa, há paixão e paixões não são fáceis. Obrigada pela paciência em entender que só tinha que dar certo e fiz ao meu máximo. À Luiza Bento de Queiroz Neta que começamos juntas trilhando essa história, vencendo desafios e buscando nossos sonhos. À Maria Valéria de Oliveira Santos, obrigada por estar sempre disposta a ajudar e por estar presente em etapas determinantes neste trabalho, sua ajuda foi essencial. Esse trio parada-dura do LBA sempre será uma amizade para vida toda! Assim, agradeço a todo o LBA em nome do meu amigo Leonardo Vitorino Costa de Aquino, que traz consigo alegria e alivia com seu sorriso todas as dificuldades, obrigada por sempre se importar comigo.

Ao meu co-orientador, Alexandre Rodrigues Silva, que me acompanha e colabora para o meu crescimento desde a iniciação científica, sempre apostando muito em mim. Sou grata as lições e desafios que me motivaram a refletir sobre ser um PhD. Sua colaboração neste trabalho foi essencial para que esta semente germinasse e tivesse tão belos frutos. Vejo muito além das palavras às vezes duras, sempre as interpretei como zelo. Assim, agradeço o carinho e a confiança que sempre depositou.

A todos que compõe o Centro de Multiplicação de Animais Silvestres (CEMAS), por me receber tão bem e confiar em nosso trabalho. Sou muito grata em poder ter trabalhado com Senhor Almeida, um homem que é como se descreve um sertanejo, uma casca grossa, mas um coração enorme. Obrigada por tantas colheitas desde a iniciação científica. Ao professor Moacir Franco de Oliveira que tenho muita gratidão e admiração por sempre ser sereno e humilde, estando em todas as colheitas, ensinando e fazendo de tudo para dar certo, agradeço muito por acreditar no nosso trabalho e por todo carinho.

À banca examinadora pela disponibilidade em estar presente neste momento tão importante e por todas as contribuições neste trabalho.

À Universidade Federal Rural do Semi-Árido (UFERSA) por meio do Programa de Pós-Graduação em Ciência Animal (PPGCA) pela formação profissional.

Aos laboratórios parceiros: Laboratório de Conservação de Germoplasma Animal (LGCA/UFERSA), Laboratório de Morfofisiologia Animal Aplicada (LMAA/UFERSA), e Laboratório de Oncologia Experimental (LOE), da Universidade Federal do Ceará (UFC) pela disponibilidade da infraestrutura física para a realização de alguns experimentos, especialmente à professora Claudia Pessoa e professora Maria Claudia dos Santos Luciano.

À Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) pela concessão de bolsa de doutorado e ao Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) pelo auxílio financeiro.

Aos animais, inicialmente, aos catetos, espécie que me doou muitos ensinamentos em minha trajetória científica. Espero poder contribuir com esta espécie o tanto que me dedicou. À minha cadela Frida, pela companhia e amor.

Finalmente, agradeço à Deus, a todos Anjos e Santos, à Santa Luzia e a todas as forças da natureza por me iluminarem nesta caminhada. Que nunca me falte a fé.

Não sei se a vida é curta ou longa para nós,
mas sei que nada do que vivemos tem sentido,
se não tocarmos o coração das pessoas.

Muitas vezes basta ser: colo que acolhe, braço
que envolve, palavra que conforta, silêncio que
respeita, alegria que contagia, lágrima que
corre, olhar que acaricia, desejo que sacia,
amor que promove.

E isso não é coisa de outro mundo, é o que dá
sentido à vida. É o que faz com que ela não
seja nem curta, nem longa demais, mas que
seja intensa, verdadeira, pura enquanto durar.

Cora Coralina

RESUMO

O direcionamento dos primeiros passos para a realização da transferência nuclear de célula somática (TNCS) em catetos garantirá uma ferramenta efetiva na conservação da espécie, perante sua acelerada diminuição populacional e sua atividade ecológica indispensável para o ecossistema. Para tanto, a presente tese foi dividida em duas etapas (três experimentos por etapa), sendo a primeira etapa o estudo das células doadoras de núcleo ou carioplastos e a segunda etapa, o estudo das células doadoras de citoplasma ou citoplastos. Assim, diante da importância de carioplastos de qualidade reconhecida para a TNCS, nós inicialmente estabelecemos cinco linhagens de fibroblastos de catetos, monitorando a viabilidade, atividade metabólica e estresse oxidativo, de acordo com os efeitos do número de passagens (primeira, terceira e décima) e criopreservação. Embora não haja efeito desses critérios sobre a viabilidade, células em décima passagem tiveram uma redução de seu metabolismo. Adicionalmente, células congeladas/descongeladas tiveram um aumento no número de espécies reativas de oxigênio e potencial de membrana mitocondrial. Além disso, sabendo da importância de manter estas células armazenadas em um biobanco de maneira adequada, nós otimizamos a solução crioprotetora utilizada na congelação lenta de fibroblastos de catetos. Deste modo, a solução composta por 10% de dimetilsulfóxido (DMSO) com 0,2 M de sacarose e 50% de soro fetal bovino (SFB) foi considerada a solução mais eficiente em manter a viabilidade, atividade proliferativa, metabolismo e níveis adequados de estresse oxidativo de células somáticas de catetos, quando comparada a soluções na ausência de sacarose e com 10% de SFB em diferentes combinações. Finalmente, um passo essencial no estabelecimento dos carioplastos para a TNCS consiste na sincronização das células em G_0/G_1 do ciclo celular. Deste modo, nós avaliamos diferentes métodos de sincronização do ciclo celular: (i) supressão de soro (SS) por um a quatro dias, (ii) inibição por contato (IC) por um a três dias e (iii) agentes químicos [DMSO, 6-dimetilaminopurina (6-DMAP), ciclohexamida (CHX), e citocalasina B (CB)] por um a dois dias, em termos de seus efeitos sobre a sincronização em G_0/G_1 e viabilidade. Assim, nós observamos que a IC por três dias foi o método mais eficiente para sincronização do ciclo celular e manutenção da viabilidade de fibroblastos de catetos. Portanto, com estes três experimentos, nós estabelecemos a etapa de carioplastos da TNCS de catetos, obtendo células de qualidade e aptas a serem usadas como doadoras de núcleo. Na segunda etapa, nós, inicialmente, adequamos as condições de maturação *in vitro* (MIV) de oócitos de catetos, avaliando o tempo de MIV e o efeito do fator de crescimento epidermal (EGF) sobre a habilidade meiótica. Assim, nós concluímos que 48 h é o período adequado para a MIV de oócitos quando comparado ao tempo de 24 h, de acordo com o potencial meiótico. Ainda, observou-se que o EGF pode ser utilizado para otimizar o meio de MIV. Finalmente, no terceiro experimento, nós avaliamos a habilidade de desenvolvimento destes oócitos após ativação artificial, usando a ionomicina como ativador primário e comparando diferentes ativadores secundários (6-DMAP, CHX e CB). Nós verificamos que a ativação química usando ionomicina e 6-DMAP foi a mais eficiente combinação, tendo esta tese alcançado como resultado significativo, uma taxa de 27,6% de blastocistos de catetos derivados da ativação oocitária artificial. Em síntese, nós obtivemos carioplastos e citoplastos que poderão ser empregados na TNCS de catetos, deixando a ponto as etapas fundamentais para a clonagem desta espécie. Ainda, destaca-se que os conhecimentos aqui gerados poderão ser aplicados em estudos de fecundação *in vitro*, compreensão do desenvolvimento embrionário, produção de células induzidas à pluripotência, e ensaios de toxicidade. Portanto, este trabalho foi um grande passo para a conservação de catetos.

Palavras-chave: Vida selvagem. Clonagem. Biobancos. Ciclo celular. Ativação oocitária. Produção de embriões.

ABSTRACT

The direction of the first steps for the achievement of somatic cell nuclear transfer (SCNT) in collared peccary will guarantee an effective tool in the conservation of the species, in view of its accelerated population decrease and its essential ecological activity for the ecosystem. Therefore, the present thesis was divided into two steps (three experiments per step), being the first step the study of the donor cells of nucleus or karyoplast and the second stage, the study of the donor cells of cytoplasm or cytoplasts. Thus, in view of the importance of acknowledge quality karyoplast for SCNT, we initially established five cell lines of collared peccary fibroblasts, monitoring viability, metabolic activity and oxidative stress, according to the effects of the number of passages (first, third and tenth) and cryopreservation. Although there is no effect of these criteria on viability, cells in tenth passage had a reduction in their metabolism. Additionally, frozen/thawed cells had an increase in the number of reactive oxygen species and mitochondrial membrane potential. Moreover, knowing the importance of maintaining these cells stored in a biobank properly, we optimize the cryoprotectant solution used in the slow freezing of collared peccary fibroblasts. Thus, the solution composed of 10% dimethyl sulfoxide (DMSO) with 0.2 M sucrose and 50% fetal bovine serum (FBS) was considered the most efficient solution in maintaining the viability, proliferative activity, metabolism and adequate levels of oxidative stress of somatic cell cells, when compared to solutions in the absence of sucrose and with 10% FBS in different combinations. Finally, an essential step in establishing the karyoplast for SCNT is the synchronization of cells in G₀/G₁ of the cell cycle. Thus, we evaluated different cell cycle synchronization methods: (i) serum suppression (SS) for one to four days, (ii) contact inhibition (CI) for one to three days and (iii) chemical agents [DMSO, 6-dimethylaminopurine (6-DMAP), cyclohexamide (CHX), and cytochalasin B (CB)] for one to two days, in terms of their effects on G₀/G₁ synchronization and viability. Thus, we observed that the IC for three days was the most efficient method for synchronizing the cell cycle and maintaining the viability of collared peccary fibroblasts. Consequently, with these three experiments, we have established karyoplast stage of SCNT in collared peccary, obtaining quality cells and able to be used as nuclear donors. In the second stage, we initially adjusted the *in vitro* maturation (IVM) conditions of collared peccary oocytes, evaluating the IVM time and the effect of the epidermal growth factor (EGF) on the meiotic ability. Thus, we concluded that 48 h is the appropriate period for oocyte IVM when compared to 24 h, according to meiotic potential. Still, it was observed that EGF can be used to optimize the IVM medium. Finally, in the third experiment, we evaluated the developmental ability of these oocytes after artificial activation, using ionomycin as the primary activator and comparing different secondary activators (6-DMAP, CHX and CB). We found that chemical activation using ionomycin and 6-DMAP was the most efficient combination, with this thesis achieving as a significant result, a rate of 27.6% of blastocysts of collared peccaries derived from oocyte artificial activation. In summary, we got karyoplast and cytoplasts that may be employed in the SCNT of collared peccary, leaving the point the fundamental steps for the cloning of this species. Furthermore, it is emphasized that the knowledge generated here can be applied for *in vitro* fertilization, studies understanding of embryonic development, production cells induced to pluripotency, and toxicity assessments. Therefore, this work was a great step for the conservation of collared peccaries.

Keywords: Wild life. Cloning. Biobanks. Cell cycle. Oocyte activation. Embryo production.

LISTA DE FIGURAS

CAPÍTULO 1 – CONSIDERAÇÕES GERAIS

Figura 1	– Estratégias de conservação <i>in situ</i> desenvolvidas em catetos.....	34
Figura 2	– Características reprodutivas de catetos a partir de estudos <i>in vitro</i> sobre a morfofisiologia do macho.....	38
Figura 3	– Evolução do conhecimento sobre aspectos reprodutivos de fêmeas de <i>P. tajacu</i> investigados <i>in vitro</i>	38
Figura 4	– Caminho percorrido no estabelecimento de amostras somáticas de catetos.....	42

CAPÍTULO 3 – ISOLATION, CHARACTERIZATION, AND CRYOPRESERVATION OF COLLARED PECCARY SKIN-DERIVED FIBROBLAST CELL LINES

Figure 1	– Outgrowth of fibroblast cells from ear skin samples of collared peccaries. Exhibit a skin explant cultured in (A) day 1, (B) day 3 and (C) day 5 of primary culture and exhibit a fibroblast population cultured in (D) day 15 and (E) day 19 of subculture. (F) exhibit cells after the trypsinization process.....	93
Figure 2	– Immunocytochemical detection of vimetin protein for identification of collared peccary fibroblasts. (A–D) cells stained with vimetin antibody. (B–E) nucleus of cells stained by Hoechst. (C–F) merged vimetin (green) and Hoechst (blue).....	95
Figure 3	– The growth dynamics of cryopreserved and non-cryopreserved cells derived from collared peccary skin. (A) growth curves of cryopreserved and non-cryopreserved cells. (B) values of population doubling time (PDT) after culture for nine days.....	97
Figure 4	– Evaluation of intracellular reactive oxygen species (ROS) and mitochondrial membrane potential ($\Delta\Psi_m$). Cell stained using fluorescent probe (A) 2',7'-dichlorodihydrofluorescein diacetate (H ₂ DCFDA) and (B) MitoTrackerRed® (CMXRos). Quantification of (C) ROS and (D) $\Delta\Psi_m$ level.....	98

CAPÍTULO 4 – CRYOPRESERVATION OF COLLARED PECCARY (*Pecari tajacu* LINNAEUS, 1758) SOMATIC CELLS IS IMPROVED BY SUCROSE AND HIGH CONCENTRATIONS OF FETAL BOVINE SERUM

- Figure 1** – Subcultures of fibroblast-like cells from ear skin samples of collared peccaries. **a)** Non-cryopreserved cells and cell cryopreserved in **b)** DMSO, **c)** DMSO-SUC, **d)** DMSO-10FBS, **e)** DMSO-50FBS, **f)** DMSO-SUC-10FBS, **g)** DMSO-SUC-50FBS. 116
- Figure 2** – Viability of collared peccary non-cryopreserved and cryopreserved cells using different extracellular cryoprotectants. Non-cryopreserved group, 10% DMSO (DMSO group), 10% DMSO and 0.2 M SUC (DMSO-SUC group), 10% DMSO and 10% FBS (DMSO-10FBS group), 10% DMSO and 50% FBS (DMSO-50FBS group), 10% DMSO, 0.2 M SUC and 10% FBS (DMSO-SUC-10FBS group), 10% DMSO, 0.2 M SUC and 50% FBS (DMSO-SUC-10FBS group).....117
- Figure 3** – The growth dynamics and metabolism of collared peccary non-cryopreserved and cryopreserved cells using different extracellular cryoprotectants. **a)** Growth curves of cryopreserved and non-cryopreserved cells. **b)** Values of population doubling time (PDT) after culture for nine days. **c)** Values of metabolic activity in percentage. Non-cryopreserved group, 10% DMSO (DMSO group), 10% DMSO and 0.2 M SUC (DMSO-SUC group), 10% DMSO and 10% FBS (DMSO-10FBS group), 10% DMSO and 50% FBS (DMSO-50FBS group), 10% DMSO, 0.2 M SUC and 10% FBS (DMSO-SUC-10FBS group), 10% DMSO, 0.2 M SUC and 50% FBS (DMSO-SUC-10FBS group).....118
- Figure 4** – Assessment of oxidative stress in collared peccary non-cryopreserved and cryopreserved cells using different extracellular cryoprotectants. Non-cryopreserved group, 10% DMSO (DMSO group), 10% DMSO and 0.2 M SUC (DMSO-SUC group), 10% DMSO and 10% FBS (DMSO-10FBS group), 10% DMSO and 50% FBS (DMSO-50FBS group), 10% DMSO, 0.2 M SUC and 10% FBS (DMSO-SUC-10FBS group), 10% DMSO, 0.2 M SUC and 50% FBS (DMSO-SUC-10FBS group). **a)** Quantification of intracellular

reactive oxygen species (ROS) levels	b) Measurement of mitochondrial membrane potential ($\Delta\Psi_m$).	c) Cells labeled with H2DCFDA for quantification of ROS levels.	d) Cells labeled with MitoTracker Red® for measurement of $\Delta\Psi_m$	119
--------------------------------------	---	--	---	-----

CAPÍTULO 5 – EFFECTS OF INCUBATION TIME AND METHOD OF CELL CYCLE SYNCHRONIZATION ON COLLARED PECCARY SKIN-DERIVED FIBROBLAST CELL LINES

Figure 1	– Culture of collared peccary fibroblast cells. (A) Fibroblasts at 70% confluency showing normal morphology. (B) Growth curve of fibroblasts.....	133
Figure 2	– Representative histograms of the flow cytometry analysis of collared peccary fibroblasts. (A) growing cells (GC, control). (B) serum starvation (SS). (C) contact inhibition (CI). (D) dimethyl sulfoxide (DMSO). (E) cycloheximide (CHX). (F) cytochalasin B (CB). (G) 6-dimethylaminopurine (6-DMAP)...	134
Figure 3	– Effect of different synchronization methods on the viability of collared peccary fibroblasts. (A) and (A') Effect of SS on fibroblast viability. (B) and (B') Evaluation of the viability of fibroblasts subjected to CI. (C) and (C') Effect of cell cycle inhibitors on the viability of synchronized fibroblasts.....	136

CAPÍTULO 6 – *IN VITRO* MATURATION OF COLLARED PECCARY (*Pecari tajacu*) OOCYTES AFTER DIFFERENT INCUBATION TIMES

Figure 1	– Macroscopic aspects of collared peccary ovaries. (A) Uniform appearance with small follicles. (B) Presence of large follicles and corpus luteum. The arrowhead indicates a small follicle and arrow shows a large follicle.....	153
Figure 2	– <i>In vitro</i> maturation of oocytes of collared peccary. (A) Immature oocytes before IVM (100X); (B) <i>Cumulus</i> cell expansion after 24 h IVM (100X); (C) <i>Cumulus</i> cell expansion after 48 h IVM (100X); (D) Oocyte with extrusion of first polar body (400X); (E) Oocyte in MII stage (400X).....	154

CAPÍTULO 7 – PRODUCTION OF COLLARED PECCARY (*Pecari tajacu* LINNAEUS, 1758) PARTHENOGENIC EMBRYOS FOLLOWING DIFFERENT OOCYTE CHEMICAL ACTIVATION AND *IN VITRO* MATURATION CONDITIONS

- Figure 1** – Representative images of (A) ovaries of collared peccary obtained after hormonal induction with PG 600®; (B) *Cumulus*-oocyte complex recovered with homogeneous cytoplasm and more one layer of *cumulus* cells; (C) COCs matured after 44 h IVM; (D) Denuded oocyte with presence of 1PB; (E) Visualization of the metaphasic plate (MII). 172
- Figure 2** – Oocytes of collared peccaries after maturation evaluated as intracellular concentration of reactive oxygen species (ROS) levels and mitochondrial membrane potential ($\Delta\Psi_m$). (A) Measurement of fluorescence of ROS levels in matured oocytes in the presence of EGF (EGF positive group) or in the absence (EGF negative group). (B) Quantification of mitochondrial membrane potential in oocytes matured in the presence of EGF (EGF positive group) or in the absence (EGF negative group). (C) Oocytes labelled with fluorescent probe H2DCFDA for ROS quantification with EGF (EGF positive group) and (D) without EGF (EGF negative group). (E) Oocytes labelled with fluorescent probe MitoTracker Red® for $\Delta\Psi_m$ with EGF (EGF positive group) and (F) without EGF (EGF negative group).....176
- Figure 3** – Kinetics of embryonic development in collared peccaries. After 3 days of IVD different embryos with (A) 2 cells; (B) 4 cells, (C) more than 8 cells. Evaluation of embryonic development on day 7 of IVD (D) blastocyst; (E) blastocyst hatching; (F) blastocyst hatched already without zona pellucida. Counting the total number of blastocyst (G) cells inner mass labelled with Hoechst 33342. (H) trophectoderm cells stained with propidium iodide. (I) merge total cells..... 178
- Figure 4** – Embryonic development of classified activated with ionomycin in combination with different secondary activators. Embryonic development in 3 days with 2 cells, 3 e 7 cells and 8 cells after activation with ionomycin in combination with secondary activators in (A) GI/GII and (B) GIII COCs. (C)

Percentage of embryos in D7 classified in different stages of blastocyst after activation of GI/GII COCs.....	181
--	-----

LISTA DE QUADROS

CAPÍTULO 1 – CONSIDERAÇÕES GERAIS

Quadro 1	–	Ocorrência e distribuição de subespécies de catetos.....	31
Quadro 2	–	Desenvolvimento de estratégias <i>ex situ in vivo</i> em catetos.....	36
Quadro 3	–	Estratégias <i>ex situ in vitro</i> aplicadas em catetos.....	40

LISTA DE TABELAS

CAPÍTULO 2 – POTENTIAL ROLE OF INTRASPECIFIC AND INTERSPECIFIC CLONING IN THE CONSERVATION OF WILD MAMMALS

Table 1	– iSCNT in some wild mammals	69
----------------	------------------------------------	----

CAPÍTULO 3 – ISOLATION, CHARACTERIZATION, AND CRYOPRESERVATION OF COLLARED PECCARY SKIN-DERIVED FIBROBLAST CELL LINES

Table 1	– Establishment of primary culture and subcultures of somatic cells derived from collared peccary ear skin.....	94
----------------	---	----

CAPÍTULO 5 – EFFECTS OF INCUBATION TIME AND METHOD OF CELL CYCLE SYNCHRONIZATION ON COLLARED PECCARY SKIN-DERIVED FIBROBLAST CELL LINES

Table 1	– Effect of serum starvation (SS) on the percentage of collared peccary fibroblasts in the G0/G1, S and G2/M phases of the cycle.....	135
Table 2	– Effect of contact inhibition (CI) on the percentage of collared peccary fibroblasts in the G0/G1, S and G2/M phases of the cycle.....	138
Table 3	– Effect of treatment with dimethyl sulfoxide (DMSO), cycloheximide (CHX), cytochalasin B (CB), or 6-dimethylaminopurin (6-DMAP) on the percentage of collared peccary fibroblasts in the G0/G1, S and G2/M phases of the cycle.....	139

CAPÍTULO 6 – *IN VITRO* MATURATION OF COLLARED PECCARY (*Pecari tajacu*) OOCYTES AFTER DIFFERENT INCUBATION TIMES

Table 1	– <i>In vitro</i> maturation of collared peccary oocytes after different incubation times.....	155
----------------	--	-----

CAPÍTULO 7 – PRODUCTION OF COLLARED PECCARY (*Pecari tajacu* LINNAEUS, 1758) PARTHENOGENIC EMBRYOS FOLLOWING DIFFERENT OOCYTE CHEMICAL ACTIVATION AND *IN VITRO* MATURATION CONDITIONS

Table 1	– Effects of EGF supplementation on <i>in vitro</i> maturation of collared peccary <i>cumulus</i> -oocyte complex (COCs).....	174
Table 2	– Morphometric assessment (in μm and μm^2) in of matured COCs derived from collared peccaries (mean $\pm$ standard error).....	175
Table 3	– Embryonic development of collared peccary GI/GII and GIII <i>cumulus</i> -oocyte complexes (COCs) activated with ionomycin in combination with different secondary activators.....	177
Table 4	– Cell count of <i>cumulus</i> -oocyte complexes (COCs) as viable and activated with ionomycin in combination with different secondary activators.....	180

LISTA DE ABREVIATURAS E SIGLAS

° C	Grau Celsius
µg	Micrograma
µL	Microlitro
µm	Micrometro
µM	Micromolar
1 PB	First polar body
10th	Tenth
15th	Fiftieth
4th	Fourth
5th	Fifth
6D	6-dimethylaminopurine
6-DMAP	6-dimethylaminopurine
ANOVA	Variance analysis
art.	Artigo
ARTs	Assisted Reproductive Techniques
Av.	Avenida
B	Blastocyst
Bak	Pro-apoptotic
Bcl-xL	Anti-apoptotic
BSA	Bovine Serum Albumin
CA	Califórnia
Ca ²⁺	Íon cálcio
cAMP	3',5' cyclic adenosine monophosphate
CAPES	Coordenação de Aperfeiçoamento de Pessoal de Nível Superior
CB	Cytochalasin B
CCM	<i>Cumulus</i> -oocyte complex collection medium
CEUA	Committee for Ethics in Animal Use
CHX	Cycloheximide
CI	Contact inhibition
cm	Centímetro
cm ²	Centímetro quadrado

CMXRos	MitoTracker Red®
CNPq	Conselho Nacional de Desenvolvimento Científico e Tecnológico
CO ₂	Dióxido de carbono
COC	<i>Cumulus</i> -oocyte complex
D0	Day 0
D3	Day 3
D7	Day 7
DMEM	Dulbecco's modified Eagle medium
DMSO	Dimethyl sulfoxide
DNA	Deoxyribonucleic acid
eB	Expanded blastocyst
eCG	Equine chorionic gonadotropin
EDTA	Ethylenediamine tetraacetic acid
EGF	Epidermal growth factor
et al.	E outros
FBS	Fetal bovine serum
Fig.	Figure
FSH	Follicle-stimulating hormone
g	G force
g	Grama
G	Gauge
Gbif	Biodiversity Information Facility
GC	Growing cells
GI	Grade I
GI	Grade I
GII	Grade II
GIII	Grade III
GIV	Grade IV
GSH	Glutathione
h	Hour
H2DCFDA	2',7'-dichlorodihydrofluorescein diacetate
H ₂ O ₂	Hydrogen peroxide
H3K18ac	Histone H3 at lysine 18

H3K27me3	Histone H3 at lysine 27
hB	Blastocyst hatching/hatched
hCG	Human Chorionic Gonadotropin Hormone
IA	Inner oocyte area
iB	Initial blastocyst
IBAMA	Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis
IC	Inibição de contato
ICM	Inner cellular mass
ICMBio	Instituto Chico Mendes de Conservação da Biodiversidade
ICSI	Intracytoplasmic sperm injection
IgG	Immunoglobulin G
iSCNT	Intraspecific and interspecific somatic cell nuclear transfer
IU	International units
IUCN	International Union for Conservation of Nature
IVD	<i>In vitro</i> development
IVEP	<i>In vitro</i> embryo production
IVF	<i>In vitro</i> fertilization
IVM	<i>In vitro</i> maturation
L	Litro
LH	Luteinizing hormone
M	Mol
MG	Minas Gerais
mg	Miligrama
MII	Metaphase II
min	Minute
MIV	Maturação <i>in vitro</i>
mL	Mililitro
mM	Milimolar
mm	Milímetro
mm3	Milímetro cúbico
MMA	Ministério do Meio Ambiente
MO	Missouri
MPF	Maturation Promoter Factor

MSD	Merck Sharp & Dohme
MtDNA	DNA mitochondrial
MTT	3-(4, 5-dimethylthiazolyl-2)-2, 5-diphenyltetrazolium bromide
n	Número
ng	Nanogram
nm	Nanometro
no	Número
NY	New York
OA	Ooplasm area
OD	Ooplasm diameter
OTC	Ovarian Tissue Cryosystem
P	Probabilidade de significância
PBS	Phosphate buffered saline
PDT	Population doubling time
pH	Hydrogen potential
PI	Propidium iodide
Ptskf	Fibroblast lines derived from adult collared peccaries
PVS	Perivitelline space
Rebio	Reserva Biológica
RN	Rio Grande do Norte
RNA	Ribonucleic acid
ROS	Reactive Oxygen Species
SCNT	Somatic Cell Nuclear Transfer
SOF	Synthetic Oviductal Fluid
SP	São Paulo
SS	Serum starvation
SS	Supressão de soro
SE	Standard error
TCM 199	Medium 199
TE	Trophectoderm
TNCS	Transferência Nuclear de Célula Somática
UFERSA	Universidade Federal Rural do Semi-Árido
USA	United States

UV	Ultraviolet
VDC	Vitrificação direta em criotubos
<i>vs.</i>	Versus
VSS	Vitrificação em superfície sólida
w/v	Weight per volume
ZPI	Inner oocyte diameter
ZPO	Oocyte diameter
ZPT	Zona pellucida thickness
$\Delta\Psi_m$	Mitochondrial membrane potential

LISTA DE SÍMBOLOS

@	Arroba
®	Marca registrada
%	Porcentagem
+	Soma
<	Menor que
=	Igual
>	Maior que
±	Mais ou menos
×	Multiplicação

SUMÁRIO

CAPÍTULO 1 – CONSIDERAÇÕES GERAIS	28
1 INTRODUÇÃO.....	28
2 FUNDAMENTAÇÃO TEÓRICA	30
2.1 Aspectos ecológicos e quantitativo populacional dos catetos.....	30
2.2 Estratégias de conservação <i>in situ</i> aplicadas em catetos.....	32
2.3 Estratégias de conservação <i>ex situ</i> aplicadas em catetos.....	35
2.3.1 Estratégias de conservação <i>ex situ in vivo</i> aplicadas em catetos	35
2.3.2 Estratégias de conservação <i>ex situ in vitro</i> aplicadas em catetos.....	37
3 JUSTIFICATIVA.....	43
4 HIPÓTESES CIENTÍFICAS	44
5 OBJETIVOS	45
5.1 Objetivo Geral.....	45
5.2 Objetivos Específicos.....	45
REFERÊNCIAS.....	46
CAPÍTULO 2 – POTENTIAL ROLE OF INTRASPECIFIC AND INTERSPECIFIC CLONING IN THE CONSERVATION OF WILD MAMMALS	60
CAPÍTULO 3 – ISOLATION, CHARACTERIZATION, AND CRYOPRESERVATION OF COLLARED PECCARY SKIN-DERIVED FIBROBLAST CELL LINES	82
CAPÍTULO 4 – CRYOPRESERVATION OF COLLARED PECCARY (<i>Pecari tajacu</i> LINNAEUS, 1758) SOMATIC CELLS IS IMPROVED BY SUCROSE AND HIGH CONCENTRATIONS OF FETAL BOVINE SERUM.....	108
CAPÍTULO 5 – EFFECTS OF INCUBATION TIME AND METHOD OF CELL CYCLE SYNCHRONIZATION ON COLLARED PECCARY SKIN-DERIVED FIBROBLAST CELL LINES.....	126
CAPÍTULO 6 – <i>IN VITRO</i> MATURATION OF COLLARED PECCARY (<i>Pecari tajacu</i> LINNAEUS, 1758) OOCYTES AFTER DIFFERENT INCUBATION TIMES	146
CAPÍTULO 7 – PRODUCTION OF COLLARED PECCARY (<i>Pecari tajacu</i> LINNAEUS, 1758) PARTHENOGENIC EMBRYOS FOLLOWING DIFFERENT OOCYTE CHEMICAL ACTIVATION AND <i>IN VITRO</i> MATURATION CONDITIONS.....	162
6 CONSIDERAÇÕES FINAIS E PERSPECTIVAS	191
ANEXO A – VERSÃO DO ARTIGO PUBLICADO NA ZYGOTE.....	192
ANEXO B – VERSÃO DO ARTIGO PUBLICADO NA PEERJ	200
ANEXO C – VERSÃO DO ARTIGO PUBLICADO NA CRYO-LETTERS.....	219
ANEXO D – VERSÃO DO ARTIGO PUBLICADO NA PESQUISA VETERINÁRIA BRASILEIRA	229
ANEXO E – VERSÃO DO ARTIGO PUBLICADO NA THERIOGENOLOGY	236
ANEXO F – COMPROVANTE DO ARTIGO SUBMETIDO NA ANNALS OF ANIMAL SCIENCE	245
ANEXO G – RESUMOS PUBLICADOS NO RENORBIO 2017	248
ANEXO H – RESUMO PUBLICADO NO SEMIC 2018.....	251

ANEXO I – RESUMOS PUBLICADOS NO CONERA 2018.....	253
ANEXO J – RESUMO PUBLICADO NO ISABR 2018.....	255
ANEXO L – RESUMO PUBLICADO NO CBRA 2019.....	257
ANEXO M – RESUMOS PUBLICADOS NO RENORBIO 2019	259
ANEXO N – RESUMO PUBLICADO NO SEMIC 2019	262

CAPÍTULO 1 – CONSIDERAÇÕES GERAIS

1 INTRODUÇÃO

A clonagem por transferência nuclear de célula somática (TNCS) consiste em uma biotécnica de importante papel na conservação da biodiversidade silvestre (PRIYA et al., 2014; WANI et al., 2018). Tal relevância pode ser representada por seus desmembramentos atrelados à multiplicação de reprodutores com melhores características genéticas (HUAN et al., 2015), obtenção de células induzidas à pluripotência (NOVO et al., 2016), investigação de mecanismos do desenvolvimento embrionário (HUANG et al., 2017), e edição genética (LIU et al., 2019). Por essas razões, a TNCS tem se tornado uma alternativa interessante para a conservação de diferentes espécies de mamíferos silvestres, como o elefante asiático (TECHAKUMPHU et al., 2018) e camelo bactriano (WANI et al., 2018), podendo assim ser também aplicada em catetos e em outros taiassuídeos.

Em geral, os catetos são subestimados quanto a sua importância ecológica, na qual na maioria das vezes utilizam-se de argumentos como a sua classificação na União Internacional da Conservação da Natureza (IUCN), a qual afirma que a espécie é classificada como pouco preocupante em nível mundial (GONGORRA et al., 2011). Tal classificação muitas vezes se mostra defasada e não representativa em vários biomas (NAGY-REIS et al., 2019). Assim, a urgência pelo desenvolvimento de biotécnicas que garantam estratégias de conservação para a espécie é um tema relevante. Adicionalmente, os catetos possuem alta adaptação a cativeiros, sendo ainda indivíduos interessantes economicamente pela apreciação da sua carne e couro (BODMER et al., 1990). Diante deste cenário, ter a ponto as etapas iniciais da TNCS é uma alternativa importante, uma vez que as etapas de preparo de células doadoras de núcleo (carioplastos) e citoplasmas receptores (citoplastos) poderão ser também ampliadas em outras biotécnicas, além de permitir a geração de conhecimento na biologia reprodutiva dos catetos e possivelmente ser extrapolada para outros taiassuídeos.

Embora várias pesquisas tenham demonstrado o sucesso da TNCS pelo nascimento de crias nascidas, tanto em mamíferos domésticos (BUOMO et al., 2016), quanto silvestres (FOLCH et al., 2009), a eficiência da técnica ainda é reduzida, sendo atribuída a seleção e a manipulação dos carioplastos (KWONG et al., 2014) e citoplastos (GARCIA-MENGUAL et al., 2008) como alguns dos fatores que devem ser estabelecidos na espécie de interesse. Ainda, é importante ressaltar que as etapas da TNCS possuem íntima relação com as

1 características biológicas da espécie, sendo importante definir e estabelecer modificações
2 adequadas nos protocolos desenvolvidos (MOHAMMED et al., 2019).

3 Assim, a otimização da TNCS pela escolha de carioplastos eficientes envolve o
4 conhecimento das alterações sofridas nas células durante seu cultivo *in vitro*, ou seja, o
5 estabelecimento das condições de incubação *in vitro* e a avaliação do potencial de
6 conservação das características normais após esse período (SONG et al., 2007). Portanto, ter
7 linhagens celulares definidas e um protocolo de criopreservação adequado que garantam que
8 essas células possam ser armazenadas para uso futuro consiste em um fator determinante para
9 uma maior variabilidade genética e amostragem da espécie (LEÓN-QUINTO et al., 2011).
10 Além disso, o estágio do ciclo celular é um dos fatores mais relevantes para que o carioplasto
11 seja reprogramado eficientemente pelo oócito receptor enucleado (CAMPBELL et al., 1996).
12 Nesse sentido, células devem ser sincronizadas em G₀/G₁ do seu ciclo celular. Contudo, a
13 resposta celular a diferentes protocolos tem sido variável em mamíferos domésticos
14 (GOISSIS et al., 2007) e silvestres (MAHESH et al., 2012), necessitando da avaliação de cada
15 protocolo para uma determinada espécie e tipo celular (WITTAYARAT et al., 2013,
16 VERAGUAS et al., 2017).

17 Já no que se refere à seleção de citoplastos aptos à TNCS, o estabelecimento de
18 protocolos de maturação *in vitro*, bem como da capacidade de ativação desses oócitos *in vitro*
19 possui também variação em cada espécie, sendo relevante o estudo desses passos em catetos.
20 Portanto, os protocolos de maturação partem principalmente da proximidade taxonômica entre
21 catetos e suínos, levando em consideração as variações de tempo utilizadas em suínos
22 (PEREYRA-BONNET et al., 2008) e suas diferentes suplementações (WU et al., 2017). Além
23 disso, diferentes protocolos de ativação já foram avaliados e variações em respostas foram
24 observadas em espécies domésticas (CHOI et al., 2013) e silvestres (LOI et al., 2011).

25 Portanto, o objetivo desta tese foi avaliar as condições de manipulação de carioplastos
26 e citoplastos, visando à TNCS em catetos. Especificamente, para atender esse objetivo foram
27 propostos: **i)** desenvolver as condições de manipulação de carioplastos pelo estabelecimento
28 de linhagens celulares, associado à definição de uma solução de criopreservação dessas
29 células e de um método de sincronização do ciclo celular em G₀/G₁ e; **ii)** avaliar diferentes
30 condições de maturação *in vitro* e ativação artificial de oócitos.

31
32
33
34

2 FUNDAMENTAÇÃO TEÓRICA

2.1 Aspectos ecológicos e quantitativo populacional dos catetos

Os catetos (*Pecari tajacu*) também conhecidos como caititu ou porco-do-mato são pertencentes à família Tayassuidae, da ordem Artiodactyla, da subordem Suiformes e da superfamília Suoidea (DESBIEZ et al., 2012). De acordo com estudos genéticos, os catetos podem conter pelo menos dois grandes clados, ou seja, um grupo de organismos originados de um único ancestral comum exclusivo, incluindo espécimes das regiões da América do Norte, Central e do Sul (GONGORA et al., 2006; 2011) com diferenças estruturais cromossômicas, mas com semelhanças fenotípicas (ADEGA et al., 2006; GONGORA et al., 2005).

Os catetos são mamíferos com 17–35 kg, de hábito diurno-crepuscular (EMMONS; FEER, 1997) que possuem hábitos alimentares herbívoros o que proporcionam a eles uma importância ecológica como dispersores de sementes (DESBIEZ et al., 2011). Em geral, os catetos vivem em grupos de 5–30 indivíduos entre machos e fêmeas de várias faixas etárias, caracterizando-se como indivíduos altamente sociáveis e cooperativos (GONGORRA et al., 2011; BIONDO et al., 2014; KEUROGHLIAN et al., 2004). Em um estudo desenvolvido por Byers; Bekoff (1981), os autores observaram um comportamento de cooperação quando vários catetos se alimentaram próximos um do outro, consumindo a mesma planta, sendo observado ainda uma relação de parentesco com alta relação genética e filopatria feminina (BYERS; BEKOFF, 1981). Adicionalmente, a estrutura do rebanho é organizada com os mais jovens centralizados e os adultos, e principalmente fêmeas, em posições mais periféricas, tanto para a remoção de alimentos quanto para proteção de agressões ao grupo (BIONDO et al., 2014).

Quanto à distribuição populacional de catetos, segundo dados da Global Biodiversity Information Facility (GBIF, 2019), há 11.131 ocorrências de indivíduos, registradas por diferentes fontes entre os anos de 1800 a 2020, como observação não especificada (106; 1,0%), observação por máquina (433; 3,9%), amostra de material (14; 0,1%), espécime preservado (3.632; 32,6%), observação humana (6.489; 58,3%), espécime fóssil (21; 0,2%) e fontes desconhecidas (436; 3,9%). Dessas ocorrências, quatorze subespécies de catetos foram observadas (**Quadro 1**), tendo ainda mais quatro espécimes registradas a serem classificadas ou não como subespécies (GBIF, 2019). Nessas ocorrências, um alto número de locais foi identificado, sendo observada uma dispersão dos catetos nos últimos anos pela América Latina, com grande concentração no México. Assim, os catetos podem ser encontrados em

diferentes habitats em virtude de sua resistência a ações antrópicas, sendo encontrados em florestas tropicais úmidas, regiões semiáridas, áreas desertas, e em áreas de temperaturas noturnas menores que 0 °C (GONGORRA et al., 2011).

Quadro 1. Ocorrência e distribuição de subespécies de catetos.

Subespécies de catetos	Nº de ocorrências	Países
<i>angulatus</i> (Cope, 1889)	232	EUA, México, Bolívia, Guatemala
<i>bangsi</i> Goldman, 1917	45	Panamá, Colômbia
<i>crassus</i> (Merriam, 1901)	42	México, Guatemala
<i>crusnigrum</i> (Bangs, 1902)	90	Costa Rica, Nicarágua, Panamá
<i>humeralis</i> (Merriam, 1901)	72	México
<i>nanus</i> (Merriam, 1901)	135	México
<i>nelsoni</i> Goldman, 1926	154	México, Guatemala, Belize,
<i>niger</i> (Allen, 1913)	10	Equador
<i>nigrescens</i> Goldman, 1926	9	Honduras, Nicarágua, Guatemala
<i>patira</i> (Kerr, 1792)	41	Venezuela, Suriname, Guiana, Brasil
<i>sonoriensis</i> (Mearns, 1897)	189	México, EUA
<i>tajacu</i> (Linnaeus, 1758)	110	Brasil, Colômbia, Venezuela, Peru, Argentina
<i>torvus</i> (Bangs, 1898)	86	Colômbia, Venezuela, Trindade Tobago
<i>yucatanensis</i> (Merriam, 1901)	122	México, Guatemala

Fonte: Global Biodiversity Information Facility (GBIF, 2019).

Já quanto à distribuição populacional de catetos, segundo a União Internacional para Conservação da Natureza (IUCN, GONGORRA et al., 2011), a espécie é classificada como menos preocupante e com distribuição em vários habitats. Por outro lado, em um estudo apresentado por Desbiez et al. (2011), os autores detalharam a situação da espécie em diferentes biomas brasileiros, classificando os catetos como menos preocupantes, mas também considerando os indivíduos como quase ameaçados na Mata Atlântica em decorrência da fragmentação e perda da qualidade de habitat bem como da caça predatória. Tal detalhamento traz a perspectiva de um possível declínio, possivelmente alterando a classificação da espécie para vulnerável à extinção (DESBIEZ et al., 2011).

Com relação ao Nordeste brasileiro, tem sido observada uma escassez de informações sobre a distribuição de catetos (CASSANO et al., 2017). Especificamente no Rio Grande do Norte, Marinho et al. (2019) apresentaram o primeiro registro dos espécimes de catetos em uma área de ecótono Caatinga-Mata Atlântica. Anteriormente, a espécie era apontada apenas por registros fósseis por Araújo-Júnior e Porpino (2011) e, posteriormente, em 2014, foi sugerido por Faria que a espécie havia sido extinta na região centro-sul do Rio Grande do Norte (Seridó). Adicionalmente, um registro de 2018 por Marinho et al. consideraram que os catetos poderiam ou estar extintos ou em locais intocados, pois não foi registrado nenhum cateto no inventário de amostragem de mamíferos de médio a grande porte do estado.

Apesar destes animais terem uma tolerância a habitats degradados, estes são considerados como indicadores da qualidade ambiental, o que demonstra que a ausência destes animais em paisagens alteradas é sinal de muita perturbação ambiental (DESBIEZ et al., 2011). Em nosso bioma Caatinga foram listadas as principais ameaças aos catetos que são ocasionadas por atividades como a geração de energia eólica, a produção de carvão, utilização de lenha, cultura de mamona e pinhão como fonte de biocombustíveis, ampliação de plantações de sisal, área de pastagem que também agravam o desmatamento e fragmentação do habitat, bem como, a criação extensiva de caprinos, ovinos e bovinos (DESBIEZ et al., 2011).

Portanto, diante da importância ecológica dos catetos e as inúmeras ações antropológicas que ameaçam o quantitativo populacional desta espécie, estratégias de conservação representam ferramentas essenciais para a manutenção da biodiversidade. Nesse contexto, em catetos, estratégias de conservação *in situ* e *ex situ* já foram desenvolvidas por diferentes grupos, objetivando desta forma a conservação da espécie.

2.2 Estratégias de conservação *in situ* aplicadas em catetos

A conservação *in situ* consiste em proteger toda área onde a espécie está inserida, preservando não somente a espécie em si, mas todo o ecossistema o qual a espécie pertence, permanecendo assim todo o habitat natural (MMA, 2020). Em geral, o desenvolvimento de estratégias *in situ* segue os pressupostos constituídos pela Convenção sobre Diversidade Biológica (BRASIL, 2000), no qual estabelecem as condições de áreas protegidas, regulamentações dos recursos biológicos, e uso sustentável dos recursos naturais. Assim, a conservação *in situ* tem como vantagens permitir que espécies continuem seus processos

1 evolutivos, mantendo as melhores condições para a conservação da vida silvestre (MMA,
2 2020).

3 No Brasil, o Ministério do Meio Ambiente estabeleceu as estratégias de conservação
4 *in situ* baseadas na formação de Unidades de Conservação, as quais são definidas como 10%
5 da área de cada bioma destinada para a proteção ambiental. Assim, para catetos, diferentes
6 regiões mantêm unidades de conservação com a espécie, tais como o Parque Nacional de
7 Brasília (ICMBio, 2012), Reserva Biológica do Tinguá, Rio de Janeiro (ICMBio, 2015), e
8 Reserva Biológica do Gurupi na Amazônia Maranhense (ICMBio, 2017). Contudo, somente
9 na Estação Ecológica de Pirapitinga do Bioma Cerrado e na Reserva Biológica União do
10 bioma Mata Atlântica, o cateto é descrito dentre a Lista de Espécies Ameaçadas protegidas da
11 Unidade de Conservação (ICMBio, 2020).

12 Adicionalmente, a criação comercial de animais silvestres é uma alternativa para
13 produtores rurais de forma sustentável, possibilitando o aproveitamento de áreas e igualmente
14 configurando-se como uma ferramenta ecológica de conservação desses indivíduos. Além
15 disso, o manejo dos catetos é previsto pelo Instituto Brasileiro do Meio Ambiente e dos
16 Recursos Naturais Renováveis (IBAMA) pelo Decreto nº 6.099, de 26 de abril de 2007 (nº
17 02001.005418/2007-11). Portanto, a criação em cativeiro nos locais onde esses animais são
18 encontrados em maior quantidade pode também ser usada de forma econômica e sustentável.

19 Não somente no Brasil, mas em outros países onde catetos podem ser encontrados,
20 uma série de estudos já foram desenvolvidos *in situ* (**Figura 1**) em áreas preservadas ou em
21 florestas onde estes animais possuem habitats. Adicionalmente, tem sido observado que
22 embora haja uma ampla distribuição desta espécie nas Américas, a maioria dos trabalhos tem
23 sido realizada no Brasil, ressaltando a sua distribuição em diferentes biomas.

24 Contudo, o número de trabalhos ainda se mostra bem reduzido em virtude
25 provavelmente por a conservação *in situ* ser onerosa, mediante das necessidades de manejo e
26 monitoramento constantes. Além disso, é de extrema importância destacar que a conservação
27 de uma espécie em um ou poucos locais de ocorrência não significa a conservação de toda a
28 sua variabilidade genética (MMA, 2020). Assim, apesar de todas as vantagens da conservação
29 *in situ* ela não mostra uma garantia ampla de preservar a variabilidade genética que é um dos
30 pontos que muitas vezes torna a discussão sobre métodos *ex situ* questionáveis.
31 Adicionalmente, é interessante destacar que a conservação da biodiversidade só poderá ser
32 alcançada por uma soma de estratégias atuando conjuntamente para aliar forças na
33 manutenção da biodiversidade.



Figura 1. Estratégias de conservação *in situ* desenvolvidas em catetos.

2.3 Estratégias de conservação *ex situ* aplicadas em catetos

A conservação *ex situ* consiste na manutenção de espécies fora de habitats naturais, tendo como principais objetivos a conservação do material genético por tempo indeterminado, permitindo que em um único local haja uma amostragem genética de muitas procedências e garantindo que haja a proteção de uma mesma espécie que tem uma distribuição geográfica muito ampla em um mesmo espaço (MMA, 2020). Neste intuito, a conservação *ex situ* pode ser considerada de duas maneiras: *ex situ in vivo* que se trata de medidas que visem a conservação em uma área restrita onde o animal geralmente é mantido em cativeiros ou fazendas (SONGSASEN; COMIZZOLI, 2019); e a *ex situ in vitro* que engloba biotecnologias reprodutivas ou técnicas de reprodução assistida (ANDRABI; MAXWELL, 2007).

No Brasil, estratégias de conservação *ex situ* são desenvolvidas de acordo com a Convenção sobre Diversidade Biológica (BRASIL, 2000), a qual são considerados componentes de diversidade biológica, estabelecimento de instalações adequadas, medidas para recuperação, regeneração e reintrodução de espécies, e colheita de recursos biológicos. Em catetos, estratégias tanto *in vivo* quanto *in vitro* já foram desenvolvidas, conforme descrição a seguir.

2.3.1 Estratégias de conservação *ex situ in vivo* aplicadas em catetos

A conservação *ex situ in vivo* consiste em estratégias que são realizadas em um ambiente delimitado, que não é o habitat natural da espécie, como fazendas de criação e cativeiros. Em catetos, as estratégias de conservação *ex situ in vivo* estão relacionadas, principalmente, ao desenvolvimento do conhecimento biológico acerca da espécie e em respostas fisiológicas e comportamentais (**Quadro 2**). A partir deste conhecimento tem sido possível desenvolver técnicas de manejo e biotecnologias que sejam mais adequadas para a espécie.

Assim, desde os primeiros estudos buscando conhecer os hábitos e a fisiologia em catetos (HELLDREN et al., 1985), trabalhos relacionados ao conhecimento fisiológico do macho (HELLDREN et al., 1989; FILGUEIRA et al., 2005; PEIXOTO et al., 2012a) e da fêmea (MAYOR et al., 2005; 2006; 2007; BATISTA et al., 2007; GARNERO et al., 2013; MAIA et al., 2014a,b; SILVA et al., 2016; AHUJA-AGUIRRE et al., 2017; MAYOR et al., 2019) têm sido desenvolvidos. Adicionalmente, estudos relacionados à aplicação destes

1 conhecimentos em técnicas de manejo e ferramentas biotecnológicas têm sido estabelecidos
 2 (GARBOR et al., 1997; SOUZA et al., 2009).

3

4 **Quadro 2.** Desenvolvimento de estratégias *ex situ in vivo* em catetos.


Finalidade	Autores
Resposta endócrina e metabólica à anestésico	Helldren et al. (1985)
Comparação do perfil dos níveis de testosterona, características seminais e mensuração testicular em animais de cativeiro e de vida livre	Helldren et al. (1989)
Avaliação de um adequado protocolo anestésico	Gabor et al. (1997)
Avaliação de parâmetros reprodutivos	Pinheiro et al. (2001)
Monitoramento reprodutivo por ultrassonografia e perfil hormonal em fêmeas	Mayor et al. (2005;2012)
Avaliação do primeiro estro pós parto e gestação em indivíduos da Amazônia	Mayor et al. (2006b)
Descrição das características estrais em indivíduos do leste da Amazônia	Mayor et al. (2007a,b)
Biometria e alterações histopatológicas dos testículos	Filgueira et al. (2005)
Padrões de atividades em cativeiro	Venturieri; Pendu (2006)
Patologias do sistema genital feminino em cativeiro	Batista et al. (2007)
Níveis de estresse produzido por densidade populacionais	Montes-Pérez et al. (2009)
Avaliação de protocolo anestésico para eletroejaculação	Souza et al. (2009)
Determinação do ciclo estral por aspectos colpocitológicos e clínicos	Guimarães et al. (2011)
Avaliação do regime de enriquecimento alimentar em cativeiro	Nogueira et al. (2011)
Análise da morfologia e ultrassonografia de órgãos abdominais em machos	Peixoto et al. (2012a)
Variação sazonal na disponibilidade dos frutos	Santos; Nogueira-Filho 2012
Avaliação da atividade do hormônio adrenocorticotrófico	Coradello et al. (2012)
Relação entre a biometria testicular e características seminais	Peixoto et al. (2012b)
Descrição da curva de crescimento de fêmeas criadas em cativeiro	Garnero et al. (2013)
Achados <i>post-mortem</i> em cativeiro no Nordeste do Brasil	Batista et al. (2014)

Comportamento social em cativeiro	Biondo et al. (2014)
Monitoramento do ciclo estral em cativeiro no semiárido nordestino	Maia et al. (2014a, b)
Influência de anestésicos sobre a ereção e ejaculação após eletroejaculação	Paiva et al. (2014)
Comportamento sexual no período periovulatório e início da gestação	Silva et al. (2016a)
Relação de dominância em cativeiro	Silva et al. (2016b)
Métodos não invasivos para avaliação da biometria testicular	Peixoto et al. (2016)
Perfis hormonais em diferentes estágios reprodutivos de fêmeas	Ahuja-Aguirre et al. (2017)
Monitoramento reprodutivo por análise dos metabólitos fecais	Mayor et al. (2019)
Variações climáticas sobre características reprodutivas do macho	Maia et al. (2019)
Sincronização do estro e inseminação artificial	Peixoto et al. (2019)
Habilidades anti-predatórias em cativeiro	Faria et al. (2020)


Além disso, trabalhos têm sido realizados para investigar os aspectos reprodutivos, como as relações da biometria testicular e características seminais (PEIXOTO et al., 2012b), monitoramento do estro (MAIA et al., 2014a), sincronização do estro (MAIA et al., 2014b), influência de variações climáticas sobre o macho (MAIA et al., 2019), e inseminação artificial (PEIXOTO et al., 2019). Tais estudos evidenciam a necessidade da continuidade de investigações nesta espécie.

2.3.2 Estratégias de conservação *ex situ in vitro* aplicadas em catetos

As técnicas de conservação *ex situ in vitro* consistem no uso de técnicas de criopreservação para a manutenção de recursos biológicos, bem como seu emprego em biotecnologias assistidas. Em catetos, as primeiras pesquisas foram voltadas para a caracterização fisiológica reprodutiva, tanto em machos (SONNER et al., 2004; CAMPOS et al., 2017) (**Figura 2**), quanto em fêmeas (LIMA et al., 2012; SILVA et al., 2016) (**Figura 3**). Adicionalmente, a conservação de amostras somáticas tem sido realizada com intuito de cultivar (SANTOS et al., 2016) e criopreservar amostras somáticas (BORGES et al., 2017b), visando aplicações futuras.



1 **Figura 2.** Características reprodutivas de catetos a partir de estudos *in vitro* sobre a
2 morfofisiologia do macho.



3 **Figura 3.** Evolução do conhecimento sobre aspectos reprodutivos de fêmeas de *P. tajacu*
4 investigados *in vitro*.

1 Paralelamente, estratégias reprodutivas foram empregadas nesses animais (**Quadro 3**),
2 especialmente quanto à conservação de germoplasma masculino (SILVA et al., 2019), e
3 feminino (CAMPOS et al., 2019). Interessantemente, os trabalhos evidenciaram a importância
4 de desenvolver técnicas específicas para esta espécie, uma vez que apesar de muitas vezes o
5 ponto de partida ser protocolos de suínos, devido algumas similaridades com os catetos, não
6 se pode extrapolar o conhecimento acerca dos suínos para a espécie em questão.

7 Especificamente, em amostras somáticas (**Figura 4**), a partir de 2016, Santos et al.
8 iniciaram os estudos visando o melhor conhecimento sobre o material somático de catetos, a
9 fim de conhecer as características do cultivo *in vitro* de células somáticas desta espécie.
10 Posteriormente, Borges et al. (2017a) identificaram os aspectos do tecido somático e as
11 células que o compõe, sendo possível conhecer as peculiaridades deste tecido e desenvolver
12 um protocolo de criopreservação específico. Ainda em 2017, foi estabelecido a melhor técnica
13 de criopreservação para tecidos somáticos, a qual a vitrificação em superfície sólida foi mais
14 adequada que a vitrificação direta em criotubos (BORGES et al., 2017b) e a melhor solução
15 crioprotetora, a qual consistiu de 3,0 M de etilenoglicol, 0,2 M de sacarose e 10% de soro
16 fetal bovino (BORGES et al., 2018a; BORGES et al., 2018b). Tais estudos permitiram
17 desenvolver o armazenamento adequado de tecidos somáticos da espécie. Adicionalmente, em
18 situações onde os indivíduos encontram-se distantes dos laboratórios de cultivo, técnicas de
19 refrigeração dos tecidos somáticos de catetos foram também avaliados (QUEIROZ NETA et
20 al., 2018).

Quadro 3. Estratégias *ex situ in vitro* aplicadas em catetos.

AVANÇO	AUTORES	AVANÇO	AUTORES
Conservação de germoplasma		Outras biotécnicas/ferramentas	
Influência da taxa de descongelação sobre a criopreservação de sêmen	Castelo et al. (2010b)	Análise evolutiva por relações genéticas e moleculares	Theimer; Keim (1994); Gongora; Morgan (2005); Gongora et al. (2006); Adegá et al. (2006); Lee et al. (2017)
Efeito da centrifugação e suplementação com sacarídeos sobre a criopreservação de sêmen	Castelo et al. (2010a)	Análise da origem do plexo braquial	Moura et al. (2007)
Conservação de sêmen a 17 °C	Garcia et al. (2012)	Avaliação de protozoários gastrointestinais	Farret et al. (2010)
Uso da água de coco em pó na criopreservação de sêmen	Silva et al. (2012)	Avaliação da variabilidade genética	Silva et al. (2010)
Avaliação de diferentes concentrações de glicerol e gema de ovo na criopreservação de sêmen	Alves et al. (2013)	Aspectos hematológicos	Almeida et al. (2011)
Avaliação de sêmen usando diferentes curvas de congelamento, tamanhos de palheta e taxas de descongelação	Silva et al. (2013)	Ensaio metabólico por glicocorticoides fecais	Coradello et al. (2012)
Conservação por curtos períodos de folículos ovarianos pré-antrais	Lima et al. (2014)	Diferenciação de células-tronco mesenquimais de tecido adiposo	Pessoa et al. (2014)

Uso de diferentes concentrações de <i>Aloe Vera</i> na criopreservação de sêmen	Souza et al. (2016)	Descrição histológica da glândula dorsal	Morales et al. (2015)
Vitrificação de tecido ovariano utilizando Ovarian Tissue Cryosystem (OTC)	Campos et al. (2019)	Diferenciação de células-tronco mesenquimais do sangue	Argôlo-Neto et al. (2016)
Vitrificação de tecido ovariano usando diferentes crioprotetores	Lima et al. (2019)	Morfologia e função da retina	Ezra-Elia et al. (2018)
Vitrificação de tecido testicular usando diferentes crioprotetores	Silva et al. (2019)	Suscetibilidade ao vírus da síndrome reprodutiva e respiratória porcina (PRRSV)	Molina-Barrios et al. (2018)




Figura 4. Caminho percorrido no estabelecimento de amostras somáticas de catetos.

Diante deste contexto, é possível entender que ainda há lacunas sobre a pesquisa básica na espécie, bem como a necessidade de novos estudos no desenvolvimento das biotecnologias a serem continuadas. Assim, avanços importantes têm norteados os próximos passos na conservação desta espécie por diferentes grupos de pesquisa.

3 JUSTIFICATIVA

A alta degradação ambiental, especialmente do bioma Caatinga, junto com a redução de mamíferos silvestres que compõem este ambiente, como os catetos, torna urgente o desenvolvimento de estratégias de conservação que possam ser aplicadas a esta espécie. Diante do potencial da transferência nuclear de célula somática (TNCS) na manutenção da biodiversidade, o esclarecimento dos primeiros passos desta técnica em catetos são fundamentais para a aplicação da TNCS visando sua conservação e criação sustentável.

Assim, é essencial o isolamento, caracterização e criopreservação de fibroblastos de catetos que possam ser utilizados como células doadoras de núcleos (carioplastos). Além disso, um protocolo mais efetivo para sincronização do ciclo celular G_0/G_1 garante que a reprogramação celular ocorra com sucesso durante a TNCS. Ainda, para que haja o sucesso na clonagem é necessário que o gameta feminino utilizado como célula receptora do núcleo (citoplasto) esteja preparado para orquestrar a reprogramação embrionária. Partindo desta necessidade, o ambiente de cultivo de oócitos deve garantir a maturação ideal para alcançar o seu potencial meiótico. Subsequentemente, o protocolo de ativação oocitária artificial necessita ser eficiente para proporcionar o desenvolvimento do embrião reconstruído.

Com base na importância destas etapas, este trabalho se preocupou em estabelecer o passo a passo do desenvolvimento inicial da técnica de TNCS, tanto dos carioplastos quanto dos citoplastos.

4 HIPÓTESES CIENTÍFICAS

4.1 Fibroblastos derivados da pele de catetos resultam em linhagens celulares com padrões proliferativos e metabólicos não alterados pelo tempo de cultivo e pela criopreservação;

4.2 A adição de altas concentrações de soro fetal bovino (SFB) na solução de congelamento lenta promovem uma melhor conservação de células somáticas de catetos após a descongelação;

4.3 A inibição por contato, independente da duração, promove a sincronização das células somáticas de catetos no estágio de G_0/G_1 do ciclo celular, não afetando ainda a viabilidade destas células;

4.4 Oócitos de catetos necessitam de um tempo de 48 h de maturação *in vitro* para o completo desenvolvimento meiótico, apresentando similaridade com oócitos suínos;

4.5 Oócitos de catetos aumentam sua habilidade meiótica quando maturados na presença de fator de crescimento epidermal (EGF);

4.6 A habilidade de desenvolvimento embrionário de oócitos de catetos após a ativação artificial é dependente do tipo de ativador secundário.

5 OBJETIVOS

5.1 Objetivo Geral

Definir as condições de manipulação de carioplastos e citoplastos, visando à transferência nuclear de célula somática (TNCS) em catetos.

5.2 Objetivos Específicos

5.2.1 Estabelecer linhagens celulares de catetos a partir de ensaios que garantam a qualidade das células para uso em TNCS;

5.2.2 Avaliar a criotolerância de células somáticas de catetos usando um agente intracelular [dimetilsulfóxido (DMSO)] em associação a diferentes crioprotetores extracelulares (sacarose e soro fetal bovino) e suas concentrações;

5.2.3 Comparar a eficiência da sincronização nas fases G₀/G₁ e viabilidade de fibroblastos de catetos, usando as estratégias: inibição por contato *vs.* privação do soro *vs.* agentes químicos [DMSO, cicloheximida (CHX), citocalasina B (CB) ou 6-dimetilaminopurina (6D)] a diferentes tempos de incubação;

5.2.4 Investigar a relação entre o potencial meiótico e os períodos (24 h *vs.* 48 h) de maturação *in vitro* de oócitos derivados de catetos;

5.2.5 Identificar a influência do fator de crescimento epidermal (EGF) no meio de maturação *in vitro* de oócitos de catetos;

5.2.6 Avaliar a ativação artificial de oócitos de catetos usando agentes químicos como ativador primário (ionomicina) e secundários (6D *vs.* 6D + CB *vs.* CHX *vs.* CHX + CB) sobre o desenvolvimento embrionário.

REFERÊNCIAS

- 1
- 2
- 3 ADEGA, F.; CHAVES, R.; KOFLER, A.; KRAUSMAN, P.R.; MASABANDA, J.;
- 4 WIENBERG, J.; GUEDES-PINTO, H. High-resolution comparative chromosome painting in
- 5 the Arizona collared peccary (*Pecari tajacu*, Tayassuidae): a comparison with the karyotype
- 6 of pig and sheep. **Chromosome Research**, v. 14, p. 243–251, 2006.
- 7 AHUJA-AGUIRRE, C.; LÓPEZ-DEBUEN, L.; ROJAS-MAYA, S.; HERNÁNDEZ-CRUZ,
- 8 B.C. Progesterone and estradiol profiles in different reproductive stages of captive collared
- 9 peccary (*Pecari tajacu*) females assessed by fecal metabolites. **Animal Reproduction**
- 10 **Science**, v. 180, p. 121–126, 2017.
- 11 ALMEIDA A.M.B.; NOGUEIRA-FILHO S.L.G.; NOGUEIRA S.S.C.; MUNHOZ A.D.
- 12 Aspectos hematológicos de catetos (*Tayassu tajacu*) mantidos em cativeiro. **Pesquisa**
- 13 **Veterinária Brasileira**, v. 31, n. 2, p. 173–177, 2011.
- 14 ALVES, H.M., OLIVEIRA, I.R.S., CASTELO, T.S., LIMA, G.L., SOUZA, A.L.P.,
- 15 MOREIRA, M.A.P., PAULA, V.V.; SILVA, A.R. Comparison of different glycerol and egg
- 16 yolk concentrations added to tris-based extender for the collared peccaries (*Tayassu tajacu*)
- 17 semen freezing. **Reproduction in Domestic Animals**, v. 48, n. 3, p. 506–511, 2013.
- 18 ANDRABI, S.M.H.; MAXWELL, W.M.C. A review on reproductive biotechnologies for
- 19 conservation of endangered mammalian species. **Animal Reproduction Science**, v. 99, p.
- 20 223–243, 2007.
- 21 ARAÚJO-JÚNIOR, H.I.; PORPINO, K.O. Assembleias fossilíferas de mamíferos do
- 22 Quaternário do Estado do Rio Grande do Norte, Nordeste do Brasil: diversidade e aspectos
- 23 tafonômicos e paleoecológicos. **Pesquisas em Geociências**, v. 38, p. 67–83, 2011.
- 24 ARGÔLO-NETO, N.M.A.; FEITOSA, M.L.T.; SOUSA, S.S.; FERNANDES, P.B.; PESSOA,
- 25 G.T.; BEZERRA, D.O.; ALMEIDA, H.M.; CARVALHO, Y.K.P.; ROCHA, A.R.; SILVA,
- 26 L.M.C.; CARVALHO, M.A.M. Isolation, expansion, differentiation and growth kinetics
- 27 essay in mesenchymal stem cells culture from the bone marrow of collared peccaries (*Tayassu*
- 28 *tajacu*). **Acta Scientiae Veterinariae**, v. 44, p. 1–11, 2016.
- 29 BATISTA, J.S.; OLINDA, R.G.; RODRIGUES, C.M.; SILVA, T.M.; VALE, R.G.; VIANA,
- 30 G.A.; OLIVEIRA, A.F.; OLIVEIRA, M. F. *Postmortem* findings in collared peccaries raised
- 31 in captivity in Northeastern Brazil. **Pesquisa Veterinária Brasileira**, v. 34, p. 1101–1108,
- 32 2014

- 1 BATISTA, J.S.; OLIVEIRA, A.F.; PORTELA, M.; BARRETO, V. Patologias do sistema
2 genital feminino de catetos (*Tayassu tajacu*) criados em cativeiro. **Revista Caatinga**, v. 20, p.
3 133–136, 2007.
- 4 BIONDO, C.; IZAR, P.; MIYAKI, C. Y.; BUSSAB, V.S. Social structure of collared peccaries
5 (*Pecari tajacu*): Does relatedness matter? **Behavioural Processes**, v. 109, p. 70–78, 2014.
- 6 BODMER, R.E.; BENDAYAN, N.Y.; MOYA, L.; FANG, T.G. Manejo de ungulados en la
7 Amazonia Peruana: Analisis de su caza y comercializacion. **Boletín de Lima**, v. 70, p. 49–
8 56, 1990.
- 9 BORGES, A.A.; LIRA, G.P.O.; NASCIMENTO, L.E.; QUEIROZ NETA, L.B.; SANTOS,
10 M.V.; OLIVEIRA; M.F., SILVA, A.R.; PEREIRA, A.F. Influence of cryopreservation
11 solution on the *in vitro* culture of skin tissues derived from collared peccary (*Pecari tajacu*
12 Linnaeus, 1758). **Biopreservation and Biobanking**, v. 16, p. 77–81, 2018a.
- 13 BORGES, A.A.; BEZERRA, F.V.; COSTA, F.N.; QUEIROZ NETA, L.B.; SANTOS, M.V.O.;
14 OLIVEIRA, M.F.; SILVA, A.R.; PEREIRA, A.F. [Histomorphological characterization of
15 collared peccary (*Pecari tajacu* Linnaeus, 1758) ear integumentary system (*Pecari tajacu*
16 Linnaeus, 1758). **Arquivo Brasileiro de Medicina Veterinária e Zootecnia**, v. 69, p. 948–
17 954, 2017a.
- 18 BORGES, A.A.; LIMA, G.L.; QUEIROZ NETA, L.B.; SANTOS, M.V.O.; OLIVEIRA, M.F.;
19 SILVA, A.R.; PEREIRA, A.F. Conservation of somatic tissue derived from collared peccaries
20 (*Pecari tajacu* Linnaeus, 1758) using direct or solid–surface vitrification techniques.
21 **Cytotechnology**, v. 69, p. 643–654, 2017b.
- 22 BORGES, A.A.; QUEIROZ NETA, L.B.; SANTOS, M.V.; OLIVEIRA, M.F.; SILVA, A.R.;
23 PEREIRA, A.F. Combination of ethylene glycol with sucrose increases survival rate after
24 vitrification of somatic tissue of collared peccaries (*Pecari tajacu* Linnaeus, 1758). **Pesquisa**
25 **Veterinária Brasileira**, v. 38, p. 350–356, 2018b.
- 26 BYERS, J.A.; BEKOFF, M. Social, spacing, and cooperative behavior of the collared peccary,
27 *Tayassu-tajacu*. **Journal Mammal**, v.62, p. 767–785, 1981.
- 28 CAMPBELL, K.H.; LOI, P.; OTAEGUI, P.J.; WILMUT, I. Cell cycle co-ordination in embryo
29 cloning by nuclear transfer. **Reviews of Reproduction**, v. 1, p. 40–46, 1996.
- 30 CAMPOS, L.B.; SILVA, A.M.; PRAXEDES, E.C.G.; BEZERRA, L.G.P.; LINS, T.L.B.G.;
31 MENEZES, V.G.; MATOS, M.H.T.; SILVA, A.R. Vitrification of collared peccary ovarian
32 tissue using open or closed systems and different intracellular cryoprotectants. **Cryobiology**,
33 v. 91, p. 77–83, 2019.

- 1 CAMPOS, L.B.; SILVA, A.M.; PEIXOTO, G.C.X.; SOUZA, A.L.P.; CASTELO, T.S.; MAIA,
2 K.M.; PEREIRA, A.F.; SILVA, A.R. Estimating the binding ability of collared peccary
3 (*Pecari tajacu* Linnaeus, 1758) sperm using heterologous substrates. **Theriogenology**, v. 92,
4 p. 57–62, 2017.
- 5 CAMPOS–JÚNIOR, P.H.A.; COSTA, G.M.J.; AVELAR, G.F.; LACERDA, S.M.S.N.;
6 COSTA, N.N.; OHASHI, O.M.; MIRANDA, M.S.; BARCELOS, L.S.; JORGE, E.C.;
7 GUIMARÃES, D.A.; FRANÇA, L.R. Derivation of sperm from xenografted testis cells and
8 tissues of the peccary (*Tayassu tajacu*). **Reproduction**, v. 147, p. 291–299, 2014.
- 9 CASSANO, C.R.; ALMEIDA–ROCHA, J.M.; ALVAREZ, M.R.; BERNARDO, C.S.S.;
10 BIANCONI, G.V.; CAMPIOLO, S.; CAMPOS, C.B.; DANILEWICZ, D.; FALCÃO, F.C.;
11 GARCÍA, F.V.; GINÉ, G.A.F.; GUIDORIZZI, C.E.; JERUSALINSKY, L.; PENDU, Y.L.;
12 MARCONDES, M.C.C.; MELO, V.L.; MENDES, F.R.; MIRETZKI, M.; NEVES, L.G.;
13 OLIVEIRA, L.C.; PEREIRA, A.R.; SILVA, K.F.M.; REIS, M.S.; VLEESCHOUWER, K.M.;
14 VIEIRA, E.M.; IACK–XIMENES, G.E. Primeira avaliação do status de conservação dos
15 mamíferos do estado da Bahia, Brasil. **Oecologia Australis** v. 21, p. 156–170, 2017.
- 16 CASTELO, T.S.; BEZERRA, F.S.B.; LIMA, G.L.; ALVES, H.M.; OLIVEIRA, I.R.S.;
17 SANTOS, E.A.A.; PEIXOTO, G.C.X.; SILVA, A.R. Effect of centrifugation and sugar
18 supplementation on the semen cryopreservation of captive collared peccaries (*Tayassu*
19 *tajacu*). **Cryobiology**, p. 275–279, 2010a.
- 20 CASTELO, T.S.; BEZERRA, F.S.B.; SOUZA, A.L.P.; MOREIRA, M.A.P.; PAULA, V.V.;
21 OLIVEIRA, M.F.; SILVA, A.R. Influence of the thawing rate on the cryopreservation of
22 semen from collared peccaries (*Tayassu tajacu*) using Tris–based extenders. **Theriogenology**,
23 v. 74, p. 1060–1065, 2010b.
- 24 CHOI, I.; ZHU, J.; CAMPBELL, K.H. The combined treatment of calcium ionophore with
25 strontium improves the quality of ovine SCNT embryo development. **Zygote**, v. 21, p. 139–
26 150, 2013.
- 27 CORADELLO, M.A.; MORAIS, R.N.; ROPER, J.; SPERCOSKI, K.M.; MASSUDA, T.;
28 NOGUEIRA, S.S.C.; NOGUEIRA–FILHO, S.L.G. Validation of a fecal glucocorticoid
29 metabolite assay for collared peccaries (*Pecari tajacu*). **Journal of Zoo and Wildlife**
30 **Medicine**, v. 43, p. 275–282, 2012.
- 31 COSTA, D.S.; HENRY, M.R.G.M.; PAULA, T.A.R. Espermatogênese de catetos (*Tayassu*
32 *tajacu*). **Arquivo Brasileiro de Medicina Veterinária e Zootecnia**. v. 56, p. 46–51, 2004.
- 33 COSTA, D.S.; PAULA, T.A.R. Cinética da espermatogênese de catetos (*Tayassu tajacu*).
34 **Revista Ceres**, v. 53, p. 512–522, 2006.

- 1 COSTA, D.S.; SILVA, J.F.S.; DA SILVEIRA, L.S. Morphometry of Leydig cells in the
2 collared peccary (*Tayassu tajacu*). **Brazilian Journal of Veterinary Research and Animal**
3 **Science**, v. 44, p. 384–389, 2007.
- 4 COSTA, G.M.; LEAL, M.C.; SILVA, J.V.; CÁSSIA, A.; FERREIRA, S.; GUIMARÃES, D.A.;
5 FRANÇA, L.R. Spermatogenic cycle length and sperm production in a fetal pig species
6 (collared peccary, *Tayassu tajacu*). **Journal of Andrology**, v. 31, p. 221–230, 2010.
- 7 DESBIEZ, A.L.J.; KEUROGHLIAN, A.; DE MELLO BEISIEGEL, B.; MEDICI, E.P.;
8 GATTI, A.; PONTES, A.R.M.; DE ALMEIDA, L.B. Avaliação do risco de extinção do cateto
9 *Pecari tajacu* Linnaeus, 1758, no Brasil. **Biodiversidade Brasileira**, v. 1, p. 74–83, 2012.
- 10 EZRA-ELIA, R.; ROSS, M.; AVNI-MAGEN, N.; BERKOWITZ, A.; OFRI, R. The retina of
11 the collared peccary (*Pecari tajacu*): structure and function. **Veterinary Ophthalmology**, v.
12 21, p. 577–585, 2018.
- 13 FARIA, O.L. A Caça nos Sertões do Seridó. In: **Sebo Vermelho**, p. 75, 2014.
- 14 FARIA, C.M.; SOUZA S.F.; COSTA, D.D.L.; SILVA, M.M.; SILVA, B.C.; YOUNG, R.J.;
15 AZEVEDO, C.S. Captive-born collared peccaries learning about their predators: Lessons
16 learnt but not remembered. **Behavioural Processes**, v. 171, p. 104031, 2020.
- 17 FARRET, M.H.; ROSA, F.V.; SILVA, A.S.; MONTEIRO, S.G. Gastrointestinal protozoa in
18 *Tayassu pecari* kept in captivity in Brazil. **Semina: Ciências Agrárias**, v. 31, p. 1041–1044,
19 2010.
- 20 FILGUEIRA, K.D.; MOURA, C.E.B.; BATISTA, J.S.; SILVA, S.M.M.S.; OLIVEIRA, M.F.;
21 ALBUQUERQUE, J.F.G.; MIGLINO, M.A. Biometria e alterações histopatológicas em
22 testículos de catetos (*Tayassu tajacu*) criados em cativeiro no semiárido nordestino, Brasil.
23 **Brazilian Journal of Veterinary Research and Animal Science**, v. 42, p. 19–25, 2005.
- 24 FOLCH, J.; COCERO, M.J.; CHESNÉ, P.; ALABART, J.L.; DOMÍNGUEZ, V.; COGNIÉ, Y.;
25 VIGNON, X. First birth of an animal from an extinct subspecies (*Capra pyrenaica pyrenaica*)
26 by cloning. **Theriogenology**, v. 71, p. 1026–1034, 2009.
- 27 GABOR, T.M.; HELLGREN, E.C.; SILVY, N.J. Immobilization of collared peccaries
28 (*Tayassu tajacu*) and feral hogs (*Sus scrofa*) with Telazol® and xylazine. **Journal of Wildlife**
29 **Diseases**, v. 33, p. 161–164, 1997.
- 30 GARCIA, A.R.; KAHWAGE, P.R.; GUIMARÃES, D.A.A.; OHASHI, O.M. Chilled semen of
31 captive collared peccaries (*Pecari tajacu*): Effects of preservation at 17 °C on semen quality.
32 **Journal of Agricultural Science and Technology**, v. 2, p. 921–929, 2012.
- 33 GARCÍA-MENGUAL, E.; ALFONSO, J.; SALVADOR, I.; DUQUE, C.C.; SILVESTRE,
34 M.A. Oocyte activation procedures and influence of serum on porcine oocyte maturation and

- 1 subsequent parthenogenetic and nuclear transfer embryo development. **Zygote**, v. 16, p. 279–
- 2 284, 2008.
- 3 GARNERO, A.D.V.; MARCONDES, C.R.; ALBUQUERQUE, N.I.; ARAÚJO, R.O.; PENDU,
- 4 Y.; GUIMARÃES, D.A. Growth curve of female collared peccaries (*Pecari tajacu*) raised in
- 5 captivity in the Brazilian Amazon Region. **Arquivo Brasileiro de Medicina Veterinária e**
- 6 **Zootecnia**, v. 65, p. 961–966, 2013.
- 7 Global Biodiversity Information Facility. *Pecari tajacu* (Linnaeus, 1758). **GBIF** 2019.
- 8 Disponível em: <https://doi.org/10.15468/39omei> accessed via GBIF.org on 2020–04–02.
- 9 Acesso em 31 de março de 2020.
- 10 GOISSIS, M.D.; CAETANO, H.V.A.; MARQUES, M.G.; BARROS, F.R.O.; FEITOSA, W.B.;
- 11 MILAZZOTTO, M.P.; BINELLI, M.; ASSUMPÇÃO, M.E.O.A.; VISINTIN, J.A. Effects of
- 12 serum deprivation and cycloheximide on cell cycle of low and high passage porcine fetal
- 13 fibroblasts. **Reproduction in Domestic Animals**, v. 42, p. 660–663, 2007.
- 14 GOMES, H.A.; CAMPOS, L.B.; PRAXEDES, E.C.; OLIVEIRA, M.F.; PEREIRA, A.F.;
- 15 SILVA, A.R.; SARAIVA, M.V. BMP-15 activity on in vitro development of collared peccary
- 16 (*Pecari tajacu* Linnaeus, 1758) preantral follicles. **Reproduction in Domestic Animals**,
- 17 2020.
- 18 GONÇALVES, A.M.M.; BORELLI, V. Histologia do funículo espermático do cateto (*Tayassu*
- 19 *tajacu*) Histology of the spermatic cord of boar (*Tayassu tajacu*). **Revista do Instituto de**
- 20 **Ciências da Saúde**, v. 26, p. 210–214, 2008.
- 21 GONGORA, J.; MORALES, S.; BERNAL, J.E.; MORAN, C. Phylogenetic divisions among
- 22 Collared peccaries (*Pecari tajacu*) detected using mitochondrial and nuclear sequences.
- 23 **Molecular Phylogenetics and Evolution**, v. 41, p. 1–11, 2006.
- 24 GONGORA, J.; MORGAN, C. Nuclear and mitochondrial evolutionary analyses of Collared,
- 25 White-lipped, and Chacoan peccaries (*Tayassuidae*). **Molecular Phylogenetics and**
- 26 **Evolution**, v. 34, p. 181–189, 2005.
- 27 GONGORA, J.; REYNA-HURTADO, R.; BECK, H.; TABER, A.; ALTRICHTER, M.;
- 28 KEUROGHLIAN, A. *Pecari tajacu*. **The IUCN Red List of Threatened Species** 2011:
- 29 e.T41777A10562361, 2011.
- 30 GUIMARÃES, D.A.; CARDOSO, D.D.L.; FERREIRA, M.A.P.; ALBUQUERQUE, N.I.D.
- 31 Puberty in male collared peccary (*Pecari tajacu*) determined by quantitative analysis of
- 32 spermatogenic cells. **Acta Amazonica**, v. 43, p. 99–103, 2013.

- 1 GUIMARÃES, D.A.; GARCIA, S.C.G.; FERREIRA, M.A.P.; SILVA, S.D.S.B.;
- 2 ALBUQUERQUE, N.I.; LE PENDU, Y. Ovarian folliculogenesis in collared peccary: *Pecari*
- 3 *tajacu* (Artiodactyla: Tayassuidae). **Revista de Biología Tropical**, v. 60, p. 437–455, 2012.
- 4 GUIMARÃES, D.A.D.; GARCIA, S.C.G.D.; PENDU, Y.L.; ALBUQUERQUE, N.I.D.
- 5 Determinação do ciclo estral em catetos *Pecari tajacu*: aspectos colpocitológicos e clínicos.
- 6 **Acta Amazonica**, v. 41, p. 583–588, 2011.
- 7 HELLGREN, E.C.; LOCHMILLER, R.L.; AMOSS J.M.S.; GRANT, W.E. Endocrine and
- 8 metabolic responses of the collared peccary (*Tayassu tajacu*) to immobilization with ketamine
- 9 hydrochloride. **Journal of Wildlife Diseases**, v. 21, p. 417–425, 1985.
- 10 HELLGREN, E.C.; LOCHMILLER, R.L.; AMOSS, M.S.; SEAGER, S.W. J.; MAGYAR, S.J.;
- 11 COSCARELLI, K.P.; GRANT, W.E. Seasonal variation in serum testosterone, testicular
- 12 measurements and semen characteristics in the collared peccary (*Tayassu tajacu*).
- 13 **Reproduction**, v. 85, p. 677–686, 1989.
- 14 HUAN, Y.; HU, K.; XIE, B.; SHI, Y.; WANG, F.; ZHOU, Y.; LIU, S.; HUANG, B.; ZHU, J.;
- 15 LIU, Z.; HE, Y.; LI, Y.; KONG, Q.; LIU, Z. Ovulation statuses of surrogate gilts are
- 16 associated with the efficiency of excellent pig cloning. **PloS One**, v. 10, p. e0142549, 2015.
- 17 HUANG, Y.; YUAN, L.; LI, T.; LIU, X.; YU, Y.; OUYANG, H.; WANG, B. IWP2 impairs the
- 18 development of porcine somatic cell nuclear transfer embryos via signaling pathway
- 19 inactivation. **Biomedical Reports**, v. 7, p. 36–40, 2017.
- 20 Instituto Chico Mendes de Conservação da Biodiversidade. **17 catetos são encontrados mortos**
- 21 **no Parque de Brasília**. ICMBio 2012. Disponível em:
- 22 [https://www.icmbio.gov.br/portal/ultimas-noticias/20-geral/3248-17-catetos-sao-](https://www.icmbio.gov.br/portal/ultimas-noticias/20-geral/3248-17-catetos-sao-encontrados-mortos-no-parque-de-brasilia)
- 23 [encontrados-mortos-no-parque-de-brasilia](https://www.icmbio.gov.br/portal/ultimas-noticias/20-geral/3248-17-catetos-sao-encontrados-mortos-no-parque-de-brasilia). Acesso em: 29 de março de 2020.
- 24 Instituto Chico Mendes de Conservação da Biodiversidade. **Esec de Pirapitinga**. ICMBio,
- 25 2020. Disponível em: <https://www.icmbio.gov.br/portal/esec-de-pirapitinga>. Acesso em: 30
- 26 de março de 2020.
- 27 Instituto Chico Mendes de Conservação da Biodiversidade. **Rebio União**. ICMBio, 2017.
- 28 Disponível em: <https://www.icmbio.gov.br/portal/rebio-uniao>. Acesso em: 29 de março de
- 29 2020.
- 30 Instituto Chico Mendes de Conservação da Biodiversidade. **Tinguá conserva pedaço de Mata**
- 31 **Atlântica**. ICMBio, 2015. Disponível em: [https://www.icmbio.gov.br/portal/ultimas-](https://www.icmbio.gov.br/portal/ultimas-noticias/4-destaques/6978-tingua-conserva-pedaco-de-mata-atlantica)
- 32 [noticias/4-destaques/6978-tingua-conserva-pedaco-de-mata-atlantica](https://www.icmbio.gov.br/portal/ultimas-noticias/4-destaques/6978-tingua-conserva-pedaco-de-mata-atlantica). Acesso em: 29 de
- 33 março de 2020.

- 1 KEUROGHLIAN, A.; EATON, D.P.; LONGLAND, W.S. Area use by white-lipped and
2 collared peccaries (*Tayassu pecari* and *Tayassu tajacu*) in a tropical forest fragment.
3 **Biological Conservation**, v. 120, p. 411–425, 2004.
- 4 KWONG, P.J.; NAM, H.Y.; KHADIJAH, W.W.; KAMARUL, T.; ABDULLAH, R.B.
5 Comparison of *in vitro* developmental competence of cloned caprine embryos using donor
6 karyoplasts from adult bone marrow mesenchymal stem cells vs ear fibroblast cells.
7 **Reproduction in Domestic Animals**, v. 49, p. 249–253, 2014.
- 8 LARA, N.L.M.; COSTA, G.M.J.; AVELAR, G.F.; GUIMARÃES, D.A.; FRANÇA, L.R.
9 Postnatal testis development in the collared peccary (*Tayassu tajacu*), with emphasis on
10 spermatogonial stem cells markers and niche. **General and Comparative Endocrinology**, v.
11 273, p. 98–107, 2019.
- 12 LEE, W.J.; LEE, J.H.; JEON, R.H.; JANG, S.J.; LEE, S.C.; PARK, J.S.; RHO, G.J. Supplement
13 of autologous ooplasm into porcine somatic cell nuclear transfer embryos does not alter
14 embryo development. **Reproduction in Domestic Animals**, v. 52, p. 437–445, 2017.
- 15 LEÓN-QUINTO, T.; SIMÓN, M.A.; SÁNCHEZ, Á.; MARTÍN, F.; SORIA, B. Cryobanking
16 the genetic diversity in the critically endangered Iberian lynx (*Lynx pardinus*) from skin
17 biopsies. Investigating the cryopreservation and culture ability of highly valuable explants and
18 cells. **Cryobiology**, v. 62, p. 145–151, 2011.
- 19 LIMA, G.L.; LUZ, V.B.; LUNARDI, F.O.; SOUZA, A.L.; PEIXOTO, G.C.; RODRIGUES,
20 A.P.R.; OLIVEIRA, M.F.; SANTOS, R.R.; SILVA, A.R. Effect of cryoprotectant type and
21 concentration on the vitrification of collared peccary (*Pecari tajacu*) ovarian tissue. **Animal**
22 **Reproduction Science**, v. 205, p. 126–133, 2019.
- 23 LIMA, G.L.; SANTOS, E.A.A.; LIMA, L.F.; LUZ, V.B.; RODRIGUES, A.P.R.; SILVA, A.R.
24 Short-term preservation of *Pecari tajacu* ovarian preantral follicles using phosphate buffered
25 saline (PBS) or powdered coconut water (ACP) media. **Arquivo Brasileiro de Medicina**
26 **Veterinária e Zootecnia**, v. 66, p. 1623–1630, 2014.
- 27 LIMA, G.L.; SANTOS, E.A.A.; LUZ, V.B.; RODRIGUES, A.P.R.; SILVA, A.R.
28 Morphological characterization of the ovarian preantral follicle population of collared
29 peccaries (*Tayassu tajacu* Linnaeus, 1758). **Anatomia, Histologia, Embryologia**, v. 42, p.
30 304–311, 2012.
- 31 LIU, Z.; CAI, Y.; LIAO, Z.; XU, Y.; WANG, Y.; WANG, Z.; JIANG, X.; LI, Y.; LU, Y.; NIE,
32 Y.; ZHANG, X.; LI, C.; BIAN, X.; POO, M.; CHANG, H.C.; SUN, Q. Cloning of a gene-
33 edited macaque monkey by somatic cell nuclear transfer. **National Science Review**, v. 6, p.
34 101–108, 2019.

- 1 LOI, P.; MODLINSKI, J.A.; PTAK, G. Interspecies somatic cell nuclear transfer: a salvage tool
2 seeking first aid. **Theriogenology**, v. 76, p. 217–228, 2011.
- 3 MAHESH, Y.U.; RAO, B.S.; KATARI, V.C.; KOMJETI, S.; CHRISTO, D.;
4 LAKSHMIKANTAN, U.; PAWAR, R.M.; SHIVAJI, S. Cell cycle synchronization of bison
5 (*Bos gaurus*) fibroblasts derived from ear piece collected *post-mortem*. **Reproduction in**
6 **Domestic Animals**, v. 47, p. 799–805, 2012.
- 7 MAIA, K.M.; PEIXOTO, G.C.; CAMPOS, L.B.; BEZERRA, J.A.B.; RICARTE, A.R.;
8 MOREIRA, N.; OLIVEIRA, M.F.; SILVA, A.R. Estrus cycle monitoring of captive collared
9 peccaries (*Pecari tajacu*) in semiarid conditions. **Pesquisa Veterinária Brasileira**, v. 34, p.
10 1115–1120, 2014a.
- 11 MAIA, K.M.; PEIXOTO, G.C.X.; CAMPOS, L.B.; SILVA, A.M.; CASTELO, T.S.;
12 RICARTE, A.R.F.; SILVA, A.R. Estrous synchronization in captive collared peccaries
13 (*Pecari tajacu*) using a prostaglandin F2 α analog. **Zoological Science**, v. 31, p. 836–839,
14 2014b.
- 15 MAIA, K.M.; SOUZA, A.L.P.; SILVA, A.M.; SOUZA, J.J.B.F.; COSTA, L.L.M.;
16 BRANDÃO, F.Z.; OLIVEIRA, M.F.; COMIZZOLI, P.; SILVA, A.R. Environmental effects
17 on collared peccaries (*Pecari tajacu*) serum testosterone, testicular morphology, and semen
18 quality in the Caatinga biome. **Theriogenology**, v. 126, p. 286–294, 2019.
- 19 MARINHO, P.H.; BEZERRA, D.; ANTONGIOVANNI, M.; FONSECA, C.R.;
20 VENTICINQUE, E.M. Mamíferos de médio e grande porte da Caatinga do Rio Grande do
21 Norte, nordeste do Brasil. **Mastozoología Neotropical**, v.25, p. 345–362, 2018.
- 22 MARINHO, P.H.; SILVA, M.; LISBOA, C.M.C.A. Presence of the collared peccary *Pecari*
23 *tajacu* (Artiodactyla, Tayassuidae) in the far northeast of its Brazilian distribution.
24 **Neotropical Biology and Conservation**, v. 14, p. 499–509, 2019.
- 25 MAYOR, P.; SILVA, G.P.; ANDRADE, R.D.S.; MONTEIRO, F.O.B.; EL BIZRI; H.R.
26 Embryonic and fetal development of the collared peccary (*Pecari tajacu*). **Animal**
27 **Reproduction Science**, v. 208, p. 106123, 2019.
- 28 MAYOR, P.; FENECH, M.; BODMER, R.E.; LOPEZ-BEJAR, M. Ovarian features of the wild
29 collared peccary (*Tayassu tajacu*) from the northeastern Peruvian Amazon. **General and**
30 **Comparative Endocrinology**, v. 147, p. 268–275, 2006a.
- 31 MAYOR, P.; GALVEZ, H.; GUIMARAES, D.A.; LOPEZ-GATIUS, F.; LOPEZ-BEJAR, M.
32 Serum estradiol-17 β , vaginal cytology and vulval appearance as predictors of estrus cyclicity
33 in the female collared peccary (*Tayassu tajacu*) from the eastern Amazon region. **Animal**
34 **Reproduction Science**, v. 97, p. 165–174, 2007a.

- 1 MAYOR, P.; GUIMARAES, D.A.; LE PENDU, Y.; SILVA, J.V.; JORI, F.; LÓPEZ-BÉJAR,
2 M. Reproductive performance of captive collared peccaries (*Tayassu tajacu*) in the eastern
3 Amazon. **Animal Reproduction Science**, v. 102, p. 88–97, 2007b.
- 4 MAYOR, P.; GUIMARÃES, D.A.; LÓPEZ-BÉJAR, M. Progesterone and estradiol-17 β as a
5 potential method for pregnancy diagnosis in the collared peccary (*Pecari tajacu*). **Research in**
6 **Veterinary Science**, v. 93, p. 1413–1417, 2012.
- 7 MAYOR, P.; GUIMARAES, D.A.; LOPEZ-GATIUS, F.; LOPEZ-BEJAR, M. First
8 *postpartum* estrus and pregnancy in the female collared peccary (*Tayassu tajacu*) from the
9 Amazon. **Theriogenology**, v. 66, p. 2001–2007, 2006b.
- 10 MAYOR, P.; LÓPEZ-GATIUS, F.; LÓPEZ-BÉJAR, M. Integrating ultrasonography within the
11 reproductive management of the collared peccary (*Tayassu tajacu*). **Theriogenology**, v. 63, p.
12 1832–1843, 2005.
- 13 MAYOR, P; JORI, F; LÓPEZ-BÉJAR, M. Anatomicohistological characteristics of the tubular
14 genital organs of the female collared peccary (*Tayassu tajacu*) from north-eastern
15 Amazon. **Anatomia, Histologia, Embryologia**, v. 33, n. 2, p. 65–74, 2004.
- 16 MEDEIROS, D.A.S.; CAMPOS, L.B.; PEREIRA, A.F.; SILVA, A.R. Recuperação oocitária
17 em catetos (*Pecari tajacu* Linnaeus, 1758) utilizando o método de fatiamento ovariano –
18 resultados preliminares. **Revista Acta Veterinaria Brasilica**, v. 8, p. 51–52, 2014.
- 19 Ministério do Meio Ambiente, **Biodiversidade**. MMA, 2020. Disponível em:
20 [https://www.mma.gov.br/biodiversidade/conservacao-e-promocao-do-uso-da-diversidade-](https://www.mma.gov.br/biodiversidade/conservacao-e-promocao-do-uso-da-diversidade-genetica/agrobiodiversidade/conserva%C3%A7%C3%A3o-in-situ,-ex-situ-e-on-farm)
21 [genetica/agrobiodiversidade/conserva%C3%A7%C3%A3o-in-situ,-ex-situ-e-on-farm](https://www.mma.gov.br/biodiversidade/conservacao-e-promocao-do-uso-da-diversidade-genetica/agrobiodiversidade/conserva%C3%A7%C3%A3o-in-situ,-ex-situ-e-on-farm).
22 Acesso em: 25 de março de 2020.
- 23 MIRANDA-MOURA, M.T.M.; OLIVEIRA, G.B.; PEIXOTO, G.C.X.; PESSOA, J.M.; PAPA,
24 P.C.; MAIA, M.S.; MOURA, C.E.B.; OLIVEIRA, M.F. Morphology and vascularization of
25 the corpus luteum of peccaries (*Pecari tajacu* Linnaeus, 1758) throughout the estrous cycle.
26 **Arquivo Brasileiro de Medicina Veterinária e Zootecnia**, v. 68, p. 87–96, 2016.
- 27 MOHAMMED, A.E.N.A.; AL-SUWAIEGH, S.; AL-SHAHEEN, T. Do the cytoplasm and
28 nuclear material of germinal vesicle oocyte support developmental competence upon
29 reconstruction with embryonic/somatic nucleus. **Cellular Reprogramming**, v. 21, p. 163–
30 170, 2019.
- 31 MOLINA-BARRIOS, R.; LUEVANO-ADAME, J.; HENAO-DÍAZ, Y.A.;
32 GIMÉNEZ-LIROLA, L.; PIÑEYRO, P.; MAGTOTO, R.; ZIMMERMAN, J. Collared
33 peccary (*Pecari tajacu*) are susceptible to porcine reproductive and respiratory syndrome
34 virus (PRRSV). **Transboundary and Emerging Diseases**, v. 65, p. 1712–1719, 2018.

- 1 MONTES-PÉREZ, R.C.; SOLÍS-SOSA, A.L.; YOKOYAMA-KANO, J.; MUKUL-YERVES,
2 J.M.; SEGURA-CORREA, J.C. Evaluación de estrés en el *Pecari tajacu* sometido a dos
3 densidades de población. **Archivos de Zootecnia**, v. 58, p. 463–466, 2009.
- 4 MOREIRA, S.S.; SILVA, A.M.; PRAXEDES, É.C.; CAMPOS, L.B.; SANTOS, C.S.; SOUZA,
5 A.L.; SILVA, A.R. Composition of collared peccary seminal plasma and sperm motility
6 kinetics in semen obtained during dry and rainy periods in a semiarid biome. **Animal**
7 **Reproduction Science**, v. 211, p. 106229, 2019.
- 8 MOURA, C.E.B.; ALBUQUERQUE, J.F.G.; MAGALHÃES, M.S.; SILVA, N.B.; OLIVEIRA,
9 M.F.; PAPA, P.C. Análise comparativa da origem do plexo branquial de catetos (*Tayassu*
10 *tajacu*). **Pesquisa Veterinária Brasileira**, v. 27, p. 357–362, 2007.
- 11 NAGY-REIS, M.B.; RIBEIRO, M.C.; SETZ, E.Z.; CHIARELLO, A.G. The key role of
12 protection status in safeguarding the ecological functions of some Neotropical
13 mammals. **Biodiversity and Conservation**, v. 28, p. 2599–2613, 2019.
- 14 NOGUEIRA, S.S.C.; NOGUEIRA-FILHO, S.L.G. Wildlife farming: an alternative to
15 unsustainable hunting and deforestation in Neotropical forests? **Biodiversity and**
16 **Conservation**, v. 20, p. 1385–1397, 2011.
- 17 NOVO, C.L.; TANG, C.; AHMED, K.; DJURIC, U.; FUSSNER, E.; MULLIN, N.P.;
18 MORGAN, N.P.; HAYRE, J.; SIENERTH, A.R.; ELDERKIN, S.; NISHINAKAMURA, R.
19 CHAMBERS, I.; ELLIS, J.; BAZETT-JONES, D.P.; RUGG-GUNN, P.J. The pluripotency
20 factor Nanog regulates pericentromeric heterochromatin organization in mouse embryonic
21 stem cells. **Genes & Development**, v. 30, p. 1101–1115, 2016.
- 22 OLIVEIRA, A.P.M.; LIMA C.D.; SARMENTO, A.C. Epididymal morphophysiology of adult
23 collared peccaries (*Pecari tajacu*). **Brazilian Journal of Veterinary Research and Animal**
24 **Science**, v. 53, p. 251–259, 2016.
- 25 PAIVA, A.L.; NUNES, T.L.; OLIVEIRA, M.G.; MORAIS, A.M.; SANTOS, E.A.; SILVA,
26 A.R.; PAULA, V.V. Effects of atipamezole and medetomidine administration on seminal
27 variables and functions of erection and ejaculation of the collared peccary (*Tayassu tajacu*)
28 after electroejaculation. **BMC Veterinary Research**, v. 10, p. 170–176, 2014.
- 29 PEIXOTO, G.C.; LIMA, G.L.; MAIA, K.M.; SOUZA, A.L.P.; CASTELO, T.S.; PAIVA, A. L.;
30 VIANA, A.C.N. Single injection of eCG/hCG leads to successful estrous synchronization in
31 the collared peccary (*Pecari tajacu* Linnaeus, 1758). **Animal Reproduction Science**, v. 208,
32 p. 106112, 2019.

- 1 PEIXOTO, G.C.X.; OLIVEIRA, I.R.S.; ALVES, N.D.; OLIVEIRA, M.F.; SILVA, A.R.
2 Abdominal Exploration in captive collared peccaries (*Tayassu tajacu*) by ultrasonography.
3 **Anatomia, Histologia, Embryologia**, v. 41, p. 256–261, 2012a.
- 4 PEIXOTO, G.C.X.; SILVA, M.A.; CASTELO, T.S.; SILVA, A.M.; BEZERRA, J.A.B.;
5 SOUZA, A.L.P.; OLIVEIRA, M.F.; SILVA, A.R. Individual variation related to testicular
6 biometry and semen characteristics in collared peccaries (*Tayassu tajacu* Linnaeus, 1758).
7 **Animal Reproduction Science**, v. 134, p. 191–196, 2012b.
- 8 PEIXOTO, G.C.X.; SILVA, M.A.; LIMA, G.L.; CAMPOS, L.B.; PAIVA, A.L.C.; FREITAS,
9 V.A.; SILVA, A.R. Use of non-invasive methods for evaluating the testicular biometry in
10 collared peccaries (*Pecari tajacu* Linnaeus, 1758). **Anatomia, Histologia, Embryologia**, v.
11 45, p. 60–66, 2016.
- 12 PESSOA, G.T.; FEITOSA, M.L.T.; NETO, N.M.A.; ROCHA, A.R.; COSTA, C.R.D.M.;
13 SILVA, G.C.; BEZERRA, D.O.; COELHO, C.J.C.; SOUSA, S.S.; CARVALHO, M.A.M.
14 Isolation, culture and differentiation potential of collared peccary (*Tayassu tajacu*) adipose–
15 derived stem cells. **Acta Scientiae Veterinariae**, v. 42, p. 1–10, 2014.
- 16 PINHEIRO, M.J.P.; SILVA, F.N.; AZEVÊDO, C.M.S.B. Avaliação de parâmetros reprodutivos
17 em catetos (*Tayassu tajacu*) criados em cativeiro. **Revista Caatinga**, v. 14, p. 71–74, 2001.
- 18 PRIYA, D.; SELOKAR, N.L.; RAJA, A.K.; SAINI, M.; SAHARE, A.A.; NALA, N.; PALTA,
19 P.; CHAUHAN, M.S.; MANIK, R.S.; SINGLA, S.K. Production of wild buffalo (*Bubalus*
20 *arnee*) embryos by interspecies somatic cell nuclear transfer using domestic buffalo (*Bubalus*
21 *bubalis*) oocytes. **Reproduction in Domestic Animals**, v. 49, p. 343–351, 2014.
- 22 QUEIROZ NETA, L.B.; OLIVEIRA LIRA, G.P.; BORGES, A.A.; OLIVEIRA SANTOS,
23 M.V.; SILVA, M.B.; OLIVEIRA, L.R.M.; SILVA, A.R.; PEREIRA, A.F. Influence of
24 storage time and nutrient medium on recovery of fibroblast-like cells from refrigerated
25 collared peccary (*Pecari tajacu* Linnaeus, 1758) skin. **In vitro Cellular & Developmental**
26 **Biology–Animal**, v. 54, p. 486–495, 2018.
- 27 SANTOS, C.A.B; NOGUEIRA–FILHO S.L.G. Disponibilidade de frutos para caititus, *Tayassu*
28 *tajacu* L. 1758 (Mammalia, Artiodactyla) na região cacauieira do Sul da Bahia. **Revista**
29 **Semiárido de Visu**, v. 2, p. 265–273, 2012.
- 30 SANTOS, E.A.A.; SOUSA, P.C.; MARTINS, J.A.M.; MOREIRA, R.A.; MONTEIRO–
31 MOREIRA, A.C.; OLIVEIRA, M.F.; MOURA, A.A.A.; SILVA, A.R. Protein profile of the
32 seminal plasma of collared peccaries (*Pecari tajacu*). **Reproduction**, v. 147, p. 753–764,
33 2014.

- 1 SANTOS, M.L.T.; BORGES, A.A.; QUEIROZ NETA, L.B.; SANTOS, M.V.O.; OLIVEIRA,
2 M.F.; SILVA, A.R.; PEREIRA, A.F. *In vitro* culture of somatic cells derived from ear tissue
3 of collared peccary (*Pecari tajacu* Linnaeus, 1758) in medium with different requirements.
4 **Pesquisa Veterinária Brasileira**, v. 36, p. 1194–1202, 2016.
- 5 SANTOS, T.C.; DANTZER, V.; JONES, C.J.P.; OLIVEIRA, M.F.; MIGLINO, M.A.
6 Macroscopic and microscopic aspects of collared peccary and white-lipped peccary placenta.
7 **Placenta**, v. 27, p. 244–257, 2006.
- 8 SANTOS, T.C.; MIGLINO, M.A.; MACHADO, G.V.; SOUZA, W.M. Vascularização arterial
9 dos ovários, tubas uterinas e útero em catetos (*Tayassu tajacu*, Linnaeus, 1758) e queixadas
10 (*Tayassu pecari*, Link, 1795). **Brazilian Journal of Veterinary Research and Animal**
11 **Science**, v. 37, p. 278–285, 2000.
- 12 SANTOS, T.C.; TONARELLI, C.; TEIXEIRA, F.A.; OLIVEIRA, M.F.; MARIA, D.;
13 REGINATO, P.; MIGLINO, M.A. Histomorphometrical and proliferative aspects of placenta
14 and uterus of the collared peccary (*Tayassu tajacu*). **Histology and Histopathology**, v. 27, p.
15 793, 2012.
- 16 SILVA, A.M.; BEZERRA, L.G.P.; PRAXEDES, E.C.G.; MOREIRA, S.S.J.; SOUZA, C.M.P.;
17 OLIVEIRA, M.F.; SILVA, A.R. Combination of intracellular cryoprotectants preserves the
18 structure and the cells proliferative capacity potential of adult collared peccary testicular
19 tissue subjected to solid surface vitrification. **Cryobiology**, v. 91, p. 53–60, 2019.
- 20 SILVA, A.M.; PEIXOTO, G.C.X.; BEZERRA, J.A.B.; CASTELO, T.S.; SANTOS, E.A.A.;
21 SILVA, A.R. Relações entre a câmara de Neubauer a espectrofotometria utilizada para a
22 determinação da concentração espermática de catetos (*Pecari tajacu*). **Ciência Rural**, v. 44,
23 p. 1494–1498, 2014.
- 24 SILVA, M.A.; PEIXOTO, G.C.X.; CASTELO, T.S.; LIMA, G.L.; SILVA, A.M.; OLIVEIRA,
25 M.F.; SILVA, A.R. Cryopreservation of collared peccary (*Peccary tajacu*) semen using
26 different freezing curves, straw sizes, and thawing rates. **Cryobiology**, v. 67, p. 50–55, 2013.
- 27 SILVA, M.A.; PEIXOTO, G.C.X.; LIMA, G.L.; BEZERRA, J.A.B.; CAMPOS, L.B.; PAIVA,
28 A.L.C.; PAULA, V.V.; SILVA, A.R. Cryopreservation of collared peccaries (*Tayassu tajacu*)
29 semen using a powdered coconut water (ACP–116c) based extender plus various
30 concentrations of egg yolk and glycerol. **Theriogenology**, v. 78, p. 605–611, 2012.
- 31 SILVA, R.W.; FREITAS, T.R.O.; SBALQUEIRO, I.J. Evaluation of genetic variability in the
32 collared peccary *Pecari tajacu* and the white-lipped peccary *Tayassu pecari* by microsatellite
33 markers. **Genetics and Molecular Biology**, v. 33, p. 62–67, 2010.

- 1 SILVA, S.D.S.B.; GUIMARÃES, D.A.; BIONDO, C.; OHASHI, O.M.; ALBUQUERQUE,
2 N.I.; DALLA VECCHIA, A.C.; LE PENDU, Y. Dominance relationships between collared
3 peccaries *Pecari tajacu* (Cetartiodactyla: Tayassuidae) in intensive breeding system. **Applied**
4 **Animal Behaviour Science**, v. 184, p. 117–125, 2016a.
- 5 SILVA, S.D.S.B.; LE PENDU, Y.; OHASHI, O.M.; OBA, E.; ALBUQUERQUE, N.I.;
6 GARCIA, A.R.; MAYOR, P.; GUIMARÃES, D.A.A. Sexual behavior of *Pecari tajacu*
7 (Cetartiodactyla: Tayassuidae) during periovulatory and early gestation periods. **Behavioural**
8 **Processes**, v. 131, p. 68–73, 2016b.
- 9 SONG, J.S.; HUA, S.; ZHANG, Y. Culture, characteristics, and chromosome complement of
10 Siberian tiger fibroblasts for nuclear transfer. **In vitro Cell Development and Biology**, v. 43,
11 p. 203–209, 2007.
- 12 SONGSASEN, N.; COMIZZOLI, P. Protecting and extending fertility for females of wild and
13 endangered mammals. In: **Textbook of Oncofertility Research and Practice**. Springer,
14 Cham, 2019. p. 401–412.
- 15 SONNER, J.B.; MIGLINO, M.A.; SANTOS, T.C.D.; CARVALHAL, R.; NETO, A.;
16 MOURA, C.E.B.D.; OLIVEIRA, M.F.D. Aspectos macroscópicos e morfométricos dos
17 testículos em catetos e queixadas. **Biota Neotropica**, v.4, p. 1–13, 2004.
- 18 SOUSA, P.C.; SANTOS, E.A.A.; SOUZA, A.L.P.; LIMA, G.L.; BARROS, F.F.P.C.;
19 OLIVEIRA, M.F.; SILVA, A.R. Sperm morphological and morphometric evaluation in
20 captive collared peccaries (*Pecari tajacu*). **Pesquisa Veterinária Brasileira**, v. 33, p. 924–
21 930, 2013.
- 22 SOUZA, A.L.P.; CASTELO, T.S.; QUEIROZ, J.P.A.F.; BARROS, I.O.; PAULA, V.V.;
23 OLIVEIRA, M.F.; SILVA, A.R. Evaluation of anesthetic protocol for the collection of semen
24 from captive collared peccaries (*Tayassu tajacu*) by electroejaculation. **Animal**
25 **Reproduction Science**, v. 116, p. 370–375, 2009.
- 26 SOUZA, A.L.P.; LIMA, G.L.; PEIXOTO, G.C.X.; SILVA, A.M.; OLIVEIRA, M.F.; SILVA,
27 A.R. Use of Aloe vera-based extender for chilling and freezing collared peccary (*Pecari*
28 *tajacu*) semen. **Theriogenology**, v. 85, p. 1432–1438, 2016.
- 29 TECHAKUMPHU, M.; RUNGSIWIWUT, R.; NUMCHAIKRIKA, P.; THONGPHAKDEE, A.
30 Cloned Asian elephant (*Elephas maximus*) embryos reconstructed from rabbit recipient
31 oocytes. **The Thai Journal of Veterinary Medicine**, v. 40, p. 63–68, 2010.
- 32 THEIMER, T.C.; KEIM, P. Geographic patterns of mitochondrial-DNA variation in collared
33 peccaries. **Journal of Mammalogy**, v. 75, p. 121–128, 1994.

- 1 VENTURIERI, B; PENDU, L.Y. Padrões de atividades de caíaitus (*Tayassu tajacu*) em
2 cativeiro. **Revista de Etologia**, v. 8, p. 35–43, 2006.
- 3 VERAGUAS, D.; GALLEGOS, P.F.; CASTRO, F.O.; RODRIGUEZ-ALVAREZ, L. Cell cycle
4 synchronization and analysis of apoptosis-related gene in skin fibroblasts from domestic cat
5 (*Felis silvestris catus*) and kodkod (*Leopardus guigna*). **Reproduction in Domestic Animals**,
6 v. 52, p. 881–889, 2017.
- 7 WANI, N.A.; HONG, S.; VETTICAL, B.S. Cytoplasm source influences development of
8 somatic cell nuclear transfer (SCNT) embryos *in vitro* but not their development to term after
9 transfer to synchronized recipients in dromedary camels (*Camelus dromedarius*).
10 **Theriogenology**, v. 118, p. 137–143, 2018.
- 11 WITTAYARAT, M.; THONGPHAKDEE, A.; SAIKHUN, K.; CHATDARONG, K.; OTOI, T.;
12 TECHAKUMPHU, M. Cell cycle synchronization of skin fibroblast cells in four species of
13 family Felidae. **Reproduction in Domestic Animals**, v. 48, p. 305–310, 2013.
- 14 WU, G.Q.; QUAN, G.B.; SHAO, Q.Y.; LV, C.R.; JIANG, Y.T.; ZHAO, Z.Y.; HONG, Q.H.
15 Cryotop vitrification of porcine parthenogenetic embryos at the early developmental stages.
16 **Theriogenology**, v. 85, p. 434–440, 2017.
- 17
- 18

**CAPÍTULO 2 – POTENTIAL ROLE OF INTRASPECIFIC AND INTERSPECIFIC
CLONING IN THE CONSERVATION OF WILD MAMMALS (Artigo de revisão)**

PUBLICADO NA ZYGOTE

QUALIS: Qualis 2013-2016: B1/ Novo Qualis: B2

FATOR DE IMPACTO: 1,278

DOI: <https://doi.org/10.1017/S0967199419000170>

Potential role of intraspecific and interspecific cloning in the conservation of wild mammals

Alana Azevedo Borges¹ and Alexsandra Fernandes Pereira¹⁻²

¹Laboratory of Animal Biotechnology, Federal Rural University of Semi-Arid, Mossoro, RN, Brazil.

Running headline: Conservation of wild mammals.

²All correspondence to: Alexsandra Fernandes Pereira

Laboratory of Animal Biotechnology

Federal Rural University of Semi-Arid

Av. Francisco Mota, 572, Mossoró, RN, 59625-900, Brazil

Phone: +55 84 3317 8361

E-mail address: alexsandra.pereira@ufersa.edu.br

Summary

Intraspecific and interspecific cloning via somatic cell nuclear transfer (iSCNT) is a biotechnique with great possibilities for wild mammals because it allows for the maintenance of biodiversity by recovering species, nuclear reprogramming for the production of pluripotency-induced cells, and studies related to embryonic development. Nevertheless, many areas in cloning, especially those associated with wild mammals, are still in question because of the difficulty in obtaining cytoplasmic donor cells (or cytoplasts). On the other hand, donor cell nuclei (or karyoplasts) are widely obtained from the skin of living or post-mortem individuals and often maintained in somatic cell banks. Moreover, the creation of karyoplast-cytoplast complexes by fusion followed by activation and embryo development is one of the most difficult steps that require further clarification to avoid genetic failures. Although difficult, cloning different species, such as wild carnivores and ungulates, can be successful via iSCNT with embryo development and the birth of offspring. Thus, novel research in the area that contributes to the conservation of biodiversity and knowledge of the physiology of species continues. The present review presents the failures and successes that occurred with the application of the technique in wild mammals, with the goal of helping future work on cloning via iSCNT.

Keywords: Biodiversity, Biotechnology, Embryo development, SCNT, Somatic cells.

Introduction

The decrease in the biodiversity of wild mammals has been caused mainly by human activity, resulting in an increase in research aimed at the development of conservation strategies (Pereira *et al.*, 2016). In general, several techniques to help with the conservation of wild animals are available, including the formation of biobanks (León-Quinto *et al.*, 2009), artificial insemination (Howard *et al.*, 2016), embryo transfer (Goeritz *et al.*, 2012), in vitro fertilization (Herrick *et al.*, 2010), and cloning using somatic cell nuclear transfer (SCNT, Folch *et al.*, 2009). Because of the low availability of oocytes for SCNT, interspecific cloning using intraspecific and interspecific nuclear transfer techniques (iSCNT) has been shown to be an important tool in conservation (Wani *et al.*, 2017).

The main argument for the application of iSCNT is the rapid decrease in the number of species. Any tool that can avoid this decrease is important. iSCNT preserves and even expands genetic variability when somatic cells of different individuals representative of the

original biodiversity of a population are collected for its use (Loi *et al.*, 2001). In addition, interest in cloning has increased not only for the conservation of endangered species, but also for the multiplication of reproducers with better genetic characteristics (Saini *et al.*, 2015), basic research on cell epigenetic status (Saragusty *et al.*, 2016), embryonic development (González-Grajales *et al.*, 2016), and the production of induced pluripotent cells (Sukparangsi *et al.*, 2018).

Therefore, in all applications of cloning, studies related to the improvement of iSCNT, as well as its wide use in different individuals, are important.

Overview of the iSCNT technique and its limitations

The iSCNT technique involves embryo reconstruction by fusing a nucleus of a donor cell (karyoplast) derived from a wild mammal with an enucleated oocyte (cytoplast) from a domestic mammal of a different species, family, order, or class (Do & Taylor-Robinson, 2014). The nucleus in G0/G1 is exposed to reprogramming by the oocyte, followed by the fusion and activation of the reconstructed embryo (Loi *et al.*, 2011). Subsequently, the resultant embryo can be transplanted into the uterus of a recipient for term development (Pereira & Freitas, 2009).

Different steps are involved in the production of clones via iSCNT. Therefore, it is interesting to highlight the steps of the technique and its peculiarities that can define the success of cloning by iSCNT.

Preparation of cytoplasts

Whether using the oocyte from a domestic or a wild mammal, some fundamental criteria must be met to obtain a cytoplast suitable for cloning, such as oocyte selection, in vitro maturation, and enucleation systems (Loi *et al.*, 2011). In general, follicular size, the oocyte collection method, and the culture environment are factors that can affect the quality of mature oocytes, and different responses to these factors can be observed in wild mammals. In some cervid species, Brahmansani *et al.* (2013) observed that low maturation rates could probably be caused by the slicing. In this method, non-competent oocytes can be recovered since the technique can result in the recovery of structures of small diameter follicles (Rho *et al.*, 2001). Additionally, the quality of ovaries obtained post-mortem may have been one of the factors that reduce the quality of oocytes in these species.

Thus, studies have shown that enriched culture media and ovarian transport conditions may result in good results using ovaries from post-mortem animals for recovery of immature oocytes. Macías-García *et al.* (2018) verified that oocytes of Iberian red deer (*Cervus elaphus hispanicus*) obtained from ovaries maintained for 16 h in a holding medium increased the oocyte meiotic competence. Moreover, these authors observed that the epidermal growth factor (EGF) demonstrated a differential effect depending upon oocyte gradient and conditions of ovary transportation. Additionally, for ovaries derived from Hokkaido sika deer (*Cervus nippon yesoensis*), maturation rates of oocytes were highest when ovaries were kept for 12 h at 20–25 °C, when compared to 24 h (Tulake *et al.*, 2014).

Specifically, with respect to the culture environment, the requirements for both composition and maturation time should be established for the in vitro maturation of each species. In the Indian blackbuck (*Antelope cervicapra*), oocytes cultured in the presence of gonadotropins (follicle-stimulating hormone, FSH and luteinizing hormone, LH) showed higher rates of expansion of the cumulus oophorus (79.3%) and extrusion of the first polar body (46.1%) compared to oocytes cultured without gonadotropins (60.4% and 33.3%, respectively) (Rao *et al.*, 2010). In the Sika deer (*Cervus nippon hortulorum*), oocytes cultured in medium supplemented with fetal bovine serum (FBS), FSH, LH, cysteamine and EGF resulted in a higher maturation rate (75.4%) compared to medium without supplementation (30.1%; Yin *et al.*, 2013). Already, different hormonal combinations of FSH, LH and 17 β -estradiol did not alter the maturation rates in oocytes derived from lions (*Panthera leo*, Fernandez-Gonzalez *et al.*, 2015). In the collared peccary (*Pecari tajacu*), we proved that oocytes need 48 h to achieve maturation instead of 24 h, according to the expansion of the cumulus cells (100% vs. 38.1%), the presence of first polar body (90.5% vs. 52.4%), and the status of the nucleus in the second metaphase (76.2% vs. 52.4%), respectively (Borges *et al.*, 2018c).

In addition to obtaining mature oocytes, the preparation of cytoplasts depends on the method of enucleation of these structures. The amount of ooplasm present in the reconstructed embryo is related to the enucleation technique that removes the nucleus from the oocyte. Matured oocytes can be enucleated in different ways, including squeezing the first polar body and the surrounding cytoplasm through a cleft in the zona pellucida of the oocyte (Lee *et al.*, 2003). Another method is manual removal in which zona-free oocytes are enucleated with a bisection blade that handbisect the metaphase II chromosomes along with a small volume of

the surrounding cytoplasm. Oocytes can also be aspirated using a micromanipulator at the location of the metaphase II chromosomes and the polar body via brief exposure to ultraviolet light (Pereira *et al.*, 2015).

Selection of karyoplasts

To obtain karyoplasts appropriate for cloning, their type and age and the manipulation techniques used are important for their future reprogramming (Kim *et al.*, 2007). Karyoplasts can be obtained from fresh or cryopreserved somatic tissues (Folch *et al.*, 2009, Pan *et al.*, 2014), from an adult (Moulavi *et al.*, 2017) or a fetus (Liu *et al.*, 2018), and in vivo or post-mortem (Pereira *et al.*, 2014). Although the recovery of these cells is not a difficult task, their processing and preservation until use in iSCNT require attention (Pereira *et al.*, 2014). In general, skin cells have been the most used cell type for karyoplasts (Song *et al.*, 2007). The skin has an abundance of cells of interest that may have different efficiencies in cloning, as observed in wild buffalo (*Bubalus arnee*). Saini *et al.* (2015) detected that fibroblasts of this species are easier to reprogram than epithelial cells.

After harvest, cells used as nuclei donors need to be characterized with respect to their culture conditions, cryopreservation, and cell cycle synchronization (Pereira *et al.*, 2014). For these steps, cells are evaluated for the number of passages, nutritional requirements during in vitro culture (Santos *et al.*, 2016), and the damage done during cryopreservation (Song *et al.*, 2007). Thus, karyoplasts have been established in vitro in Dulbecco's modified Eagle's medium (DMEM) supplemented with FBS and growth factors (Santos *et al.*, 2016).

Karyoplasts have been routinely cryopreserved by slow freezing (Sharma *et al.*, 2018) using a combination of dimethyl sulfoxide (DMSO), FBS, and sucrose as the cryoprotectant, as observed with Iberian lynx (*Lynx pardinus*, León-Quinto *et al.*, 2014). Although it is more desirable to use a somatic cell bank after tissue culture, the absence of in vitro culture conditions sometimes makes these banks unfeasible, resulting in the immediate formation of the targets for those somatic tissues (Borges *et al.*, 2017a,b; Queiroz Neta *et al.*, 2018). The three somatic tissue conservation techniques used for wild animals are slow-freezing cryopreservation (Mestre-Citrinovitz *et al.*, 2016), vitrification (Borges *et al.*, 2018a,b), and cooling at 4–6°C (Queiroz Neta *et al.*, 2018). In collared peccaries, we compared two techniques of vitrification and we observed that the solid-surface vitrification was found to be more efficient method for vitrifying skin tissue when compared to direct vitrification in

cryovials, probably due tissues not be involved in a large amount of cryoprotectants before passing through a drastic change in temperature during the solid-surface vitrification (Borges *et al.*, 2017b).

Finally, the third step in the preparation of the karyoplasts is cell synchronization in the G0/G1 stage (Gómez *et al.*, 2003; Yelisetti *et al.*, 2016). In general, nuclear reprogramming is controlled by epigenetic modification. For this to occur, the somatic cells must be in G0/G1 to allow the removal of reversible epigenetic changes acquired during cell differentiation (Song *et al.*, 2007). Thus, cells can be subjected to different treatments for synchronization during culture. Inhibition by contact (Moulavi *et al.*, 2017), serum deprivation (Wani *et al.*, 2017), and chemicals that inhibit the cell cycle (Gómez *et al.*, 2003) are methods used for synchronization. Serum deprivation and inhibition by contact are the most commonly used (Moulavi *et al.*, 2017).

Under high confluence or serum privation, fibroblast cells derived from the skin of adult argali (*Ovis ammon*) were efficiently synchronized at G0/G1; nevertheless, cells were in lower proportion in the growing stage (Pan *et al.*, 2014). Authors observed that the highest proportion of cells from the African wild cat (*Felis silvestris lybica*) at G0/G1 was obtained by serum deprivation compared with that obtained by inhibition by contact and the inhibitor roscovitine (Gómez *et al.*, 2003). Leopard (*Panthera pardus*) skin cells treated with chemical inhibitors such as sodium butyrate have a greater propensity to undergo alterations (Yelisetti *et al.*, 2016).

Embryonic reconstruction stages

After the transfer of the nucleus into the enucleated oocyte, the cytoplasm-karyoplast complex is subjected to an electric pulse that not only induces the fusion of the somatic cell nucleus with the enucleated oocyte to form a new complex, but also promotes the release of intracellular calcium that initiates cellular activation (Pereira & Freitas, 2009). In general, the successful development of a reconstructed embryo depends on the complex interactions between the cytoplasm and the nuclear structure during embryonic development; failures in this interaction can cause problems during early cleavage and embryonic development (González-Grajales *et al.*, 2016).

1 The activation of the cytoplasm-karyoplast complexes guarantees adequate embryonic
2 reconstruction (Yamochi *et al.*, 2013). Because the iSCNT technique reprograms the nucleus
3 of a somatic cell of one species using the oocyte cytoplasm of another species, it is essential
4 that the activation protocol be able to activate the reconstructed embryo (Zhao *et al.*, 2006).
5 Physiologically, a mammalian oocyte is activated during fusion with a sperm, releasing
6 meiotic cell cycle arrest and enabling the resumption of the oocyte meiotic cell cycle
7 (Sparman *et al.*, 2010). Therefore, a well-developed protocol allows a high rate of blastocyst
8 formation by promoting good embryonic development through activation.

9
10 Activation protocols, including physical methods such as electrical pulses and alteration of
11 osmolarity, and chemical methods such as calcium-mobilizing compounds like strontium
12 chloride, ionomycin, and ethanol, to promote the initial release of calcium have been
13 evaluated in different species, as sika deer (Yin *et al.*, 2013), alpaca (*Vicugna pacos*) and
14 llama (*Lama glama*, Ruiz *et al.*, 2015), with blastocyst rates of 32.4%, 22.5% and 18.7%,
15 respectively. In general, calcium mobilizers are used in combination with kinase protein
16 inhibitors or protein synthesizers such as cyclohexamide and 6-dimethylaminopurine (6-
17 DMAP). In addition, a cytostatic factor inactivator and microfilament inhibitor such as
18 cytochalasin B are used to prevent extrusion of the second polar body and maintain the
19 diploidy of the presumed embryo (Ruiz *et al.*, 2015).

20
21 For red deer (*Cervus elaphus*), electrical activation before chemical activation with ionomycin
22 and 6-DMAP was efficient for the production of clone embryos (32–44%), obtaining
23 genetically healthy calves (Berg *et al.*, 2007). Nevertheless, the same protocol resulted in a
24 low developmental rate (5.7%) of activated oocytes in swamp deer and 0.0% embryos in
25 spotted deer, sambar deer, and Brow-antlered deer after oocyte parthenogenetic activation
26 (Brahmasani *et al.*, 2013). Blackbuck (*Antilope cervicapra*) oocytes activated with ionomycin
27 and 6-DMAP resulted in 58% cleaved embryos and 13% blastocysts (Rao *et al.*, 2010).
28 Therefore, the artificial activation method (chemical, electrical protocols or your
29 combination) can result in different responses among species. In this sense, it is necessary to
30 evaluate the type of artificial activation that promotes the best rates of embryonic
31 development in the species of interest.

32
33 In vitro culture systems are essential for early embryonic development and nuclear
34 reprogramming (Gómez *et al.*, 2008; Pereira *et al.*, 2013). Choosing the appropriate culture

medium for each species is considered the initial step in proper embryonic development (Zhao *et al.*, 2006). Lee *et al.* (2003) used somatic cells of the mountain bongo (*Tragelaphus eurycerus isaaci*) and domestic cow (*Bos taurus*) oocytes and observed that a chemically defined, protein-free medium of TCM199 supplemented with FBS supported embryonic development. Nonetheless, there is no one culture medium suitable for all species that allows better embryonic development for a given species under study.

Finally, the effect of epigenetic reprogramming is a very relevant factor in the success of iSCNT (Gómez *et al.*, 2008). Some epigenetic markers were characterized with respect to their function during embryonic reprogramming and their influence on the chromatin structure from post-translational modifications (Song *et al.*, 2007). The overall level of the acetylation of histone H3 at lysine 18 (H3K18ac) and trimethylation of histone H3 at lysine 27 (H3K27me3), and the expression level of some important apoptosis proteins (caspase 3 and caspase 7), and p53 were evaluated. The hyperacetylated state of histones is associated with transcriptionally active domains, while the hypoacetylated state is associated mainly with silenced chromatin regions of histone acetyl transferases and histone deacetylases. The methylation pattern of the DNA is determined by DNA methyltransferases. OCT3/4, NANOG, and CDX2 are very important because of their close association with pluripotency and early embryonic development (Saini *et al.*, 2015).

Advances and perspectives of iSCNT in wild mammals

Several works aimed at cloning different wild mammals have been conducted (Table 1). Among these studies, those that obtained offspring were on wild bovine (Lanza *et al.*, 2000), sheep (Loi *et al.*, 2001), felid (Gómez *et al.*, 2004; Li *et al.*, 2007), canid (Kim *et al.*, 2007; Oh *et al.*, 2008), and goat (Folch *et al.*, 2009). Therefore, several families have proven the success of using iSCNT for the recovery and reintroduction of wild mammals.

An important point to remember is that as the taxonomic distance between donor and recipient species increases, the production of blastocysts decreases because of the decreased ability of somatic cells to be reprogrammed (Priya *et al.*, 2014). In general, enucleated oocytes are from a domestic species that is phylogenetically close to the wild species that donates the nucleus. For example, domestic sheep cytoplasts were able to reprogram argali fibroblast nuclei (Pan *et al.*, 2014) and domestic buffalo cytoplasm was able to reprogram wild buffalo karyoplast (Priya *et al.*, 2014).

Table 1 iSCNT in some wild mammals.

Scientific Name	IUCN Category*	Country	Karyoplast	Cytoplast	Outcome	Authors
Carnivores						
<i>Canis lupus</i>	Least Concern	Seoul, Korea	Fibroblast	<i>Canis lupus familiaris</i>	17% pregnancies, 2 pups	Kim <i>et al.</i> (2007)
<i>Canis lupus</i>	Least Concern	Seoul, Korea	Fibroblast	<i>Canis lupus familiaris</i>	23.5% pregnancies, 4 pups	Oh <i>et al.</i> (2008)
<i>Felis margarita</i>	Least Concern	USA	Fibroblast	<i>Felis catus</i>	3 pups	Gómez <i>et al.</i> (2008)
<i>Felis silvestris lybica</i>	Least Concern	USA	Fibroblast	<i>Felis catus</i>	28% blastocyst	Gómez <i>et al.</i> (2003)
<i>Felis silvestris lybica</i>	Least Concern	USA	Fibroblast	<i>Felis catus</i>	75% pregnancies, 17 pups	Gómez <i>et al.</i> (2004)
Ungulates						
<i>Acinonyx jubatus</i>	Vulnerable	Argentina	Fibroblast	<i>Felis catus</i>	27.4% blastocyst	Moro <i>et al.</i> (2015)
<i>Acinonyx jubatus venaticus</i>	Vulnerable	Iran	Fibroblast	<i>Felis catus</i>	5.9% morula	Moulavi <i>et al.</i> (2017)
<i>Bos gaurus</i>	Vulnerable	USA	Fibroblast	<i>Bos taurus</i>	25% pregnancies, none term	Lanza <i>et al.</i> (2000)
<i>Bos javanicus</i>	Endangered	USA	Fibroblast	<i>Bos taurus</i>	17% pregnancies, none term	Sansinena <i>et al.</i> (2005)
<i>Bubalus arnee</i>	Endangered	India	Fibroblast	<i>Bubalus</i>	38.7% blastocyst	Priya <i>et al.</i> (2014)

<i>Bubalus arnee</i>	Endangered	India	Fibroblast/Epithelial cell	<i>bubalis</i> <i>Bubalus</i>	50.6 vs 20.5% blastocyst	Saini <i>et al.</i> (2015)
<i>Capra ibex</i>	Least Concern	China	Fibroblast	<i>bubalis</i> <i>Capra</i>	11% blastocyst	Wang <i>et al.</i> (2007)
<i>Capra pyrenaica pyrenaica</i>	Absente	Spain	Fibroblast	<i>hircus</i> <i>Capra</i>	1 born	Folch <i>et al.</i> (2009)
<i>Ovis ammon</i>	Near Threatened	China	Fibroblast/ <i>Cumulus</i> cell	<i>pyrenaica</i> <i>Ovis aries</i>	22.1% blastocyst	Pan <i>et al.</i> (2014)
<i>Ovis orientalis musimon</i>	Vulnerable	Italy	Granulosa cell	<i>Ovis aries</i>	1 pup	Loi <i>et al.</i> (2001)
<i>Tragelaphus eurycerus isaaci</i>	Critically Endangered	USA	Fibroblast	<i>Bos taurus</i>	24% blastocyst	Lee <i>et al.</i> (2003)
Others species						
<i>Macaca fascicularis</i>	Least Concern	Thailand	Fibroblast	<i>Bos taurus</i>	33% blastocyst	Lorthongpanich <i>et al.</i> (2008)
<i>Macaca fascicularis</i>	Least Concern	China	Fibroblast	<i>Macaca</i> <i>fascicularis</i>	2 born	Liu <i>et al.</i> (2018)
<i>Balaenoptera bonaerensis</i>	Data deficient	Japan	<i>Cumulus</i> cells	<i>Bos taurus</i> / <i>Sus scrofa</i> <i>domesticus</i>	27.5–52.8% cleavage	Ikumi <i>et al.</i> (2004)

*IUCN: International Union for Conservation of Nature and Natural Resources. USA: United States of America

1 *Carnivores*

2 Some works have shown the advances achieved by iSCNT in wild canine species. These species,
 3 including the gray wolf (*Canis lupus*), have gradually become endangered or extinct. Thus, in
 4 2007, with the goal of canid conservation, Kim *et al.* (2007) cultured fibroblasts derived from the
 5 ear of an adult female gray wolf that were then used as donor cells of nuclei. Using domestic
 6 canine oocytes, the authors produced a pregnancy with cloned embryos of two genetic identities
 7 of the cloned wolves, but there were no births. In 2008, Oh *et al.* (2008) obtained three wolf pups
 8 from cloned embryos using cells obtained from a male gray wolf 6 h after death and domestic
 9 canine oocytes. These studies demonstrated the successful cloning of endangered wild canines.

10
 11 In felid species, the main oocyte source has been the domestic cat. In species from the Felidae
 12 subfamily, some progress has been achieved. Thus, synchronized nuclei were donated by the
 13 African wild cat and transferred to enucleated domestic cat oocytes resulting in a high rate of
 14 blastocyst formation but no pregnancies (Gómez *et al.*, 2003). In another study in which embryos
 15 were constructed using somatic cells derived from the African wild cat and domestic cat oocytes,
 16 75% of the embryos developed to term and 25% underwent fetal resorption or abortion (Gómez
 17 *et al.*, 2004). Of the 17 cloned kittens born, seven were stillborn, eight died within hours of
 18 delivery or up to 6 weeks of age, and two are alive and healthy. Additionally, some studies on
 19 wild felids have shown the establishment of somatic resource banks. There is a bank with somatic
 20 samples of 69 individual Iberian lynx, considered the most endangered felid in the world, with
 21 the aim of future cloning (León-Quinto *et al.*, 2009; 2014).

22
 23 In addition, works on the cheetah (*Acinonyx jubatus*), a species of the Pantherinae subfamily,
 24 have been performed in South America and Asia. Somatic cells from a cheetah raised in South
 25 America were transferred to domestic cat oocytes, and, after embryo aggregation during in vitro
 26 culture, high blastocyst formation rates were obtained (16.7%–28.3%) (Moro *et al.*, 2015).
 27 Moulavi *et al.* (2017) used nonviable frozen cells derived from frozen tissue from an Asiatic
 28 cheetah (*Acinonyx jubatus venaticus*) and in vitro-matured domestic cat oocytes and obtained
 29 morula rates of 5.9%. Although no blastocyst was obtained, this study demonstrated that
 30 enucleated cat oocytes can partially remodel and reactivate nonviable nuclei of the Asiatic
 31 cheetah and support its reprogramming back to the embryonic stage.

1 *Ungulates*

2 Some studies with ungulates have been performed with significant success, especially for species
3 already extinct. The first animal derived from an extinct subspecies was obtained using
4 fibroblasts from skin biopsies collected before the death of the last female *Capra pyrenaica*
5 *pyrenaica*. After a year under cryopreservation, these cells were used as karyoplasts and fused
6 with the cytoplasts of a domestic goat to reconstruct embryos. The rate of cleaved embryos after
7 36 h was 47.3%, of which 65.5% were transferred. Five recipients were pregnant at 45 days but
8 only one pregnancy went to term. Unfortunately, a few minutes after birth the animal died from
9 pulmonary complications (Folch *et al.*, 2009).

10
11 Experiments were carried out with wild yak (*Bros grunniens*) with the goal of evaluating the
12 parameters that affect the success of iSCNT (Li *et al.*, 2007). Fibroblasts and cumulus cells were
13 used as donor cells, but the cell type and different ages were found to have no significant effect
14 on iSCNT.

15
16 In 2017, the birth of a Bactrian camel cloned by iSCNT was first reported (Wani *et al.*, 2017).
17 The fibroblasts used to donate nuclei were obtained from ear skin biopsy samples from an adult
18 male Bactrian camel (*Camelus bactrianus*) and the cytoplast of dromedary camel (*Camelus*
19 *dromedaries*) was the oocyte recipient. Twenty-six blastocysts were transferred to 23
20 synchronized dromedary recipients yielding five pregnancies with one going to term. This work
21 has great importance because the Bactrian camel is the eighth most endangered large mammal on
22 Earth.

23
24 Finally, the woolly mammoth (*Mammuthus primigenius*) is perhaps the one wild mammal of the
25 ungulates whose cloning arouses the greatest interest. This animal became extinct about 10,000
26 years ago. However, epithelial and muscular cells from 14,000–15,000-year-old mammoth
27 tissues were cryopreserved, with the goal of producing embryos of this species (Kato *et al.*,
28 2009). In this study, the authors injected cell nucleus-like structures into mature mouse
29 enucleated oocytes; however, the oocytes did not form pronuclear-like structures at 7 h after
30 injection.

1 *Other species*

2 The ability of bovine enucleated oocytes to support dedifferentiation of nuclei from monkey
 3 fibroblasts in interspecies cloned monkey embryos has been observed (Lorthongpanich *et al.*,
 4 2008). These embryos were cultured in conditions different from the medium used for cattle with
 5 monkey-specific alterations, but the embryos were not able to develop past 16 cells under any
 6 culture condition. Nevertheless, OCT-4 was detected, demonstrating the ability of bovine
 7 ooplasm to support dedifferentiation but not embryonic development. Therefore, the culture
 8 medium promotes dedifferentiation but is not able to support complete embryonic development
 9 (Lorthongpanich *et al.*, 2008). In another work that used porcine cytoplasts and donor cells from
 10 a rhesus monkey, it was possible to obtain blastocysts despite the low rate (2.04%) (Zhu *et al.*,
 11 2014). Although being a SCNT study, the cloning of cynomolgus monkeys (*Macaca fascicularis*)
 12 is cited here because of recent advances in this species. Thus, in a study on cynomolgus monkeys
 13 using SCNT, Liu *et al.* (2018) applied histone demethylase Kdm4d mRNA and histone
 14 deacetylase inhibitor trichostatin A after activation. Embryonic development improved followed
 15 by a greater number of pregnancies, which resulted in the birth of two monkeys via the SCNT
 16 technique using fetal fibroblasts and oocytes of cynomolgus monkeys.

17
 18 With respect to aquatic mammals, a study performed on the minke whale (*Balaenoptera*
 19 *bonaerensis*) compared different conditions of iSCNT, including the ability of porcine and bovine
 20 ooplasm to produce reconstructed embryos and the effects of different donor cell types (viable
 21 or nonviable cells) on whale SCNT embryos (Ikumi *et al.*, 2004). The authors concluded that
 22 whale iSCNT embryos can develop to at least the four-cell stage, regardless of the survivability
 23 of the donor cells and the porcine or bovine ooplasm.

24 25 **Final considerations**

26 Although cloning has several technical limitations that require greater attention to improve the
 27 technique, iSCNT has been applied to numerous species of wild mammals and has achieved
 28 positive results with respect to embryonic stages in pregnancies and offspring born. The works
 29 cited in this paper have made it possible to analyze the state of the art and to perform specific
 30 studies the problems in the technique that can be fixed according to the species being studied.

This review has shown that there is no rule that says several species should be cloned following the same protocol, but that each species has different needs at each stage of the technique. In addition, all the papers referred to in this review point to the need for improvement and study at a certain stage, which will lead to improvement of the technique. Thus, to achieve a satisfactory result with iSCNT, each step involved in cloning must be suitable for the species being studied.

Although iSCNT is not the main tool for the reestablishment of endangered wild mammals, its use to increase the possibilities of reproduction and multiplication of individuals has been proposed. It should be refined so that it can be an alternative when traditional techniques cannot be applied. In addition, cloning helps elucidate the embryonic development of a wild species and the subsequent application of this knowledge.

Finally, this biotechnology can help generate more ways to maintain individual species. Therefore, the improvement of protocols to potentiate this technique is of interest because although it has low efficiency rates, iSCNT shows promise because of the pups of different species that have been born.

Acknowledgements

Alana Azevedo Borges is a recipient of a grant from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES, Financial Code 001). Alexsandra Fernandes Pereira is recipient of a grant from CNPq (no. 306963/2017-5). None of the authors has any conflict of interest to declare.

References

- Berg, D.K., Li, C., Asher, G., Wells, D.N. & Oback, B. (2007). Red deer cloned from antler stem cells and their differentiated progeny. *Biol. Reprod.* **77**, 384–94.
- Borges, A.A., Bezerra, F.V.B., Costa, F.N., Queiroz Neta, L.B., Santos, M.V.O., Oliveira, M.F., Silva, A.R. & Pereira, A.F. (2017a). Histomorphological characterization of collared peccary (*Pecari tajacu* Linnaeus, 1758) ear integumentary system. *Arq. Bras. Med. Vet. Zootec.* **69**, 948–54.
- Borges, A.A., Lima, G.L., Queiroz Neta, L.B., Santos, M.V.O., Oliveira, M.F., Silva, A.R. &

- 1 Pereira, A.F. (2017b). Conservation of somatic tissue derived from collared peccaries (*Pecari*
2 *tajacu* Linnaeus, 1758) using direct or solid-surface vitrification techniques. *Cytotechnology*
3 **69**, 643–54.
- 4 Borges, A.A., Lira, G.P.O., Nascimento, L.E., Queiroz Neta, L.B., Santos, M.V.O., Oliveira, M.F.,
5 Silva, A.R. & Pereira, A.F. (2018a). Influence of cryopreservation solution on the *in vitro*
6 culture of skin tissues derived from collared peccary (*Pecari tajacu* Linnaeus, 1758).
7 *Biopreserv. Biobank*. **16**, 77–81.
- 8 Borges, A.A., Queiroz Neta, L.B., Santos, M.V.O., Oliveira, M.F., Silva, A.R. & Pereira, A.F.
9 (2018b). Combination of ethylene glycol with sucrose increases survival rate after vitrification
10 of somatic tissue of collared peccaries (*Pecari tajacu* Linnaeus, 1758). *Pesq. Vet. Bras.* **38**,
11 350–56.
- 12 Borges, A.A., Santos, M.V.O., Queiroz Neta, L.B., Oliveira, M.F., Silva, A.R. & Pereira A.F.
13 (2018c). *In vitro* maturation of collared peccary (*Pecari tajacu* Linnaeus, 1758) oocytes after
14 different incubation times. *Pesq. Vet. Bras.* **38**, 1863–68.
- 15 Brahmasani, S.R., Yelisetti, U.M., Katari, V., Komjeti, S., Lakshmikantan, U., Pawar, R.M. &
16 Sisinthy, S. (2013). Developmental ability after parthenogenetic activation of *in vitro* matured
17 oocytes collected postmortem from deers. *Small Rumin. Res.* **113**, 128–35.
- 18 Do, V.H. & Taylor-Robinson, A. (2014). Somatic cell nuclear transfer in mammals reprogramming
19 mechanism and factors affecting success. *Clon. Transgen.* **3**, 129.
- 20 Fernandez-Gonzalez, L., Hribal, R., Stagegaard, J., Zahmel, J. & Jewgenow, K. (2015). Production
21 of lion (*Panthera leo*) blastocysts after *in vitro* maturation of oocytes and intracytoplasmic
22 sperm injection. *Theriogenology* **83**, 995–9.
- 23 Folch, J., Cocero, M., Chesné, P., Alabart, J., Domínguez, V., Cognié, Y., Roche, A., Fernández-
24 Arias, A., Martí, J. & Sánchez, P. (2009). First birth of an animal from an extinct subspecies
25 (*Capra pyrenaica pyrenaica*) by cloning. *Theriogenology* **71**, 1026–34.
- 26 Goeritz, F., Painer, J., Jewgenow, K., Hermes, R., Rasmussen, K., Dehnhard, M. & Hildebrandt, T.
27 (2012). Embryo retrieval after hormonal treatment to control ovarian function and non-surgical
28 artificial insemination in African lions (*Panthera leo*). *Reprod. Domest. Anim.* **47**, 156–60.
- 29 Gómez, M.C., Jenkins, J.A., Giraldo, A., Harris, R.F., King, A., Dresser, B.L. & Pope, C.E. (2003).
30 Nuclear transfer of synchronized African wild cat somatic cells into enucleated domestic cat
31 oocytes. *Biol. Reprod.* **69**, 1032–41.

- 1 Gómez, M.C., Pope, C.E., Giraldo, A., Lyons, L.A., Harris, R.F., King, A.L., Cole, A., Godke,
2 R.A. & Dresser, B.L. (2004). Birth of African wild cat cloned kittens born from domestic cats.
3 *Cloning Stem Cells*. **6**, 247–58.
- 4 Gómez, M.C., Pope, C.E., Kutner, R.H., Ricks, D.M., Lyons, L.A., Ruhe, M., Dumas, C., Lyons,
5 J., López, M. & Dresser, B.L. (2008). Nuclear transfer of sand cat cells into enucleated
6 domestic cat oocytes is affected by cryopreservation of donor cells. *Cloning Stem Cells* **10**,
7 469–84.
- 8 González-Grajales, L.A., Favetta, L.A., King, W.A. & Mastromonaco, G.F. (2016). Lack of effects
9 of ooplasm transfer on early development of interspecies somatic cell nuclear transfer bison
10 embryos. *BMC Dev. Biol.* **16**, 36.
- 11 Herrick, J., Campbell, M., Levens, G., Moore, T., Benson, K., D’Agostino, J., West, G., Okeson,
12 D., Coke, R. & Portacio, S. (2010). *In vitro* fertilization and sperm cryopreservation in the
13 black-footed cat (*Felis nigripes*) and sand cat (*Felis margarita*). *Biol. Reprod.* **82**, 552–62.
- 14 Howard, J., Lynch, C., Santymire, R., Marinari, P. & Wildt, D. (2016). Recovery of gene diversity
15 using long-term cryopreserved spermatozoa and artificial insemination in the endangered
16 black-footed ferret. *Anim. Conserv.* **19**, 102–11.
- 17 Ikumi, S., Sawai, K., Takeuchi, Y., Iwayama, H., Ishikawa, H., Ohsumi, S. & Fukui, Y. (2004).
18 Interspecies somatic cell nuclear transfer for *in vitro* production of Antarctic minke whale
19 (*Balaenoptera bonaerensis*) embryos. *Cloning Stem Cells*. **6**, 284–93.
- 20 Kato, H., Anzai, M., Mitani, T., Morita, M., Nishiyama, Y., Nakao, A., Kondo, K., Lazarev, P.A.,
21 Ohtani, T. & Shibata, Y. (2009). Recovery of cell nuclei from 15,000 years old mammoth
22 tissues and its injection into mouse enucleated matured oocytes. *Proc. Jpn. Acad Ser. B. Phys.*
23 *Biol. Sci.* **85**, 240–47.
- 24 Kim, M.K., Jang, G., Oh, H.J., Yuda, F., Kim, H.J., Hwang, W.S., Hossein, M.S., Kim, J.J., Shin,
25 N.S. & Kang, S.K. (2007). Endangered wolves cloned from adult somatic cells. *Cloning Stem*
26 *Cells*. **9**, 130–37.
- 27 Lanza, R.P., Cibelli, J.B., Diaz, F., Moraes, C.T., Farin, P.W., Farin, C.E., Hammer, C.J., West,
28 M.D. & Damiani, P. (2000). Cloning of an endangered species (*Bos gaurus*) using interspecies
29 nuclear transfer. *Cloning*. **2**, 79–90.
- 30 Lee, B., Wirtu, G.G., Damiani, P., Pope, E., Dresser, B.L., Hwang, W. & Bavister, B.D. (2003).
31 Blastocyst development after intergeneric nuclear transfer of mountain bongo antelope somatic

- 1 cells into bovine oocytes. *Cloning Stem Cells*. **5**, 25–33.
- 2 León-Quinto, T., Simon, M.A., Cadenas, R., Jones, J., Martinez-Hernandez, F.J., Moreno, J.M.,
3 Vargas, A., Martinez-Hernandez, F.J. & Soria, B. (2009). Developing biological resource
4 banks as a supporting tool for wildlife reproduction and conservation: the Iberian lynx bank as
5 a model for other endangered species. *Anim. Reprod. Sci.* **112**, 347–61.
- 6 León-Quinto, T., Simón, M.A., Cadenas, R., Martínez, Á. & Serna, A. (2014). Different
7 cryopreservation requirements in foetal *versus* adult skin cells from an endangered mammal,
8 the Iberian lynx (*Lynx pardinus*). *Cryobiology*. **68**, 227–33.
- 9 Li, Y., Dai, Y., Du, W., Zhao, C., Wang, L., Wang, H., Liu, Y., Li, R. & Li, N. (2007). *In vitro*
10 development of yak (*Bos grunniens*) embryos generated by interspecies nuclear transfer. *Anim.*
11 *Reprod. Sci.* **101**, 45–59.
- 12 Liu, Z., Cai, Y., Wang, Y., Nie, Y., Zhang, C., Xu, Y., Zhang, X., Lu, Y., Wang, Z. & Poo, M.
13 (2018). Cloning of macaque monkeys by somatic cell nuclear transfer. *Cell*. **172**, 881–7.
- 14 Loi, P., Modlinski, J. & Ptak, G. (2011). Interspecies somatic cell nuclear transfer: a salvage tool
15 seeking first aid. *Theriogenology*. **76**, 217–28.
- 16 Loi, P., Ptak, G., Barboni, B., Fulka Jr, J., Cappai, P. & Clinton, M. (2001). Genetic rescue of an
17 endangered mammal by cross-species nuclear transfer using *post-mortem* somatic cells. *Nat.*
18 *Biotech.* **19**, 962.
- 19 Lorthongpanich, C., Laowtammathron, C., Chan, A.W.S., Ketudat-Cairns, M. & Parnpai, R.
20 (2008). Development of interspecies cloned monkey embryos reconstructed with bovine
21 enucleated oocytes. *J. Reprod. Dev.* **54**, 306–13.
- 22 Macías-García, B., González-Fernández, L., Matilla, E., Hernández, N., Mijares, J., & Sánchez-
23 Margallo, F.M. (2018). Oocyte holding in the Iberian red deer (*Cervus elaphus hispanicus*):
24 Effect of initial oocyte quality and epidermal growth factor addition on *in vitro* maturation.
25 *Reprod. Domest. Anim.* **53**, 243–48.
- 26 Mestre-Citrinovit, A.C., Sestelo, A.J., Ceballos, M.B., Baranao, J.L. & Saragueta, P. (2016).
27 Isolation of primary fibroblast culture from wildlife: the *Panthera onca* case to preserve a
28 South American endangered species. *Curr. Protoc. Mol. Biol.* **116**, 28.7.1–28.7.14.
- 29 Moro, L.N., Hiriart, M.I., Buemo, C., Jarazo, J., Sestelo, A., Veraguas, D., Rodriguez-Alvarez, L.
30 & Salamone, D.F. (2015). Cheetah interspecific SCNT followed by embryo aggregation
31 improves *in vitro* development but not pluripotent gene expression. *Reproduction* **150**, 1–10.

- 1 Moulavi, F., Hosseini, S.M., Tanhaie-Vash, N., Ostadhosseini, S., Hosseini, S.H., Hajinasrollah,
2 M., Asghari, M., Gourabi, H., Shahverdi, A. & Vosough, A. (2017). Interspecies somatic cell
3 nuclear transfer in Asiatic cheetah using nuclei derived from *post-mortem* frozen tissue in
4 absence of cryo-protectant and *in vitro* matured domestic cat oocytes. *Theriogenology*. **90**,
5 197–203.
- 6 Oh, H., Kim, M., Jang, G., Kim, H., Hong, S., Park, J., Park, K., Park, C., Sohn, S. & Kim, D.Y.
7 (2008). Cloning endangered gray wolves (*Canis lupus*) from somatic cells collected
8 *postmortem*. *Theriogenology*. **70**, 638–47.
- 9 Pan, X., Zhang, Y., Guo, Z. & Wang, F. (2014). Development of interspecies nuclear transfer
10 embryos reconstructed with argali (*Ovis ammon*) somatic cells and sheep ooplasm. *Cell Biol.*
11 *Inter.* **38**, 211–8.
- 12 Pereira, A.F. & Freitas, V.J.F. (2009). Cloning in ruminants: progress and current perspectives.
13 *Rev. Bras. Reprod. Anim.* **33**, 118–28.
- 14 Pereira, A.F., Feltrin, C., Almeida, K.C., Carneiro, I.S., Avelar, S.R.G., Alcântara Neto, A.S.,
15 Sousa, F.C., Melo, C.H.S., Moura, R.R., Teixeira, D.I.A., Bertolini, L.R., Freitas, V.J.F. &
16 Bertolini, M. (2013). Analysis of factors contributing to the efficiency of the *in vitro*
17 production of transgenic goat embryos (*Capra hircus*) by handmade cloning (HMC). *Small*
18 *Rumin. Res.* **109**, 163–72.
- 19 Pereira, A.F., Melo, L.M., Freitas, V.J.F. & Salamone, D.F. (2015). Phosphorylated H2AX in
20 parthenogenetically activated, *in vitro* fertilized and cloned bovine embryos. *Zygote* **23**, 485–
21 93.
- 22 Pereira, A.F., Santos, M.L.T., Borges, A.A., Queiroz Neta, L.B., Santos, M.V.O. & Feitosa, A.K.N.
23 (2014). Isolation and characterization of skin-derived donor cells for nuclear transfer. *Acta Vet.*
24 *Brasilica*. **8**, 311–6.
- 25 Pereira A.F., Silva A.R., Lima G.L. & Silva A.M. (2016). Somatic and gonadal tissue
26 cryopreservation: an alternative tool for the germplasm conservation in wild mammals. In:
27 Melanie Walton Editor. Germplasm: characteristics, diversity and preservation. Nova Science
28 Publishers, New York, pp. 80–117.
- 29 Priya, D., Selokar, N., Raja, A., Saini, M., Sahare, A., Nala, N., Palta, P., Chauhan, M., Manik, R.
30 & Singla, S. (2014). Production of wild buffalo (*Bubalus arnee*) embryos by interspecies
31 somatic cell nuclear transfer using domestic buffalo (*Bubalus bubalis*) oocytes. *Reprod. Dom.*

- 1 *Anim.* **49**, 343–51.
- 2 Queiroz Neta, L.B., Lira, G.P.O., Borges, A.A, Santos, M.V.O., Silva, M.B., Oliveira, L.R.M.,
3 Silva, A.R., Oliveira, M.F. & Pereira, A.F. (2018). Influence of storage time and nutrient
4 medium on recovery of fibroblast-like cells from refrigerated collared peccary (*Pecari tajacu*
5 Linnaeus, 1758) skin. *In vitro Cell Dev. Biol. Anim.* **54**, 486–95.
- 6 Rao, B.S., Mahesh, Y.U., Lakshmikantan, U.R., Suman, K., Charan, K.V. & Shivaji, S. (2010).
7 Developmental competence of oocytes recovered from *postmortem* ovaries of the endangered
8 Indian blackbuck (*Antilope cervicapra*). *J. Reprod. Dev.* **56**, 623–9.
- 9 Rho, G.J., Hahnel, A.C. & Betteridge, K.J. (2001). Comparison of oocyte maturation times and of
10 three methods of sperm preparation for their effects on the production of goat embryos *in vitro*.
11 *Theriogenology*. **56**, 503–16.
- 12 Ruiz, J., Landeo, L., Mendoza, J., Correa, J., Silva, M. & Ratto, M.H. (2015). *In vitro*
13 developmental competence of alpaca (*Vicugna pacos*) and llama (*Lama glama*) oocytes after
14 parthenogenetic activation. *Small Rumin. Res.* **133**, 148–52.
- 15 Saini, M., Selokar, N., Raja, A., Sahare, A., Singla, S., Chauhan, M., Manik, R. & Palta, P. (2015).
16 Effect of donor cell type on developmental competence, quality, gene expression, and
17 epigenetic status of interspecies cloned embryos produced using cells from wild buffalo and
18 oocytes from domestic buffalo. *Theriogenology*. **84**, 101–8.
- 19 Sansinena, M., Hylan, D., Hebert, K., Denniston, R. & Godke, R. (2005). Banteng (*Bos javanicus*)
20 embryos and pregnancies produced by interspecies nuclear transfer. *Theriogenology*. **63**,
21 1081–91.
- 22 Santos, M.L.T., Borges, A.A., Queiroz Neta, L.B., Santos, M.V.O., Oliveira, M.F., Silva, A.R. &
23 Pereira, A.F. (2016). *In vitro* culture of somatic cells derived from ear tissue of collared
24 peccary (*Pecari tajacu* Linnaeus, 1758) in medium with different requirements. *Pesq. Vet.*
25 *Bras.* **36**, 1194–202.
- 26 Saragusty, J., Diecke, S., Drukker, M., Durrant, B., Friedrich Ben-Nun, I., Galli, C., Göritz, F.,
27 Hayashi, K., Hermes, R. & Holtze, S. (2016). Rewinding the process of mammalian extinction.
28 *Zoo Biol.* **35**, 280–92.
- 29 Sharma, R., Sharma, H., Ahlawat, S., Aggarwal, R., Vij, P. & Tania, M. (2018). First attempt on
30 somatic cell cryopreservation of critically endangered *Camelus bactrianus* of India. *Gene*
31 *Reports*. **10**, 109–15.

- 1 Song, J., Hua, S., Song, K. & Zhang, Y. (2007). Culture, characteristics and chromosome
2 complement of Siberian tiger fibroblasts for nuclear transfer. *In vitro Cell. Dev. Biol. Anim.* **43**,
3 203–9.
- 4 Sparman, M.L., Tachibana, M. & Mitalipov, S.M. (2010). Cloning of non-human primates: the
5 road “less traveled by”. *Int. J. Dev. Biol.* **54**, 1671.
- 6 Sukparangsi, W., Bootsri, R., Sikeao, W., Karoon, S. & Thongphakdee, A. (2018). Establishment
7 of induced pluripotent stem cells from fishing cat and clouded leopard using integration-free
8 method for wildlife conservation. *Reprod. Fertil. Dev.* **30**, 230.
- 9 Tulaki, K., Yanagawa, Y., Takahashi, Y., Katagiri, S., Higaki, S., Koyama, K., Wang, X. & Li, H.
10 (2014). Effects of ovarian storage condition on *in vitro* maturation of Hokkaido sika deer
11 (*Cervus nippon yesoensis*) oocytes. *Jpn. J. Vet. Res.* **62**, 187–92.
- 12 Wang, L., Peng, T., Zhu, H., Lv, Z., Liu, T., Shuai, Z., Gao, H., Cai, T., Cao, X. & Wang, H.
13 (2007). *In vitro* development of reconstructed ibex (*Capra ibex*) embryos by nuclear transfer
14 using goat (*Capra hircus*) oocytes. *Small Rumin. Res.* **73**, 135–41.
- 15 Wani, N.A., Vettical, B.S. & Hong, S.B. (2017). First cloned Bactrian camel (*Camelus bactrianus*)
16 calf produced by interspecies somatic cell nuclear transfer: A step towards preserving the
17 critically endangered wild Bactrian camels. *PloS One.* **12**, e0177800.
- 18 Yamochi, T., Kida, Y., Oh, N., Ohta, S., Amano, T., Anzai, M., Kato, H., Kishigami, S., Mitani, T.
19 & Matsumoto, K. (2013). Development of interspecies cloned embryos reconstructed with
20 rabbit (*Oryctolagus cuniculus*) oocytes and cynomolgus monkey (*Macaca fascicularis*)
21 fibroblast cell nuclei. *Zygote* **21**, 358–66.
- 22 Yelisetti, U.M., Komjeti, S., Katari, V.C., Sisinthy, S. & Brahmasani, S.R. (2016). Interspecies
23 nuclear transfer using fibroblasts from leopard, tiger, and lion ear piece collected *postmortem*
24 as donor cells and rabbit oocytes as recipients. *In vitro Cell. Dev. Biol. Anim.* **52**, 632–45.
- 25 Yin, Y., Tang, L., Zhang, P., Kong, D., Wang, Z., Guan, J. & Li, Z. (2013). Optimizing the
26 conditions for *in vitro* maturation and artificial activation of sika deer (*Cervus nippon*
27 *hortulorum*) oocytes. *Reprod. Domest. Anim.* **48**, 27–32.
- 28 Zhao, Z.J., Ouyang, Y.C., Nan, C.L., Lei, Z.L., Song, X.F., Sun, Q.Y. & Chen, D.Y., (2006).
29 Rabbit oocyte cytoplasm supports development of nuclear transfer embryos derived from the
30 somatic cells of the camel and Tibetan antelope. *J. Reprod. Dev.* **52**, 449–59.
- 31 Zhu, H.Y., Kang, J.D., Li, S., Jin, J.X., Hong, Y., Jin, L., Guo, Q., Gao, Q.S., Yan, C.G. & Yin,

- 1 X.J. (2014). Production of rhesus monkey cloned embryos expressing monomeric red
- 2 fluorescent protein by interspecies somatic cell nuclear transfer. *Biochem. Biophys. Res.*
- 3 *Commun.* 638–43.
- 4

**CAPÍTULO 3 – ISOLATION, CHARACTERIZATION, AND CRYOPRESERVATION
OF COLLARED PECCARY SKIN-DERIVED FIBROBLAST CELL LINES**

(Artigo experimental)

PUBLICADO NA PEERJ

QUALIS: Quadriênio 2013-2016: Não classificado pela Medicina Veterinária/ Novo Qualis: A2

FATOR DE IMPACTO: 2,353

DOI: <https://doi.org/10.7717/peerj.9136>

Isolation, characterization, and cryopreservation of collared peccary skin-derived fibroblast cell lines

Alana Azevedo Borges¹, Gabriela Pereira de Oliveira Lira¹, Lucas Emanuel Nascimento¹, Maria Valéria de Oliveira Santos¹, Moacir Franco de Oliveira², Alexandre Rodrigues Silva³ and Alexsandra Fernandes Pereira¹

¹ Laboratory of Animal Biotechnology, Universidade Federal Rural do Semi-Árido, Mossoró, Rio Grande do Norte, Brazil

² Laboratory of Applied Animal Morphophysiology, Universidade Federal Rural do Semi-Árido, Mossoró, Rio Grande do Norte, Brazil

³ Laboratory of Animal Germplasm Conservation, Universidade Federal Rural do Semi-Árido, Mossoró, Rio Grande do Norte, Brazil

Corresponding Author:

Alexsandra F Pereira¹

Av. Francisco Mota, 572, Mossoro/RN, 59625 900, Brazil

Phone: +55 84 3317 8361

E-mail address: alexsandra.pereira@ufersa.edu.br

Running title: Peccary skin-derived fibroblast lines.

1 Abstract

2 **Background.** Biobanking of cell lines is a promising tool of support for wildlife conservation. In
3 particular, the ability to preserve fibroblast cell lines derived from collared peccaries is of
4 significance as these wild mammals are unique to the Americas and play a large role in
5 maintaining the ecosystem. We identified collared peccary fibroblasts by immunofluorescence
6 and evaluated their morphology, growth, and adherence capacity. Further, we monitored the
7 viability and metabolic activity of the fibroblasts to determine the effects of passage number and
8 cryopreservation on establishment of cell lines.

9
10 **Methods.** Skin biopsies were collected from the peripheral ear region from five adult animals in
11 captivity. Initially, cells were isolated from fragments and cultured in the Dulbecco's modified
12 Eagle medium (DMEM) supplemented with 10% fetal bovine serum and 2% antibiotic-
13 antimycotic solution under a controlled atmosphere (38.5°C, 5% CO₂). We evaluated the
14 maintenance of primary cells for morphology, adherence capacity of explants, explants in
15 subconfluence, cell growth, and absence of contamination. Moreover, we identified the fibroblast
16 cells by immunofluorescence. Additionally, to evaluate the influence of the number of passages
17 (first, third, and tenth passage) and cryopreservation on establishment of cell lines, fibroblasts
18 were analysed for the viability, metabolic activity, population doubling time (PDT), levels of
19 reactive oxygen species (ROS), and mitochondrial membrane potential ($\Delta\Psi_m$).

20
21 **Results.** All explants (20/20) adhered to the dish in 2.4 days $\pm$ 0.5 with growth around the
22 explants in 4.6 days $\pm$ 0.7, and subconfluence was observed within 7.8 days $\pm$ 1.0. Moreover, by
23 morphology and immunocytochemistry analyses, cells were identified as fibroblasts presenting
24 oval nuclei, a fusiform shape, and positive vimentin staining. No contamination was observed
25 after culture without antibiotics and antifungals for 30 days. While no difference was observed
26 for cell viability after the passages (first vs. third: $P = 0.98$; first vs. tenth: $P = 0.76$; third vs.
27 tenth: $P = 0.85$), metabolic activity was found to be reduced in the tenth passage ($23.2\% \pm$
28 12.1%), when compared to that in the first and third passage ($100.0\% \pm 24.4\%$, $P = 0.006$).
29 Moreover, the cryopreservation did not influence the viability ($P = 0.11$), metabolic activity ($P =$
30 0.77), or PDT ($P = 0.11$). Nevertheless, a greater $\Delta\Psi_m$ ($P = 0.0001$) was observed for the
31 cryopreserved cells (2.12 ± 0.14) when compared to that in the non-cryopreserved cells ($1.00 \pm$

0.05). Additionally, the cryopreserved cells showed greater levels of intracellular ROS after thawing (1.69 ± 0.38 vs. 1.00 ± 0.22 , $P = 0.04$).

Conclusions. This study is the first report on isolation, characterization, and cryopreservation of fibroblasts from collared peccaries. We showed that adherent cultures were efficient for obtaining fibroblasts, which can be used as donor cells for nuclei for species cloning and other applications.

Subjects: Cell Biology, Veterinary Medicine.

Keywords: Biological characterization, cellular conservation, cryobanking, wild mammals.

Introduction

Collared peccaries (*Pecari tajacu* Linnaeus, 1758) are wild mammals found only in the Americas and show a distribution from southern United States to northern Argentina, inhabiting the most diverse environments (Santos *et al.*, 2009). Currently, their population is considered to be stable (Gongora *et al.*, 2011); however, a significant reduction of their population has been seen in some biomes, such as the Caatinga (Desbiez *et al.*, 2012) and the Atlantic forest (Lazure *et al.*, 2010). As excellent seed dispersers (Redford 1992), they are very important for the maintenance of our ecosystem, whereas, economically, they have been commercialized for their meat and in leather production (Santos *et al.*, 2009). Scientifically, collared peccaries can be used as experimental models for closely related species such as the *Tayassu peccary* and *Catagonus wagneri* that have been listed as “vulnerable” in the IUCN Red List of Threatened Species (Keuroghlian *et al.*, 2013; Altrichter *et al.*, 2015).

In this sense, studies related to the conservation of the collared peccary have been intensified, especially aimed at improving the techniques related to the preservation of somatic samples. Using this study, we established a culture condition for explants derived from the skin of adult collared peccaries (Santos *et al.*, 2016) and developed a protocol for cryopreservation (Borges *et al.*, 2017; Borges *et al.*, 2018a; Borges *et al.*, 2018b) and refrigeration of these explants (Queiroz Neta *et al.*, 2018). In order to conduct the cloning experiments on this species by a somatic cell nuclear transfer, as well as to produce induced pluripotent cells, it is necessary to establish properly characterized cell lines.

In general, as observed in other mammals (*Guan et al., 2010; Kwong et al., 2014*), establishment of an adequate cell line is a prerequisite step for the success of cloning and producing induced pluripotent cells (*Borges & Pereira 2019*). For these techniques, fibroblasts and epithelial cells derived from the skin have been widely used (*Jyotsana et al., 2016; Siengdee et al., 2018*). Initially, epithelial and fibroblast cells were grown simultaneously; nevertheless, fibroblasts can more easily adhere as well as detach by trypsinization as compared to the epithelial cells (*Bai et al., 2012; Saadeldin et al., 2019; Siengdee et al., 2018*). In these methods, the culture after the second passage has been considered to contain mainly fibroblasts (*Mehrabani et al., 2014*).

Additionally, for the confirmation of a fibroblast line, it is necessary to verify the possible changes that occur in these cells during culture (*Guan et al., 2010; Song et al., 2007*) and cryopreservation (*Magalhães et al., 2017*). In general, the number of passages throughout an *in vitro* study can modify the cellular epigenetic state, affecting the embryonic development after cloning (*Rodriguez-Osorio et al., 2012; Trokovic et al., 2015*). *Magalhães et al. (2017)* observed a reduced viability and metabolic activity in the cells derived from the skin of the brown brocket deer in the tenth passage. Thus, the establishment of a cell line ensures a complete knowledge of the parameters that confer quality to the nucleus of the donor cell, named the karyoplast (*Guan et al., 2010*). Moreover, identification of the damages occurring during cryopreservation is essential for establishment of a cell line. Cryo-variables may affect several cellular processes, including survival, functionality, and the cytoskeleton, which may compromise the reprogramming ability of the karyoplasts (*Chatterjee et al., 2017*). Therefore, we aimed to isolate, characterize, and cryopreserve the fibroblast cells derived from the skin of the ear of collared peccaries for their future application in cloning strategies by a somatic cell nuclear transfer and production of induced pluripotent cells.

Materials & Methods

Chemicals and media

The Dulbecco's modified Eagle medium (DMEM), fetal bovine serum (FBS), penicillin, streptomycin, and amphotericin solutions were obtained from Gibco-BRL (Carlsbad, USA). Fluorescent probes were purchased from Invitrogen (Carlsbad, USA). Anti-vimentin antibody

and goat anti-mouse IgG (Alexa Fluor® 488) were purchased from Abcam (Cambridge, USA). The other chemicals were obtained from Sigma-Aldrich (St. Louis, USA). Media were filtered using a 0.22-µm system (Corning, New York, USA) and adjusted to pH of 7.2–7.4.

Bioethics and animals

This study was approved by the Ethics Committee of Animal Use of the Federal Rural University of Semi-Arid (CEUA/UFERSA, no. 23091.001072/2015-92) and the Chico Mendes Institute for Biodiversity Conservation (ICMBio, no. 48633-2). All animals belonged to the Centre of Multiplication of Wild Animals (CEMAS/UFERSA, Mossoró, RN, Brazil, 5°10'S, 37°10'W), registered at the Brazilian Institute of Environment and Renewable Natural Resources (IBAMA) as a scientific breeding site (no. 1478912). The breeder stocks 100 collared peccaries on an average, and for this research four females and one male at ages of 26.8 months ± 2.9 months were used.

Ear tissue explant collection and primary culture

Peripheral skin (1 cm²–2 cm²) was recovered from the ear sections used to identify collared peccaries kept in captivity. After the collection, a trichotomy of the tissue followed by a sterilization with 70% alcohol was performed. Samples were transported to the laboratory in DMEM supplemented with 2% antibiotic-antimycotic solution (10,000 units/mL penicillin, 10,000 µg/mL streptomycin, and 25 µg/mL amphotericin B) at 37°C within 30 min.

In the laboratory, fragments (9.0 mm³) were washed sequentially under laminar flow in the following media: 1) DMEM supplemented with 10% FBS, and 10% antibiotic-antimycotic solution; 2) alcohol; and 3) DMEM plus 10% FBS, and 2% antibiotic-antimycotic solution. Then, the samples were fragmented (four fragments per animal) and placed in polystyrene culture dishes treated for cell adhesion with the latter medium for cell culture. The skin was cultured at 38.5°C under a controlled environment with 5% CO₂ and 95% air, according to a method described by *Santos et al. (2016)*.

Evaluation of the somatic cells in primary cultures and subcultures

During primary culture, the medium was changed every 24 h. For evaluation of the somatic cells, the primary culture was analyzed before reaching confluency and until it reached a confluency of

70%–80%. Using an inverted microscope (Nikon TS100, Tokyo, Japan), the cells were evaluated for the following parameters: cell morphology, number of adhered samples, number of samples, evident subconfluency, day of sample adherence, day of subconfluent growth of the samples, and total time to reach 70%–80% confluence (*Borges et al., 2017*).

When the cells reached 70%–80% subconfluency, they were subcultured and distributed for other analyses. The 70%–80% subconfluence was defined as the stage when 70%–80% of the culture dishes consisted of somatic cells (*Santos et al., 2016*). Subconfluent cells were washed with PBS then trypsinized with a trypsin/EDTA solution (0.25%/0.2%) for 7 min and centrifuged at 600 x g for 10 min. The supernatant was removed, the cell pellet was resuspended in culture medium, and the cell suspension was transferred to another dish for subculturing (*Borges et al., 2018b*). The medium was replaced with a fresh medium every other day and the cells were monitored daily. With the successful passaging of the cultures, the cells are considered a cell line, following the convention of the Society of In Vitro Biology (*Schaeffer, 1990*). The cell line was designated as Ptskf.

Thus, in addition to an evaluation of the maintenance of cells in the primary culture, the subcultured cells were initially evaluated for the confirmation of fibroblasts using morphology and immunofluorescence analyses. Moreover, the possibility of contamination was also evaluated. Subsequently, the influence of the number of passages (first, third, and tenth passage) and the metabolic activity of the cells were analyzed by a viability assay using trypan blue and the 3-(4, 5-dimethylthiazolyl-2)-2, 5-diphenyltetrazolium bromide (MTT) assay, respectively. Moreover, the cells were also evaluated for the effects of a slow freezing cryopreservation. Other than the above-mentioned tests, growth dynamics by quantification of the population doubling time (PDT), oxidative stress analysis for quantification of intracellular reactive oxygen species (ROS) levels using the fluorescent probe 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA), and assessment of the mitochondrial membrane potential ($\Delta\Psi_m$) using the fluorescent probe MitoTrackerRed® were performed.

Morphological characterization of the fibroblasts

Morphological characteristics were observed throughout the in vitro culture under light microscopy for cellular and nuclear shapes and cytoplasmic extensions.

Vimentin immunofluorescence

For a morphological confirmation, the cells were subjected to an immunocytochemistry protocol based on the method described by *Amoli et al. (2017)*. Briefly, the cells were fixed using 4% paraformaldehyde for 10 min at 25°C, then washed with chilled PBS. Subsequently, cells were incubated with an antigen-retrieval buffer (100 mM Tris, 5% urea, pH 9.5), and then permeabilized for 1 h in 0.4% Triton X-100. Afterwards, the cells were incubated in 0.1% Tween-20 for 1 h to block non-specific binding of the antibodies. Finally, the cells were immunostained with mouse anti-vimentin antibody (ab8979, 1:200) for 24 h at 4°C, and, then incubated with the secondary antibody (goat anti-mouse IgG, Alexa Fluor® 488, ab150113, 1:400) for 1 h at 25°C in the dark. Cells were counter-marked with 1 µg/mL Hoechst for 1 min and observed under a fluorescence microscope (Olympus BX51TF, Tokyo, Japan).

Confirmation of the absence of bacterial and fungal contamination

Cells of the third passage were cultured for 30 days in DMEM containing 10% FBS in the absence of an antibiotic-antimycotic solution, at 38.5°C, 5% CO₂ and 95% air. Daily evaluation was performed under light microscopy for the identification of bacterial and fungal contamination.

Influence of the passage number on the quality of fibroblast lines

Initially, the fibroblast cells were analyzed for the effect of the number of passages (first, third, and tenth passage) by a viability assay using trypan blue, according to the method described by *Magalhães et al. (2017)*. We evaluated these three cell passages specifically because both fibroblast and epithelial cells were present in the initial (first) passage, only fibroblasts were visualized at the third passage onwards, and the cells of the tenth passage were used for most of the production of embryonic clones (*Shiga et al., 1999; Kubota et al., 2000*). The evaluations were performed in triplicate for each animal for each passage.

Briefly, the cells were stained with 0.4% trypan blue in PBS and counted on a hemocytometer. Subsequently, the cells were also analyzed for a metabolic activity using the MTT assay, according to the method described by *Borges et al. (2018b)*. A concentration of 5.0×10^4 cells/mL from the first, third, and tenth passages was grown in 12-well polystyrene plates treated for cell adhesion. After 5 days, 1.5 mL of the MTT solution (5 mg/mL in DMEM) was added and the polystyrene culture dishes treated for cell adhesion were incubated for 3 h. The MTT solution was then removed and 1.0 mL of dimethyl sulfoxide (DMSO) was added for 5 min under slow stirring to solubilize the MTT. After the total dissolution of formazan crystals, samples were analyzed in a spectrophotometer (Shimadzu® UV-mini-1240, Kyoto, Japan) at an absorbance wavelength of 595 nm. The evaluations were performed in triplicate for each animal for each passage.

Influence of cryopreservation on the quality of fibroblast lines

To evaluate the effect of cryopreservation on the quality of fibroblast lines, cells of the third passage of the five animals were subjected to slow freezing in the freezing medium (DMEM supplemented with 10% DMSO as a permeating cryoprotectant and 10% FBS and 0.2 M sucrose as non-permeating cryoprotectants). Cells at a concentration of 5.0×10^4 cells/mL were first exposed to DMSO-FBS solution for 15 min at 4°C, then sucrose solution was added followed by an additional incubation for 15 min at 4°C. The cryovials containing 1.0 mL of cells in the freezing medium were cooled in a Mr. Frosty freezing container (Thermo Scientific, Waltham, USA) at a cooling rate of 1°C/min, and later stored in a freezer at -80°C, reaching -70°C overnight before being transferred into liquid nitrogen (*León-Quinto et al., 2014*).

For thawing, the cryovials were exposed for 1 min at 25°C and immersed in a water bath at 37°C for 3 min–4 min. Then, the cell contents were removed from the cryovials and washed to remove the cryoprotectants. Initially, the first wash was performed with DMEM and 10% FBS containing 0.2 M sucrose at 4°C for 15 min and centrifuged. Subsequently, the second wash was performed using only DMEM and 10% FBS, maintained at 25°C for 15 min, centrifuged, and the cells were recovered for the evaluations as per a method described previously (*Santos et al., 2016*).

After thawing, the non-cryopreserved and the cryopreserved cells were evaluated for growth dynamics by quantification of PDT. The evaluations were performed in triplicate for each animal. Briefly, the growth kinetics was studied for nine days using 3.0×10^4 cells/mL, and cells were counted daily to determine the number of growing cells. Data on the cell growth and density were monitored and recorded, mean values of which were used to plot a growth curve and calculate PDT (Roth 2006) using the following formula:

$$PDT = T \ln 2 / \ln (X_e / X_b)$$
 where PDT is the time of the culture (in hours), T is the incubation time, X_b is the number of cells at the beginning of the time incubation, X_e is the number of cells at the end of the incubation time, and $\ln$ is the Napierian logarithm.

Moreover, for evaluation of an oxidative stress by quantification of the intracellular ROS levels, cells were stained with the fluorescent probe H_2DCFDA , according to a method described by Santos *et al.* (2019). Thawed cells were washed with PBS and placed into polystyrene culture dishes treated for cell adhesion containing 500 μ L of 5 μ M H_2DCFDA . The cells obtained after a 70% confluency were incubated at 38.5°C in 5% CO_2 for 30 min. Stained cells were washed with PBS, placed on glass slides, photographed under a fluorescence microscope (Olympus BX51TF, Tokyo, Japan), and fluorescence signal intensity (pixels) was quantified. Ten images (two/animal) obtained were evaluated using the ImageJ software (version 1.49v, Java 1.8.0_201, Wayne Rasband, U.S. National Institutes of Health, Bethesda, MD, USA; website: <http://rsb.info.nih.gov/ij/download.html>). The background signal intensity was subtracted from the values obtained for the treated samples. Measured mean value of the micrograph for the non-cryopreserved cells was taken as a calibrator. Relative expression levels (arbitrary fluorescence units) were generated by dividing the measured value of each micrograph for the cryopreserved cells by the mean of the calibrator.

Finally, for the assessment of $\Delta\Psi_m$, cells were stained using 500 nM of the fluorescent probe MitoTracker Red® (CMXRos), according to a method described by Santos *et al.* (2019). The procedure, incubation, and evaluation of the ten images (two/animal) were performed as described for the quantification of ROS.

Statistical analysis

All data have been expressed as the mean $\pm$ standard error (one animal/one repetition) and were analyzed using the StatView 5.0 software (Graph-Pad Software Incorporation, La Jolla, USA). Normality of all results was verified by the Shapiro-Wilk test and homoscedasticity was verified by the Levene's test. ROS levels, $\Delta\Psi_m$, viability, and metabolic activity were altered with arcsine and analysed by variance analysis (ANOVA) followed by the Tukey's test. PDT was compared with ANOVA followed by the unpaired t-test. Statistical significance was set at $P < 0.05$.

Results

Evaluation of the somatic cells in the primary cultures and subcultures

The total culture time was 95 days with an evaluation of cells until the tenth passage. The adhesion of the fragments (Fig. 1A), detachment of cells (Fig. 1B-1C), and proliferative capacity were observed in all the explants until reaching a confluence (and later, a subconfluence) around the adhered fragments (Fig. 1D-1F, Table 1). All explants had adhesion ability and reached subconfluence. Number of days for each explant to reach a 100% tissue adherence (2.4 days $\pm$ 0.5 days), to grow around the explants (4.6 days $\pm$ 0.7 day), and to reach subconfluence (7.8 days $\pm$ 1.0 day), were different.

Morphological characterization of the fibroblasts

In cultures, monolayers of cells with a fibroblast-like morphology were observed (Fig. 1E). The cells had an oval nuclei and extensions with a fusiform shape, showing rapid growth that replaced the epithelial cells.




Figure 1 Outgrowth of fibroblast cells from ear skin samples of collared peccaries. Exhibit a skin explant cultured in A) day 1, B) day 3 and C) day 5 of primary culture and exhibit a fibroblast population cultured in D) day 15 and E) day 19 of subculture. F) exhibit cells after the trypsinization process. Scale bar = 100 μm .

1 **Table 1 Establishment of primary culture and subcultures of somatic cells derived from**
 2 **collared peccary ear skin.**

Animal	No. samples			No. attached samples		
	Initial	Attached (%)	Day all attached explants	Grow subconfluence (%)	to Day cell grow explants	all Subconfluence total time (days)
F1	4	100	2	100	4	6
F2	4	100	2	100	4	5
F3	4	100	1	100	3	9
F4	4	100	3	100	5	10
M1	4	100	4	100	7	9
Mean	± 20	100	2.4 ± 0.5	100	4.6 ± 0.7	7.8 ± 1.0
S.E						

Vimentin immunofluorescence

Morphology of the fibroblast-like cells in the initial culture was observed by light microscopy, which was further confirmed for the cell type identification as vimentin-labeled fibroblasts under fluorescence microscopy (Fig. 2A-2F). Cells exhibited a high expression of vimentin that marked the cytoplasm completely, and the spindle-like shape and ovoid nucleus was highlighted by the Hoechst labeling. Therefore, the identification of a fibroblast cell was evident.




Figure 2 Immunocytochemical detection of vimentin protein for identification of collared peccary fibroblasts. A-D) cells stained with vimentin antibody. B-E) nucleus of cells stained by Hoechst. C-F) merged vimentin (green) and Hoechst (blue). a-b-c (x5), d-e-f (x10). Scale bar = 10 μm .

Confirmation of the absence of bacterial and fungal contamination

No sign of contamination (turbidity, colony, or hyphal growth) was observed for 30 days in the culture without antibiotics and antifungals. The culture medium did not show any change in the appearance when observed under a light microscope. We did not observe turbidity or any specific odor. In addition, there was no change in the biological characteristics of growth and proliferation indicating a complete absence of contamination.

Influence of the passage number on the quality of fibroblast lines


No significant difference was observed in the cell viability (74.5% to 84.4%) when evaluated by trypan blue staining after the passages (first vs. third: $P = 0.98$; first vs. tenth: $P = 0.76$; third vs. tenth: $P = 0.85$). However, the metabolic activity was reduced in the tenth passage ($23.2\% \pm 12.1\%$) as compared to that of the first and third passages ($100.0\% \pm 24.4\%$, $P = 0.006$).

Influence of cryopreservation on the quality of fibroblast lines

Cryopreservation did not affect the viability when evaluated by trypan blue staining ($87.4\% \pm 0.3\%$ vs. $74.0\% \pm 5.9\%$, $P = 0.11$). Moreover, after two passages of the thawed cells, the viability was $86.4\% \pm 3.2\%$. In addition, no difference ($P = 0.77$) was observed for the metabolic activity between the cryopreserved ($85.2\% \pm 10.0\%$) and the non-cryopreserved cells ($100.0\% \pm 36.4\%$).

Moreover, the cryopreserved and the non-cryopreserved cells were compared for growth dynamics (Fig. 3). The growth curve of both groups showed a typical “S-shaped” pattern from the nine-day culture of cells. The latency time was two days, followed by an exponential phase until the fourth day, the stationary phase until the seventh day, and the plateau phase from day eight (Fig. 3A). No difference was observed for the PDT values of the cryopreserved and the non-cryopreserved cells ($P = 0.11$, Fig. 3B).

Nevertheless, the cryopreserved cells showed greater levels of intracellular ROS (Fig. 4A) in arbitrary fluorescence units when compared to that of the non-cryopreserved cells (1.69 ± 0.38 vs. 1.00 ± 0.22 , $P = 0.04$) (Fig. 4C). In addition, an alteration in the $\Delta\Psi_m$ (Fig. 4B) in arbitrary fluorescence units ($P = 0.0001$) was observed for the cryopreserved cells (2.37 ± 0.07) when compared to that of the non-cryopreserved cells (1.00 ± 0.05) (Fig. 4D).



1 **Figure 3 The growth dynamics of cryopreserved and non-cryopreserved cells derived from collared peccary skin.** A) growth
 2 curves of cryopreserved and non-cryopreserved cells. B) values of population doubling time (PDT) after culture for nine days.
 3




Figure 4 Evaluation of intracellular reactive oxygen species (ROS) and mitochondrial membrane potential ($\Delta\Psi_m$). Cell stained using fluorescent probe A) 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA) and B) MitoTrackerRed® (CMXRos) (x10). Quantification of C) ROS and D) $\Delta\Psi_m$ levels. Scale bar = 10 μ m. (*) Indicate statistical difference (P < 0.05).

Discussion

In this study, we isolated, characterized, and cryopreserved the fibroblast cells derived from the skin of collared peccaries. Moreover, we established the fibroblast cell lines of these animals with an aim to use these cells in cloning experiments by a somatic cell nuclear transfer in the future. The cell line can be considered as the first constituent of the peccary invitrome and a resource for future studies in many disciplines (Barioch 2018; Bols *et al.*, 2017). Thus, the ear tissues of collared peccaries can be isolated and grown into fibroblasts in an adherent culture for establishment of cell lines and development of a cryobank. The development of these somatic cell banks has been increasing in the interest of conserving genetic samples of wild mammals to preserve valuable species, and as sources for biological research (León-Quinto *et al.*, 2009; Mehrabani *et al.*, 2014; Saadeldin *et al.*, 2019; Siengdee *et al.*, 2018).

All explants adhered to the flask surface within 2 days to 4 days, with cellular growth around the explant within 3 days, and demonstrating confluency within 5 days to 10 days after a culture initiation. These characteristics of explants during *in vitro* culture were similar to the explants derived from other domestic and wild mammals. In studies using tissues from horses, the migration of fibroblast and epithelial-like cells from explants have been observed after 5 days to 7 days of an *in vitro* culture (Amoli *et al.*, 2017). In the case of goat-derived tissues, the explants reportedly adhered to the flasks within 5 days to 7 days and the cells became confluent within 3 days to 5 days post adhesion (Bai *et al.*, 2012). In the Iranian Sistani cattle-derived tissues, the explants adhered to the culture flasks within 7 days to 14 days and were observed to allow the growth of fibroblast-like cells from the margins of explants (Gorji *et al.*, 2017).

For the Luxi cattle-derived tissues, fibroblast-like or epithelial-like cells could be seen migrating from the tissues within 5 days to 12 days post adhesion (Liu *et al.*, 2008). In the tissues derived from wild camels, fibroblast-like or epithelial-like cells could be seen migrating from the sides of explants within 8 days to 10 days post adhesion (Sharma *et al.*, 2018). In tissues derived from the domestic porcine, a species phylogenetically close to the collared peccaries, all the explants adhered within 3 days to 8 days (Silvestre, Sánchez & Gómez, 2004). The similarity among these data can be related to the culture medium because in a majority of these studies, DMEM containing FBS, antibiotic, and antimycotic solution was used (Magalhães *et al.*, 2017; Saadeldin *et al.*, 2019; Siengdee *et al.*, 2018). Since

1 primary culture needs to mimic the *in vivo* environment of the cells (Guo *et al.*, 2018), we
2 observed previously (Santos *et al.*, 2016) that the medium for growth of somatic cells derived
3 from collared peccaries was DMEM with 10% FBS and 2% antibiotic-antimycotic solution.

4
5 We showed that ear explant cultures obtained from the tissues of the collared peccaries were
6 efficient for cell recovery allowing a culture for 95 days and up to ten passages. Moreover, we
7 confirmed that the ear skin was a common source where fibroblasts cells could be separated
8 and eventually used as karyoplasts for cloning purposes (Luo *et al.*, 2014). These cells were
9 identified by vimentin, an intermediate filament that indicates the mesenchymal origin of
10 endothelial and fibroblast cells (Yajing *et al.*, 2018). Initially, during *in vitro* culture, epithelial
11 and fibroblastic cells grew simultaneously. However, fibroblasts can be trypsinized more
12 rapidly and adhere more easily as compared to the epithelial cells (Bai *et al.*, 2012; Saadeldin
13 *et al.*, 2019). Therefore, in this work, cells from the third passage were confirmed as
14 fibroblasts by morphology and immunofluorescence analyses.

15
16 The clear medium observed during the 30-day assay demonstrated the ability to allow the
17 growth of the culture without any biological contamination. The propagation of fungi causes
18 turbidity by accumulation of their metabolites. In addition, the colonies can be seen under a
19 light microscope, or sometimes with the naked eye (Li *et al.*, 2009). Bacterial contamination
20 can also be identified by the naked eye as turbidity. One simple way to avoid this
21 contamination is to filter the culture medium (Bai *et al.*, 2012). Sources of contamination may
22 include, but are not limited to, the equipment, air, culture medium, serum, and explant.
23 Therefore, microbial contaminations are quite frequent in cell culture (Bai *et al.*, 2012). Thus,
24 the use of antibiotic and antimycotic combined with careful handling, is essential to ensure the
25 absence of contamination.

26
27 After two passages, it was possible to separate fibroblast cells from other primary cells
28 because different cell types exhibit different cellular behavior upon trypsinization. The
29 fibroblast cells detach quicker than epithelial cells (Gorji *et al.*, 2017). However, fibroblasts
30 detach in response to trypsin more rapidly than epithelial cells and adhere more quickly (Bai
31 *et al.*, 2012). In wild camels, the initial lag phase of 48 h representing the adaptation of
32 fibroblasts and recovery from a protease damage is followed by the exponential phase
33 (Sharma *et al.*, 2018). In collared peccaries, the replication of cells begins to slow down after
34 seven days because of contact inhibition, which, in wild camels, has been observed after six

1 days (*Sharma et al., 2018*). No difference in the cell viability was observed among the first,
2 third, and the tenth passage, corroborating with the studies that used cells from these passages
3 for production of competent cloned embryos (*Shiga et al., 1999; Kubota et al., 2000*).

4
5 In contrast, through the metabolic activity test evaluated by the formation of formazan
6 crystals, a significant reduction in the metabolic activity at the tenth passage was observed,
7 indicating a reduced cellular functionality. Similar behavior was observed in cells from the
8 brown brocket deer in which the metabolic activity measured by the MTT assay showed
9 significantly lower values in the tenth passage than the values in the fourth passage
10 (*Magalhães et al., 2017*). Therefore, the number of passages can reduce the metabolic activity
11 rate and cell proliferation, thereby conserving cells of the early passages (*Li et al., 2009*).
12 After several passages, genetic characteristics of the cells can be modified by culture
13 conditions; hence, a minimum number of passages have been recommended to conserve the
14 cellular characteristics (*Mehrabani et al., 2014*). Owing to this reason, the cells were
15 cryopreserved in the third passage for the conservation of the somatic germplasm of collared
16 peccaries.

17
18 The cell survival rate after thawing is the most commonly used criteria to evaluate the success
19 of a cryopreservation (*Chatterjee et al., 2017*). The cellular viability and the functional
20 metabolic activity of the cells were maintained after thawing the fibroblasts isolated from the
21 collared peccaries. This factor demonstrates that optimal *in vitro* culture conditions
22 significantly influence the recovery from cellular damages caused by the freezing process
23 (*Gorji et al., 2017*). As for the growth curve, cryopreserved cells presented a very similar
24 profile to that of the non-cryopreserved cells, showing their normal proliferation capacity
25 regardless of the cryopreservation process. The establishment of somatic cell banks using
26 cryopreservation technology is an easy and effective approach towards storing the genetic
27 information of diverse species (*Li et al., 2009*). However, the cells should be handled with the
28 utmost care during cryopreservation to maintain a high-quality cell bank in the long term
29 (*Mehrabani et al., 2014*).

30
31 Moreover, epigenetic alterations, such as DNA fragmentation, free radical accumulation,
32 ionic imbalances, apoptosis, biochemical alterations, DNA methylation, and histone
33 modification can be a result of the cryopreservation (*Chatterjee et al., 2017*). These after-
34 effects of cryopreservation may have caused mitochondrial structural abnormalities, thereby

promoting an increased ROS production and H₂O₂ content, increased lipid peroxidation, and increased expression of autophagic proteins harbored by the cells (*Mata et al., 2012*). A failure in the mitochondrial membrane potential is a hallmark of apoptosis, leading to the collapse of the organelle and release of cytochrome-C into the cytoplasm, and ultimately activation of the apoptotic cascade (*Magalhães et al., 2012*). Moreover, a high $\Delta\Psi_m$ mitochondrial respiratory chain becomes a significant ROS producer (*Korshunov, Skulachev, & Starkov., 1997*). Therefore, a higher $\Delta\Psi_m$ in cryopreserved cells can be linked mainly to an increase in the oxidative stress.

Finally, parameters like cryovariables, including cooling and thawing rates, type and concentration of the cryoprotectant, cell type and shape, and nucleation temperature may affect the success of cryopreservation (*Chatterjee et al., 2017*). This suggests that the optimization of related cryopreservation methods for the collared peccary fibroblasts to minimize an altered $\Delta\Psi_m$ and increased levels of intracellular ROS production is essential.

Conclusions

To our knowledge, this study is the very first report on a successful isolation, characterization, and cryopreservation of fibroblast lines derived from adult collared peccaries (P_{tskf}). We showed that the adherent culture was efficient for obtaining fibroblasts, which can be used as donor cells for nuclei for cloning of this species. Moreover, it was possible to maintain the viability of the cells until the tenth passage. In addition, cryopreservation did not affect the viability, metabolic activity, and proliferative activity of the fibroblasts after slow freezing. However, cryopreservation altered the ROS levels and $\Delta\Psi_m$, indicating necessary optimization of the cryopreservation protocol. Lastly, the establishment of fibroblast cell lines derived from collared peccaries may be a source of experimental models for many biological studies such as nuclear reprogramming and animal cloning.

Acknowledgements

The authors thank the Centre for Wild Animals Multiplication (CEMAS/UFERSA) for providing the animals, and the Laboratory Biochemistry and Molecular Biology (BIOMOL/UERN) for technical assistance. This study was supported by the Brazilian Council of Scientific Development (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES, Financial Code 001).

References

- Altrichter, M., Taber, A., Noss, A., Maffei, L., Campo, J. 2015.** *Catagonuswagneri*. The IUCN Red list of threatened species e.T4015A72587993. DOI: 10.2305/IUCN.UK.2015-2.RLTS.T4015A72587993.en. (accessed 07 October 2019).
- Amoli AD., Mohebbali N., Farzaneh P., Fazeli SAS., Nikfarjam L., Movasagh SA., Moradmamand Z., Ganjibakhsh M., Nasimian A., Izadpanah M. 2017.** Establishment and characterization of Caspian horse fibroblast cell bank in Iran. *In vitro Cellular & Developmental Biology-Animal* **53**:337–343. DOI: 10.1007/s11626-016-0120-3.
- Bai C., Wang D., Su X., Zhang M., Guan W., Ma Y. 2012.** Establishment and biological research of the Jining Grey goat fibroblast line. *Turkish Journal of Veterinary and Animal Sciences* **36**:659–667. DOI: 10.3906/vet-1106-15.
- Barioch A. 2018.** The Cellosaurus: a cell knowledge resource. *Journal of Biomolecular Techniques* **29**:25–38.
- Bols NC., Pham PH., Dayeh VR., Lee LEJ. 2017.** Invitromatics, invitrome, and invitroomics: Introduction of three new terms for *in vitro* biology and illustration of their use with the cell lines from rainbow trout. *In vitro Cellular & Developmental Biology-Animal* **53**:383–405. DOI: 10.1007/s11626-017-0142-5
- Borges AA., Lima GL., Queiroz Neta LB., Santos MVO., Oliveira MF., Silva AR., Pereira AF. 2017a.** Conservation of somatic tissue derived from collared peccaries (*Pecari tajacu* Linnaeus, 1758) using direct or solid-surface vitrification techniques. *Cytotechnology* **69**:643–654. DOI: 10.1007/s10616-017-0074-7
- Borges AA., Lira GPO., Nascimento LE., Queiroz Neta LB., Santos MVO., Oliveira MF., Silva AR., Pereira AF. 2018a.** Influence of cryopreservation solution on the *in vitro* culture of skin tissues derived from collared peccary (*Pecari tajacu* Linnaeus, 1758). *Biopreservation and Biobanking* **16**:77–81. DOI: 10.1089/bio.2017.0090.
- Borges AA., Queiroz Neta LB., Santos MVO., Oliveira MF., Silva AR., Pereira AF. 2018b.** Combination of ethylene glycol with sucrose increases survival rate after vitrification of somatic tissue of collared peccaries (*Pecari tajacu* Linnaeus, 1758). *Pesquisa Veterinária Brasileira* **38**:350–356. DOI: 10.1590/1678-5150-PVB-5193.
- Borges, AA., Pereira, AF. 2019.** Potential role of intraspecific and interspecific cloning in the conservation of wild mammals. *Zygote* **27**:111–117. doi: 10.1017/S0967199419000170.
- Chatterjee A., Saha D., Niemann H., Gryshkov O., Glasmacher B., Hofmann N. 2017.** Effects of cryopreservation on the epigenetic profile of cells. *Cryobiology* **74**:1–7. DOI: 10.1016/j.cryobiol.2016.12.002.

- 1 **Desbiez ALJ., Keuroghlian A., Mello Beisiegel B., Medici EP., Gatti A., Pontes ARM.,**
2 **Campos CB., Tófoli CF., Júnior EAM., Azevedo FC. 2012.** Avaliação do risco de
3 extinção do cateto *Pecari tajacu* Linnaeus, 1758, no Brasil. *Biodiversida de Brasileira* **3**:74–
4 83.
- 5 **Gongora, J., Reyna-Hurtado, R., Beck, H., Taber, A., Altrichter, M., Keuroghlian, A.,**
6 **2011.** *Pecaritajacu*. The IUCN Red List of Threatened Species e.T41777A10562361.
7 <http://dx.doi.org/10.2305/IUCN.UK.20112.RLTS.T41777A10562361.en>. (accessed on 26
8 May 2019).
- 9 **Gorji ZE., Khaledi KJ., Amoli AD., Ganjibakhsh M., Nasimian A., Gohari NS.,**
10 **Izadpanah M., Vakhshiteh F., Farghadan M., Moghanjoghi SM. 2017.** Establishment
11 and characteristics of Iranian Sistani cattle fibroblast bank: a way to genetic conservation.
12 *Conservation Genetics Resources* **9**:305–312. DOI: 10.1007/s12686-016-0640-x.
- 13 **Guan W., Liu C., Li C., Liu D., Zhang W., Ma Y. 2010.** Establishment and cryopreservation
14 of a fibroblast cell line derived from Bengal tiger (*Panthera tigris tigris*). *Cryoletters*
15 **31**:130–138.
- 16 **Guo H, Xing Y, Zhang Y, He L, Deng F, Ma X, Li Y. 2018.** Establishment of an
17 immortalized mouse dermal papilla cell strain with optimized culture strategy. *PeerJ*
18 **6**:e4306 <https://doi.org/10.7717/peerj.4306>
- 19 **Jyotsana B., Sahare AA., Raja AK., Singh KP., Nala N., Singla S., Chauhan M., Manik R.,**
20 **Palta P. 2016.** Use of peripheral blood for production of buffalo (*Bubalus bubalis*) embryos
21 by handmade cloning. *Theriogenology* **86**:1318–1324. DOI:
22 10.1016/j.theriogenology.2016.04.073.
- 23 **Keuroghlian, A., Desbiez, R., Reyna-Hurtado, M., Altrichter, H., Beck, A., Taber, J.M.V.**
24 **2013.** *Tayassupecari*. IUCN Red List of Threatened Species 2013.1.
25 <http://dx.doi.org/10.2305/IUCN.UK.2013-1.RLTS.T41778A44051115.en>. (accessed 07
26 October 2019).
- 27 **Korshunov, SS., Skulachev, VP., Starkov, AA. 1997.** High protonic potential actuates a
28 mechanism of production of reactive oxygen species in mitochondria. *FEBS Lett* **416**:15–28.
29 doi: 10.1016/s0014-5793(97)01159-9
- 30 **Kubota, C., Yamakuchi, H., Todoroki, J., Mizoshita, K., Tabara, N., Barber, M., Yang, X.**
31 **2000.** Six cloned calves produced from adult fibroblast cells after long-term culture. *PNAS*
32 **97**:990–995. doi: 10.1073/pnas.97.3.990
- 33 **Kwong P., Nam H., Wan Khadijah W., Kamarul T., Abdullah R. 2014.** Comparison of *in*
34 *vitro* developmental competence of cloned caprine embryos using donor karyoplasts from

- 1 adult bone marrow mesenchymal stem cells vs. ear fibroblast cells. *Reproduction in*
2 *Domestic Animals* **49**:249–253. DOI: 10.1111/rda.12262.
- 3 **Lazure L., Bachand M., Anseau C., Almeida-Cortez J. 2010.** Fate of native and introduced
4 seeds consumed by captive white-lipped and collared peccaries (*Tayassupecari*, Link 1795
5 and *Pecaritajacu*, Linnaeus 1758) in the Atlantic rainforest, Brazil. *Brazilian Journal of*
6 *Biology* **70**:47–53. DOI: 10.1590/S1519-69842010000100008.
- 7 **León-Quinto T., Simón MA., Cadenas R., Martínez Á., Serna A. 2014.** Different
8 cryopreservation requirements in foetal versus adult skin cells from an endangered mammal,
9 the Iberian lynx (*Lynx pardinus*). *Cryobiology* **68**:227–233. DOI:
10 10.1016/j.cryobiol.2014.02.001.
- 11 **León-Quinto, T., Simón, MA., Cadenas, R., Jones, J., Martínez Hernandez, FJ., Moreno,**
12 **JM., Soria, B. 2009.** Developing biological resource banks as a supporting tool for wildlife
13 reproduction and conservation: the Iberian lynx bank as a model for other endangered
14 species. *Animal Reproduction Science* **112**:347–361. doi:10.1016/j.anireprosci.2008.05.070
- 15 **Li LF, Guan WJ, Li H, Zhou XZ, Bai XJ, Ma YH. 2009.** Establishment and characterization
16 of a fibroblast cell line derived from Texel sheep. *Biochemistry and Cell Biology* **87**:485–
17 492. DOI: 10.1139/O09-005.
- 18 **Liu C., Guo Y., Guan W., Ma Y., Zhang H., Tang X. 2008.** Establishment and biological
19 characteristics of Luxi cattle fibroblast bank. *Tissue and Cell* **40**:417–424. DOI:
20 10.1016/j.tice.2008.04.005.
- 21 **Luo J., Liang M., Yang X., Xu H., Shi D., Lu S. 2014.** Establishment and biological
22 characteristics comparison of Chinese swamp buffalo (*Bubalus bubalis*) fibroblast cell lines.
23 *In vitro Cellular & Developmental Biology-Animal* **50**:7–15. DOI: 10.1007/s11626-013-
24 9677-2.
- 25 **Magalhães LC., Bhat MH., Freitas JL., Melo LM., Teixeira DI., Pinto LC., Câmara LM.,**
26 **Duarte JM., Freitas VJ. 2017.** The effects of cryopreservation on different passages of
27 fibroblast cell culture in brown brocket deer (*Mazama gouazoubira*). *Biopreservation and*
28 *Biobanking* **15**:463–468. DOI: 10.1089/bio.2017.0060.
- 29 **Magalhães R., Nugraha B., Pervaiz S., Yu H., Kuleshova LL. 2012.** Influence of cell culture
30 configuration on the post-cryopreservation viability of primary rat hepatocytes. *Biomaterials*
31 **33**:829–836. DOI: 10.1016/j.biomaterials.2011.10.015.
- 32 **Mata M., Garrido-Maraver J., Cotán D., Cordero MD., Oropesa-Ávila M., Izquierdo LG.,**
33 **De Miguel M., Lorite JB., Infante ER., Ybot P. 2012.** Recovery of MERRF fibroblasts
34 and cybrids pathophysiology by Coenzyme Q 10. *Neurotherapeutics* **9**:446–463. DOI:

- 1 10.1007/s13311-012-0103-3.
- 2 **Mehrabani D., Mahboobi R., Dianatpour M., Zare S., Tamadon A., Hosseini SE.**
- 3 **2014.** Establishment, culture, and characterization of Guinea pig fetal fibroblast cell.
- 4 *Veterinary Medicine International* **2014**:1–5. DOI: 10.1155/2014/510328.
- 5 **Queiroz Neta LB., Oliveira Lira GPO., Borges AA., Santos MVO., Silva MB., Oliveira**
- 6 **LRM., Silva AR., Oliveira MF., Pereira AF. 2018.** Influence of storage time and nutrient
- 7 medium on recovery of fibroblast-like cells from refrigerated collared peccary (*Pecari tajacu*
- 8 Linnaeus, 1758) skin. *In vitro Cellular & Developmental Biology-Animal* **54**:486–495. DOI:
- 9 10.1007/s11626-018-0270-6.
- 10 **Redford KH. 1992.** The empty forest. *BioScience* **42**:412–422.
- 11 **Rodriguez-Osorio N., Urrego R., Cibelli J., Eilertsen K., Memili E. 2012.** Reprogramming
- 12 mammalian somatic cells. *Theriogenology* **78**:1869–1886. DOI:
- 13 10.1016/j.theriogenology.2012.05.030.
- 14 **Roth V. 2006.** Available at <http://www.doubling-time.com/compute.php> (accessed 01 September
- 15 2019).
- 16 **Saadeldin IM., Swelum AAA., Noreldin AE., Tukur HA., Abdelazim AM., Abomughaid**
- 17 **MM., Alowaimer AN. 2019.** Isolation and culture of skin-derived differentiated and stem-
- 18 like cells obtained from the Arabian Camel (*Camelus dromedarius*). *Animals* **9**:378–395.
- 19 DOI:10.3390/ani9060378
- 20 **Santos DO., Mendes A., Nogueira SS da C., Nogueira Filho SLG. 2009.** Criação comercial
- 21 de caititus (*Pecari tajacu*): uma alternativa para o agronegócio. *Revista Brasileira de Saúde*
- 22 *e Produção Animal* **10**:1–10.
- 23 **Santos MLT., Borges AA., Queiroz Neta LB., Santos MVO., Oliveira MF., Silva AR.,**
- 24 **Pereira AF. 2016.** *In vitro* culture of somatic cells derived from ear tissue of collared
- 25 peccary (*Pecari tajacu* Linnaeus, 1758) in medium with different requirements. *Pesquisa*
- 26 *Veterinária Brasileira* **36**:1194–1202.
- 27 **Santos, MVO., Nascimento, LE., Praxedes, EA., Borges, AA., Silva, AR., Bertini, LM.,**
- 28 **Pereira, AF. 2019.** *Syzygium aromaticum* essential oil supplementation during *in vitro*
- 29 bovine oocyte maturation improves parthenogenetic embryonic development.
- 30 *Theriogenology* **128**: 74–80. DOI:10.1016/j.theriogenology.2019.01.031
- 31 **Sharma R., Sharma H., Ahlawat S., Aggarwal R., Vij P., Tania M. 2018.** First attempt on
- 32 somatic cell cryopreservation of critically endangered *Camelus bactrianus* of India. *Gene*
- 33 *Reports* **10**:109–115. DOI: 10.1016/j.genrep.2017.11.007.
- 34 **Schaeffer WI. 1990.** Terminology associated with cell, tissue and organ culture, molecular

- 1 biology and molecular genetics. *In vitro Cellular & Developmental Biology* **26**:97–101.
- 2 **Shiga, K., Fujita, T., Hirose, K., Sasae, Y., Nagai, T. 1999.** Production of calves by transfer
3 of nuclei from cultured somatic cells obtained from Japanese black bulls. *Theriogenology* **52**:
4 527–535. DOI: 10.1016/S0093-691X(99)00149-1
- 5 **Siengdee P., Klinhom S., Thitaram C., Nganvongpanit K. 2018.** Isolation and culture of
6 primary adult skin fibroblasts from the Asian elephant (*Elephas maximus*). *PeerJ* **6**:e4302.
7 DOI: 10.7717/peerj.4302.
- 8 **Silvestre MA., Sánchez JP., Gómez EA. 2004.**Vitrification of goat, sheep, and cattle skin
9 samples from whole ear extirpated after death and maintained at different storage times and
10 temperatures. *Cryobiology* **49**:221–229. DOI: 10.1016/j.cryobiol.2004.08.001.
- 11 **Song J., Hua S., Song K., Zhang Y. 2007.** Culture, characteristics and chromosome
12 complement of Siberian tiger fibroblasts for nuclear transfer. *In vitro Cellular &*
13 *Developmental Biology-Animal* **43**:203–209. DOI: 10.1007/s11626-007-9043-3.
- 14 **Trokovic R., Weltner J., Noisa P., Raivio T., Otonkoski T. 2015.** Combined negative effect
15 of donor age and time in culture on the reprogramming efficiency into induced pluripotent
16 stem cells. *Stem Cell Research* **15**:254–262. DOI: 10.1016/j.scr.2015.06.001.
- 17 **Yajing S., Rajput IR., Ying H., Fei Y., Sanganyado E., Ping L., Jingzhen W., Wenhua L.**
18 **2018.** Establishment and characterization of pygmy killer whale (*Feresa attenuata*) dermal
19 fibroblast cell line. *PloS one* **13**: e0195128. DOI: 10.1371/journal. pone.0195128.

20

21

**CAPÍTULO 4 – CRYOPRESERVATION OF COLLARED PECCARY (*Pecari tajacu*
LINNAEUS, 1758) SOMATIC CELLS IS IMPROVED BY SUCROSE AND HIGH
CONCENTRATIONS OF FETAL BOVINE SERUM**

(Artigo experimental)

PUBLICADO NA CRYO-LETTERS

QUALIS: Quadriênio 2013-2016: B1/ Novo Qualis: A2

FATOR DE IMPACTO: 0,694

CRYOPRESERVATION OF COLLARED PECCARY (*Pecari tajacu* LINNAEUS, 1758) SOMATIC CELLS IS IMPROVED BY SUCROSE AND HIGH CONCENTRATIONS OF FETAL BOVINE SERUM

Gabriela Pereira de Oliveira Lira¹, Alana Azevedo Borges¹, Matheus Barbosa do Nascimento¹, Leonardo Vitorino Costa de Aquino¹, Moacir Franco de Oliveira², Alexandre Rodrigues Silva³ and Alexsandra Fernandes Pereira^{1*}

¹Laboratory of Animal Biotechnology, Federal Rural University of Semi-Arid (UFERSA), Mossoro, RN, Brazil.

²Laboratory of Applied Animal Morphophysiology, UFERSA, Mossoro, RN, Brazil.

³Laboratory of Animal Germplasm Conservation, UFERSA, Mossoro, RN, Brazil

*Corresponding author e-mail: alexsandra.pereira@ufersa.edu.br

Abstract

BACKGROUND: The formation of somatic cell banks is affected by, amongst other factors, the cryoprotectant solution used. The selection of an effective solution, therefore, is a primary parameter. **OBJECTIVE:** We optimized the cryoprotectant used for collared peccary somatic cell cryopreservation. **MATERIALS AND METHODS:** We categorized cells into different groups based on their cryopreservation and evaluated the morphology, viability, proliferative activity, metabolism, and oxidative stress. One group was cryopreserved in 10% DMSO with 10% fetal bovine serum (DMSO-10FBS), and another with 50% FBS (DMSO-50FBS). The cryopreservation of both groups included the presence of 0.2 M sucrose (DMSO-SUC-10FBS and DMSO-SUC-50FBS). Non- cryopreserved cells and cells cryopreserved with 10% DMSO (DMSO) supplemented with 0.2 M sucrose (DMSO-SUC) were used as controls. **RESULTS:** There was no difference observed in morphology or viability among the groups. Proliferative activity was reduced in DMSO-10FBS when compared to controls. Although cryopreservation reduced metabolism, no difference was observed among solutions. A lower level of reactive oxygen species was observed in cells of DMSO-SUC- 50FBS when compared to other cryoprotectants. Only cells of DMSO-SUC-50FBS had mitochondrial potential similar to non-cryopreserved cells. **CONCLUSION:** 10% DMSO supplemented with 50% FBS and 0.2 M SUC was observed to be the most efficient cryoprotectant for preserving collared peccary somatic cells.

Keywords: Peccaries, extracellular cryoprotectants, slow freezing, cryobanking.

Introduction

Collared peccaries (*Pecari tajacu* Linneaus, 1758), also known as “wild pigs”, are wild mammals belonging to the Tayassuidae family of the order Artiodactyla (21). These animals play an important role in maintaining the global ecosystem where they act as seed dispersers, promote flora renewal, and contribute to the food chain (34). Although globally classified as “Least Concern”, this species is already extinct in eastern and southern Argentina and is declining in some biomes (9), thus necessitating conservation strategies for the maintenance of its population.

One of the conservation tools applied to wild mammals has been the formation of somatic resource banks (6). These banks allow long-term storage of tissues and somatic cells to be employed in different proposals, such as multiplying individuals by somatic cell

nuclear transfer (SCNT) (43) and generating induced pluripotent cells (41). The efficiency of these procedures depends on the quality of cells after slow freezing, which is dependent on the choice of the cryoprotectant, a crucial step for the success of these cryobanks (28). In general, an effective cryoprotectant is the appropriate combination of intracellular cryoprotectants such as dimethyl sulfoxide (DMSO) (25), ethylene glycol (EG) (16), and extracellular cryoprotectants such as sucrose (SUC) (23) and/or fetal bovine serum (FBS) (15).

Specifically, our group has successfully established somatic tissue banks of collared peccaries based on our knowledge of the tissues (2), techniques (3), and solutions required (4, 5). Subsequently, we initiated somatic cell banks of this species, demonstrating that 10% DMSO in the presence of 0.2 M SUC was more efficient in somatic cell slow freezing when compared to EG with 0.2 M SUC, showing a viability of 69.8% and 58.5%, respectively (17). Thus, one way to improve the rates of cell recovery would be to optimize the extracellular agents employed. Studies have shown that FBS, when associated with 10% DMSO and 0.2 M SUC, promoted a beneficial effect in some species (*Lynx pardinus* and *Elephas maximus*) (15, 36). This agent can reduce oxidative stress, caused by excessive production of reactive oxygen species (ROS) during slow freezing (11).

Nevertheless, variations in FBS concentration can be observed in some species with its use at 10% (*Felis silvestris libica*) (8) and 50% (*Bubalus bubalis*) (18), necessitating the evaluation of the most appropriate concentration, as well as its relationship with SUC. Therefore, we aimed to optimize the cryoprotectant by varying the different extracellular combinations of SUC and FBS. We assessed the impact of the cryoprotectants on the morphology, viability, proliferative activity, metabolism, and oxidative stress of cryopreserved collared peccary cells.

Materials and Methods

Chemicals and media

All chemicals used were purchased from Sigma-Aldrich (St. Louis, MO, USA). Dulbecco's modified Eagle's medium (DMEM), FBS, trypsin-EDTA, Antibiotic-Antimycotic (ATB-ATM) solution and fluorescent probes were obtained from Gibco-BRL (Carlsbad, CA, USA). The 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) assay was purchased from Grainer (Miami, FL, USA).

Compliance with ethical standards and animals

The experimental protocols and animal handling procedures were performed with the approval of the Animal Ethics Committee of the Federal Rural University of Semi-Arid (CEUA/UFERSA, no. 23091.001072/2015-92), in compliance with the Chico Mendes Institute for Biodiversity Conservation (ICMBio, no. 48633-2). A total of four adult collared peccaries, 41.5 ± 1.8 months old, provided by the Centre of Multiplication of Wild Animals (CEMAS/UFERSA, Mossoró, RN, Brazil; 5°10'S, 37°10'W) were used.

Skin biopsy, primary culture, and subcultures

Sections (1-2 cm²) used for collared peccary identification were taken from peripheral ear tissue using pliers. After collection, tissue samples were washed in 70% ethanol and incubated in DMEM supplemented with 10% FBS and 2% ATB-ATM solution at 37 °C for 30 min.

In the laboratory, tissues were trichotomized, washed in 70% ethanol followed by DMEM supplemented with 10% FBS and 2% ATB-ATM solution. Subsequently, tissue samples were fragmented in 9.0 mm³ sections and cultured in DMEM supplemented with 10% FBS and 2% ATB-ATM solution at 38.5 °C in an incubator with 5% CO₂ and 95% air (31). During the primary culture, the medium was changed every 24 h. Cells were subcultured after reaching 70-80% confluence until the third passage, followed by cryopreservation. The subconfluence of 70-80% was defined when 70-80% of the petri dishes presented somatic cells (31).

Study design and cell cryopreservation

The morphology, viability, proliferative activity, metabolism, and oxidative stress were evaluated in non-cryopreserved and cryopreserved cells. Cells were cryopreserved in either 10% DMSO with 10% FBS (DMSO-10FBS group), or 50% FBS (DMSO-50FBS group). In both the groups, 0.2 M SUC was added and these were further designated as the DMSO-SUC-10FBS and DMSO-SUC-50FBS groups, respectively. Moreover, non-cryopreserved cells (non-cryopreserved group), those cryopreserved in 10% DMSO (DMSO group) and those cryopreserved with 10% DMSO supplemented with 0.2 M SUC (DMSO-SUC group), were used as controls.

For cryopreservation, cells were subjected to slow cooling in freezing medium (DMEM supplemented with 10% DMSO and extracellular cryoprotectants according to experimental groups). Briefly, cells at a concentration of 1.0×10^4 cells/mL were first exposed to DMSO solution for 15 min at 4 °C in DMEM, followed by the addition of SUC and FBS solutions according to experimental groups and incubated for an additional 15 min at 4 °C. The cryovials containing 1.0 mL of cells in freezing solution were cooled in a Mr. Frosty freezing container (Thermo Scientific Nalgene, Rochester, NY, USA) at a cooling rate of 1 °C/min, and kept in a freezer -80 °C until reaching -70 °C, before being transferred to liquid nitrogen (15).

After 2 weeks, cryovials were removed from liquid nitrogen and kept at room temperature of 25 °C for up to 1 min and then warmed in a water bath at 37 °C for 3-4 min. For the removal of cryoprotectants, the cells were washed twice with DMEM. Cell suspension was centrifuged for 10 min at 400×g and washed again with DMEM and centrifuged according to León-Quinto et al. (15). In the groups containing SUC, the cells were washed with DMEM with 0.2 M SUC, prior to centrifugation as previously described.

Evaluation of morphological characteristics and cell viability

The cells' morphology was evaluated daily using an inverted microscope (Nikon TS100, Tokyo, Japan). The following morphological characteristics were assessed: size, aspect, shape and adhesion (29). The viability analysis was performed by cells stained with trypan blue. For each animal and each group, the viability test was performed in duplicate. Briefly, the cells were centrifuged and suspended in 1.0 mL of the cell culture medium (DMEM supplemented with 10% FBS and 2% ATB-ATM solution); an aliquot of cells was stained with 0.4% trypan blue (in phosphate buffered saline, PBS) in the ratio 1:1 and counted in a Neubauer chamber (39).

Analysis of proliferative activity and metabolism

Proliferative activity was quantified according to the population doubling time (PDT), which was calculated by seeding cells at a density of 1.0×10^4 cells/well in a twenty-four well plate. Cells were trypsinized in duplicate and counted at 24 h intervals for up to 216 h of culture. The mean cell counts were recorded every time and the cell growth curve was delineated. Finally, the PDT was calculated using the following formula (30):

PDT = $T \ln 2 / \ln (X_e/X_b)$, where PDT is the time of the culture (in hours), T is the incubation time, X_b is the number of cells at the beginning of the time incubation, X_e is the number of cells at the end of the incubation time, and $\ln$ is Napierian logarithm.

For evaluation of metabolism, cells were seeded into 12-well dishes at a density of 5.0×10^4 cells/mL and cultured at 38.5 °C in a humid atmosphere with 5% CO₂ (4). After five days, MTT solution (5 mg/mL in DMEM) was added to each well for 3 h under the same conditions. After incubation, the MTT solution was removed, and DMSO was added to each dish to solubilize the formazan crystals. Finally, the absorbance (595 nm) of the samples was read using a spectrophotometer (Shimadzu® UV-mini-1240, Kyoto, Japan). The mean values of non-cryopreserved cells were considered 100% as compared to the other groups.

Assessment of oxidative stress

Cellular oxidative stress was evaluated by quantification of intracellular reactive oxygen species (ROS) levels and measurement of mitochondrial membrane potential ($\Delta\Psi_m$). Briefly, intracellular levels of ROS were quantified using 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA) as a fluorescent probe, according to Santos et al. (32). For this, the cells, independent of the experimental group, were seeded in glass slides and grown *in vitro* until reaching 70% confluence. Subsequently, cells were washed in PBS and incubated with 1.0 mL 5 μ M H₂DCFDA at 38.5 °C with 5% CO₂ for 30 min protected from light. After this period, the stained cells were washed twice in PBS and imaged using a fluorescence microscope (Olympus BX51TF, Tokyo, Japan). The intensity of the signal fluorescence (pixels) was measured using ImageJ software (version 1.49v, Java 1.8.0_201, Wayne Rasband, U.S. National Institutes of Health, Bethesda, MD, USA; website: <http://rsb.info.nih.gov/ij/download.html>). The background signal intensity was subtracted from the values obtained for the treated samples. Measured mean value of the micrograph for non-cryopreserved cells was used as a calibrator. Relative expression levels (arbitrary fluorescence units) were generated by dividing the measured value of each micrograph for the cryopreserved cells by the mean of the calibrator.

The $\Delta\Psi_m$ was assessed using the fluorescent probe MitoTracker Red® (CMXRos) at 500 nm according to Santos et al. (32). The procedure, incubation and evaluation of the images, were performed as described for the quantification of ROS.

1 *Statistical analysis*

2 Data were expressed as mean $\pm$ standard error (one animal/one repetition) and analyzed
3 using the GraphPad software (Graph-Pad Software Incorporated, La Jolla, CA, USA). All
4 results were verified for normality by the Shapiro-Wilk test and homoscedasticity by
5 Levene's test. Since data did not show a normal distribution, they were arcsine transformed
6 and analyzed by ANOVA followed by Tukey test. Significance was set at $P < 0.05$.


8 **Results**

9 *Evaluation of morphological characteristics and cell viability*

10 All cells derived from collared peccary tissue had consistent morphological
11 characteristics in all experimental groups (Fig. 1a-g). Slow cooling did not alter the capacity
12 of the cells to grow and reach 70-80% confluence, presenting cells with similar morphology
13 to fibroblasts, with fusiform aspect, cytoplasmic prolongations, abundant cytoplasm and
14 central nucleus, with an overall large and oval morphology.

15
16 Moreover, no difference was observed among groups for cell viability evaluated by
17 trypan blue (Fig. 2, $P > 0.05$). Thus, while non-cryopreserved cells had a viability of $87.0\% \pm$
18 5.2 , cryopreserved cells of different groups showed viability in the range of $62.5\% \pm 5.2$ to
19 $91.7\% \pm 0.0$.

1



2 **Figure 1.** Subcultures of fibroblast-like cells from ear skin samples of collared peccaries. **a)** Non-cryopreserved cells and cell cryopreserved in **b)**
3 DMSO, **c)** DMSO-SUC, **d)** DMSO-10FBS, **e)** DMSO-50FBS, **f)** DMSO-SUC-10FBS, **g)** DMSO-SUC-50FBS. Scale bar = 100 μ m.




Figure 2. Viability of collared peccary non-cryopreserved and cryopreserved cells using different extracellular cryoprotectants. Non-cryopreserved group, 10% DMSO (DMSO group), 10% DMSO and 0.2 M SUC (DMSO-SUC group), 10% DMSO and 10% FBS (DMSO-10FBS group), 10% DMSO and 50% FBS (DMSO-50FBS group), 10% DMSO, 0.2 M SUC and 10% FBS (DMSO-SUC-10FBS group), 10% DMSO, 0.2 M SUC and 50% FBS (DMSO-SUC-10FBS group). $P > 0.05$.

Analysis of proliferative activity and metabolism

Although the growth curve did not show a typical “S” shape in all cryopreserved groups, it was possible to observe from the shape of the curve the occurrence of the initial adaptation phase of the cells, followed by exponential growth until reaching the stable phase of decline (Fig. 3a). In addition, non-cryopreserved cells showed a reduction in their growth, however, this reduction stabilized from day 8. No longer observed the decline in the curve, which showed that there was no cell death only the decrease of proliferative activity of cells from this group. Additionally, the proliferative activity evaluated by population doubling time was reduced in cells derived from DMSO-10FBS group ($10.5 \text{ h} \pm 0.3$) when compared to non-cryopreserved cells (19.4 ± 2.6 , Fig. 3b).

Although cryopreservation reduced cell metabolism, no difference was observed among cryoprotectant solutions ($P > 0.05$). Thus, while cryopreserved cells had a metabolic activity of $100.0\% \pm 0.0$, cryopreserved cells among the different groups varied among $63.6\% \pm 22.6$ and $97.7\% \pm 25.1$.




Figure 3. The growth dynamics and metabolism of collared peccary non-cryopreserved and cryopreserved cells using different extracellular cryoprotectants. **a)** Growth curves of cryopreserved and non-cryopreserved cells. **b)** Values of population doubling time (PDT) after culture for nine days. Non-cryopreserved group, 10% DMSO (DMSO group), 10% DMSO and 0.2 M SUC (DMSO-SUC group), 10% DMSO and 10% FBS (DMSO-10FBS group), 10% DMSO and 50% FBS (DMSO-50FBS group), 10% DMSO, 0.2 M SUC and 10% FBS (DMSO-SUC-10FBS group), 10% DMSO, 0.2 M SUC and 50% FBS (DMSO-SUC-10FBS group). Different letters show significant differences among the groups ($P < 0.05$). Bars indicate standard error.

Assessment of oxidative stress

A lower level of ROS was observed in cells derived from DMSO-SUC-50FBS when compared to other cryopreserved groups (Fig 4a and 4c).

Moreover, only cells from the DMSO-SUC-50FBS group had mitochondrial membrane potential similar to the non-cryopreserved cells (Fig 4b and 4d).




Figure 4. Assessment of oxidative stress in collared peccary non-cryopreserved and cryopreserved cells using different extracellular cryoprotectants. Non-cryopreserved group, 10% DMSO (DMSO group), 10% DMSO and 0.2 M SUC (DMSO-SUC group), 10% DMSO and 10% FBS (DMSO-10FBS group), 10% DMSO and 50% FBS (DMSO-50FBS group), 10% DMSO, 0.2 M SUC and 10% FBS (DMSO-SUC-10FBS group), 10% DMSO, 0.2 M SUC and 50% FBS (DMSO-SUC-10FBS group). **a)** Quantification of intracellular reactive oxygen species (ROS) levels **b)** Measurement of mitochondrial membrane potential ($\Delta\Psi_m$). **c)** Cells labeled with H₂DCFDA for quantification of ROS levels. **d)** Cells labeled with MitoTracker Red[®] for measurement of $\Delta\Psi_m$. Different letters within each analysis show significant differences among the groups ($P < 0.05$). Scale bar = 200 μ m.

Discussion

The successful formation of a bank of biological resources is correlated to the efficiency of the cryopreservation protocol used. As such, the initial combination of intracellular and extracellular cryoprotectants is key in ensuring cell quality after thawing. In this study, we showed that the combination of 0.2 M SUC with 50% FBS added to the cryoprotectant solution containing 10% DMSO, ensured a better quality of the recovered collared peccary somatic cells, particularly by reducing the oxidative stress caused by slow freezing and by maintaining normal cellular characteristics. Therefore, our results demonstrated three relevant points of cryopreservation of somatic cells: (a) the cells obtained from collared peccary tissue showed characteristics similar to fibroblast cells used as the main donor nucleus in SCNT, (b) some alterations resulting from the cryopreservation may not be observed from the analysis of the viability and metabolic activity of the thawed cells; therefore, more specific assessments such as the analysis of the oxidative stress may be necessary, and (c) the favorable effect of the addition of 50% FBS on the cryoprotectant solution containing 10% DMSO and 0.2 M SUC. Thus, these findings ensure that higher quality cells are employed in future conservation strategies.

In general, cryopreservation protocols can induce lesions such as lipid peroxidation of membranes and damage to DNA in fibroblasts, resulting from the formation of ROS (33). Moreover, the natural antioxidant defense mechanism of the cells is not enough to eliminate excess ROS (11). Therefore, studies have shown that supplementation with antioxidants and/or factors that modulate the cell death process caused by excess ROS may be a potential solution to reduce oxidative stress induced by cryopreservation (14). Additionally, a decrease in $\Delta\Psi_m$ has been observed in cells with excess ROS formation (13). Thus, as suggested by Poot et al. (27), we used a fluorescent probe that has an affinity for molecules present in cells. Such probes are responsible for evaluating the relationship among mitochondria and the triggering of apoptotic mechanisms. With this, it was possible to demonstrate that among the cryoprotectant solutions tested, the cells obtained from the DMSO-SUC-50FBS treatment showed high $\Delta\Psi_m$ after thawing.

Thus, one compound that has such antioxidant potential is FBS, which has been frequently used in cryoprotectant solutions (12). In general, FBS is routinely added to stabilize cell membranes and adjust intra- and extracellular osmotic pressure; however, the

1 role of this cryoprotectant as an antioxidant has become increasingly evident, which may be
2 related to its constituents, mostly amino acids, being formed (20). This could be due to the
3 synergistic effect of their constituents, which may have attached to cell membranes,
4 promoting stabilization and avoiding excessive concentration of solutes that may be related to
5 ROS formation (12). In addition, their constituents may be responsible for the increased
6 activity of antioxidant intracellular enzymes (26). Thus, it is likely that the FBS in the
7 DMSO-SUC-FBS solution acted as a source of antioxidants. However, the mechanisms
8 underlying the antioxidant role of FBS under the formation of ROS suffered by cells during
9 cryopreservation are still unclear. Moreover, when the cells were cryopreserved with 10%
10 FBS (DMSO-10FBS and DMSO-SUC-10FBS groups), the fluorescence intensity increased,
11 indicating an increase in ROS production. This suggests that the reduction and/or absence of
12 antioxidative agents in the DMSO, DMSO-10FBS and DMSO-SUC-10FBS groups may be
13 associated with the formation of ROS (24, 42, 44).

14
15 This result may be due to FBS counteracting the cytotoxicity induced by DMSO, as
16 demonstrated by Miki et al. (22), when analyzing the $\Delta\Psi_m$ of stem cells subjected to a
17 cryoprotectant solution with 10% DMSO in association with 90% FBS, which resulted in a
18 $\Delta\Psi_m$ of 1.00. Therefore, treatments with 10% FBS caused the cells to be more exposed to
19 cytotoxicity caused by DMSO during the freeze-thaw cycle. The DMSO penetrates the cell
20 membrane and forms hydrogen bonds with the water molecules present in the cytosol. Thus,
21 the freezing point of the water decreases, and there is a lower probability of the formation of
22 ice crystals inside the cell. Furthermore, the integrity of the membrane is conserved and for
23 this reason, DMSO is an indispensable intracellular cryoprotectant. However, even though its
24 positive effect is observed, its toxicity remains high, suggesting its association with
25 extracellular cryoprotectants such as SUC and FBS. This association can accelerate the
26 cellular dehydration process and balance cytotoxicity. Therefore, the ideal concentrations for
27 achieving this cytotoxic balance must be established so that the concentrations of extracellular
28 cryoprotectants are sufficient to minimize the negative effect of the high toxicity of DMSO.
29 Additionally, the presence of SUC can promote osmotic maintenance of the medium and
30 perform synergistically with the intracellular cryoprotectant, resulting in improved effects
31 when used in combination (37).

1 The ROS levels of the DMSO-SUC-10FBS group were higher in comparison to the
2 DMSO group, probably due to the use of more cryoprotective agents in their composition. It
3 should be borne in mind that such cryoprotective agents (DMSO-SUC-10FBS), despite
4 promoting cell protection, present a certain toxicity that may be responsible for greater
5 oxidative stress and subsequent increased production of ROS. However, this does not mean
6 that the use of fewer cryoprotective agents is recommended, but rather that a combination
7 promoting balance between the concentration used, toxicity, protection of cells at cryogenic
8 temperatures and lower production of ROS, should be utilized.

9
10 In addition, the morphological and confluence aspects observed in the collared cells were
11 similar to those observed by Queiroz Neta et al. (29) and Borges et al. (4) describing collared
12 peccary cells recovered from refrigerated and vitrified tissues, respectively. Thus, proper
13 maintenance of the morphological characteristics of cultured cells may be associated with a
14 decrease in ice crystal formation due to the cryopreservation process, which when exposed to
15 cells, may have deleterious effects on cell characteristics (15).

16
17 The viability data showed that cryopreserved cells and non-cryopreserved cells were
18 stable regardless of treatments. This can be attributed to the quality of the 10% DMSO slow
19 freezing technique that is commonly associated with better results for somatic cells derived
20 from wild mammals (23, 35).

21
22 Further, the data found on higher viability rates and the performance of cells in vitro
23 culture shown in this study, corroborate the findings of Subramanian et al. (40) and Barnes et
24 al. (1). These authors affirmed that cultures derived from cryopreserved and non-
25 cryopreserved cells have a similar growth performance, since the cell population after culture
26 can recover with rates similar to non-cryopreserved cells.

27
28 Finally, all groups showed no change in PDT values except DMSO-10FBS. In general,
29 cells cultured in specific treatments including FBS in their composition required less time to
30 replicate (19). Our PDT rates and metabolism activity evaluated by MTT may be associated
31 with the use of an adequate system of cryopreservation that promoted an optimum cooling
32 rate (10) associated with compounds present in FBS such as transport proteins, adhesion

proteins, enzymes, hormones, growth factors, cytokines, fatty acids, lipids, vitamins, carbohydrates, and nitrogen of non-protein origin (7).

In conclusion, 10% DMSO supplemented with 0.2 M SUC and 50% FBS was the most efficient cryoprotectant solution for conserving collared peccary somatic cells. This work presents a breakthrough for the improvement and establishment of an ideal freezing protocol for collared peccary somatic cells that can be used in future biotechnical techniques such as cloning, since each stage involved should be optimized for the species studied.

Acknowledgements: The authors thank the Centre for Wild Animals Multiplication (CEMAS/UFERSA) for providing the animals. This study was financed in part by the Higher Education Personnel Improvement Coordination - Brazil (CAPES, Financial Code 001) and National Counsel of Technological and Scientific Development (CNPq). MF Oliveira, AR Silva and AF Pereira were CNPq investigators.

References

1. Barnes LM, Bentley CM, Dickson AJ (2003) *Biotechnol Prog* **19**, 233–237.
2. Borges AA, Bezerra FVF, Costa FN, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, Pereira AF (2017a) *Arq Bras Med Vet Zootec* **69**, 948–954.
3. Borges AA, Lima GL, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, Pereira AF (2017b) *Cytotechnology* **69**, 643–654.
4. Borges AA, Lira GPO, Nascimento LE, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, Pereira AF (2018) *Biopreserv Biobank* **16**, 77–81.
5. Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, Pereira AF (2018) *Pesq Vet Bras* **38**, 350–356.
6. Borges AA, Pereira AF (2019) *Zygote* **27**, 111–117.
7. Brunner D, Frank J, Appl H, Schöffl H, Pfaller W, Gstraunthaler G (2010) *Altex* **27**, 53–62.
8. Gómez MC, Jenkins JA, Giraldo A, Harris RF, King A, Dresser AL, Pope CE (2003) *Biol Reprod* **69**, 1032–1041.
9. Gongora J, Biondo C, Cooper JD, Taber A, Keuroghlian A, Altrichter M, Nascimento FF, Chong AY, Miyaki CY, Bodmer R, Mayor P, González S (2011) *Bonn Zoolog Bull* **60**, 95–101.

- 1 10. Gurruchaga H, Ciriza J, del Burgo LS, Rodriguez-Madoz JR, Santos E, Prosper F,
2 Hernández RM, Orive G, Pedraz JL (2015) *Int J Pharm* **485**, 15–24.
- 3 11. Gurruchaga H, del Burgo LS, Hernández RM, Orive G, Selden C, Fuller B, Ciriza J, Pedraz
4 JL (2018) *J Control Release* **281**, 119–138.
- 5 12. Kimura Y, Okitsu T, Xibao L, Teramae H, Okonogi A, Toyoda K, Uemoto S, Fukushima
6 M (2013) *Islets* **5**, 45–52.
- 7 13. Lançon R, Celeghini ECC, Alves MBR, Lemes KM, Gonella-Diaza AM, Oliveira LZ,
8 Arruda AR (2018) *J Equine Vet Sci* **69**, 78–83.
- 9 14. Len JS, Koh WSD, Tan WST (2019) *Biosci Rep* **39**, BSR20191601.
- 10 15. León-Quinto T, Simón MA, Cadenas R, Martínez A, Serna A (2014) *Cryobiology* **68**, 227–
11 233.
- 12 16. Li Y, Lu RH, Luo GF, Pang WJ, Yang GS (2006) *Cryobiology* **53**, 240–247.
- 13 17. Lira GPO, Borges AA, Nascimento LE, Santos MVO, Queiroz Neta LB, Oliveira MF,
14 Pereira AF (2017) in: *Anais do Encontro de Biotecnologia do Nordeste (RENORBIO)*.
- 15 18. Luo C, Lu C, Wang X, Wang Z, Li X, Gong F, Jiang J, Liu Q, Shi D (2013)
16 *Theriogenology* **80**, 878–886.
- 17 19. Mahesh YU, Rao BS, Katari VC, Komjeti S, Christo D, Lakshmikantan U, Pawar RM,
18 Shivaji S (2012) *Reprod Domest Anim* **47**, 799–805.
- 19 20. Marco-Jiménez F, Garzón DL, Peñaranda DS, Pérez L, Viudes-de-Castro MP, Vicente JS,
20 Jover M, Asturiano JF (2006) *Cryobiology* **53**, 51–57.
- 21 21. Mayer JJ, Wetzel RM (1987) *Mamm. Species* **293**, 1–7.
- 22 22. Miki T, Wong W, Zhou E, Gonzalez A, Garcia I, Grubbs BH (2016) *Stem Cell Res Ther* **7**,
23 8.
- 24 23. Moulavi F, Hosseini SM, Tanhaie-Vash N, Ostadhosseini S, Hosseini SH, Hajinasrollah M,
25 Asghari MH, Gourabi H, Shahverdi A, Vosough AH, Nasr-Esfahani MH (2017)
26 *Theriogenology* **90**, 197–203.
- 27 24. Nirmala JG, Akila S, RT Narendhirakannan, Chatterjee S (2017) *Adv Powder Technol* **28**,
28 1170–1184.
- 29 25. Oliveira LRM, Silva MB, Praxedes EA, Ribeiro LR, Silva HVR, Pereira AF (2020) *An*
30 *Acad Bras Ciênc* In press.
- 31 26. Piao MJ, Hyun YJ, Cho SK, Kang HK, Yoo ES, Koh YS, Lee NH, Ko MH, Hyun JW
32 (2012) *Mar Drugs* **10**, 2826–2845.
- 33 27. Poot M, Gibson LL, Singer VL (1997) *Cytometry A* **27**, 358–364.

- 1 28. Praxedes EA, Borges AA, Santos MVO, Pereira AF (2018) *Zoo Biol* **37**, 258–263.
- 2 29. Queiroz Neta LB, Lira GPO, Borges AA, Santos MVO, Silva MB, Oliveira LRM, Silva
- 3 AR, Oliveira MF, Pereira AF (2018) *In Vitro Cell Dev Biol Anim* **54**, 486–495.
- 4 30. Roth V, Available at: <http://www.doubling-time.com/compute.php>, (2006), Accessed date:
- 5 12 February 2019.
- 6 31. Santos MLT, Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, Pereira
- 7 AF (2016) *Pesq Vet Bras* **36**, 194–1202.
- 8 32. Santos MVO, Nascimento LE, Praxedes EA, Borges AA, Silva AR, Bertini LM, Pereira AF
- 9 (2019) *Theriogenology* **128**, 74–80.
- 10 33. Santos TRM, Silva ACG, Carvalho FS, Sanz G, Vaz BG, Lião LM, Menegatti R, Valadares
- 11 MC (2018) *Chem-Biol Interact* **293**, 20–27.
- 12 34. Schlichting PE (2018) *J. Wildl. Manage* **82**, 1556–1557.
- 13 35. Sharma R, Sharma H, Ahlawat S, Aggarwal RAK, Vij PK, Tania MS (2018) *Gene Rep*
- 14 **10**, 109–115.
- 15 36. Siengdee P, Klinhom S, Thitaram C, Nganvongpanit C (2018) *PeerJ* **6**, e4302.
- 16 37. Silva AR, Campos LB, Maia KM, Borges AA (2017) *Rev Bras Reprod Anim* **41**, 110–115.
- 17 38. Singh M, Degala V (2017) *Int J Biol* **9**, 9–13.
- 18 39. Strober W (2015) *Curr Protoc Immunol* **111**, A3-B.
- 19 40. Subramanian J, Aulakh RPS, Grewal PS, Sanford M, Pynn AFJ, Yuk IH (2018) *Biotechnol*
- 20 *Prog* **34**, 463–477.
- 21 41. Verma R, Holland MK, Temple-Smith P, Verma PJ (2012) *Theriogenology* **77**, 220–228.
- 22 42. von Bomhard A, Elsässer A, Ritschl LM, Schwarz S, Rotter N (2016) *PloS One* **11**,
- 23 e0149660.
- 24 43. Wani NA, Vettical BS, Hong SB (2017) *PloS One* **12**, e0177800.
- 25 44. Zhang L, Xue X, Yan J, Yan LY, Jin HH, Zhu XH, He ZZ, Liu J, Li R, Qiao J (2016) *Sci*
- 26 *Rep* **6**, 26326.
- 27
- 28

**CAPÍTULO 5 – EFFECTS OF INCUBATION TIME AND METHOD OF CELL
CYCLE SYNCHRONIZATION ON COLLARED PECCARY SKIN-DERIVED
FIBROBLAST CELL LINES**

(Artigo experimental)

ACEITO NOS ANNALS OF ANIMAL SCIENCE em 20/07/2020

QUALIS:

Quadriênio 2013-2016: Não classificado pela Medicina Veterinária/ Novo Qualis: A3

FATOR DE IMPACTO: 1,586

Effects of incubation time and method of cell cycle synchronization on collared peccary skin-derived fibroblast cell lines*

Alana Azevedo Borges¹, Maria Claudia dos Santos Luciano², Matheus Barbosa do Nascimento¹, Gabriela Pereira de Oliveira Lira¹, Fátima de Cássia Evangelista de Oliveira², Claudia Pessoa², Alessandra Fernandes Pereira^{1,♦}

¹Laboratory of Animal Biotechnology, Universidade Federal Rural do Semi-Árido, Mossoro, RN, Brazil.

²Experimental Oncology Laboratory, Universidade Federal do Ceará, Fortaleza, CE, Brazil.

♦Corresponding author: Alessandra Fernandes Pereira

Laboratory of Animal Biotechnology

Universidade Federal Rural do Semi-Árido

Av. Francisco Mota, 572, Mossoró, RN, 59625-900, Brazil

Phone: +55 84 3317 8361

E-mail address: alexsandra.pereira@ufersa.edu.br

Abbreviated title: Cell cycle of peccary fibroblasts

*Financed by Brazilian Council of Scientific Development (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brazil (CAPES, Financial Code 001). AF Pereira and C Pessoa are recipients of CNPq grants.

Abstract

The success of cloning by somatic cell nuclear transfer depends on the efficiency of nuclear reprogramming, with the cycle stage of the donor cell playing a crucial role. Therefore, the aim was to evaluate three different approaches for cell cycle synchronization: (i) serum starvation (SS) for 1 to 4 days, (ii) contact inhibition (CI) for 1 to 3 days, and (iii) using cell cycle regulatory inhibitors (dimethyl sulfoxide, cycloheximide, cytochalasin B, or 6-dimethylaminopurine) for 1 and 2 days, in terms of their effects on synchronization in G₀/G₁ phases and viability of collared peccary skin fibroblasts. Flow cytometry analysis revealed that SS for 4 days ($79.0\% \pm 1.6$) and CI for 3 days ($78.0\% \pm 1.4$) increased the percentage of fibroblasts in G₀/G₁ compared to growing cells GC, ($68.1\% \pm 8.6$). However, SS for 3 and 4 days reduced the viability evaluated by differential staining ($81.4\% \pm 0.03$ and $81.6\% \pm 0.06$) compared to growing cells (GC, $95.9\% \pm 0.06$). CI did not affect the viability at any of the analyzed time intervals. No cell cycle inhibitors promoted synchronization in G₀/G₁. These results indicate that CI for 3 days was the most efficient method for cell cycle synchronization in peccary fibroblasts.

Key words: G₀/G₁ phase, karyoplast, somatic cell nuclear transfer, wild mammals.

Introduction

The collared peccaries are wild mammals endemic to the American continent, where they play a crucial ecological role in maintaining their habitats (Desbiez et al., 2012). Among the various interactions of these animals with the ecosystem are their functions as seed dispersers and prey for larger mammals (Desbiez and Keuroghlian, 2009). Although their population is not considered to be of concern internationally (Gongora et al., 2011), in some biomes, such as Caatinga and the Atlantic forest, these individuals' populations are declining (Desbiez et al., 2012), thus requiring conservation strategies that ensure their maintenance in biodiversity.

Fortunately, this species' high adaptation capacity and zootechnical performance in captivity (Briceño-Méndez et al., 2016) suggest that reproductive biotechniques such as cloning by somatic cell nuclear transfer (SCNT) may be successfully used for their multiplication, as well as for research on embryonic development mechanisms of the species. SCNT involves embryo reconstruction by fusing the nucleus of a donor cell (karyoplast) with

an enucleated oocyte (cytoplasm), where the karyoplast in cell cycle phase G₀/G₁ is exposed to reprogramming by the oocyte, followed by fusion and activation of the reconstructed embryo (Loi et al., 2011). Usually, the success of SCNT depends on the efficiency of nuclear reprogramming and each species has different needs at each stage of the technique (Borges and Pereira, 2019). Although this process has not yet been completely clarified, it is known that the cell cycle phase of the karyoplast affects the results (Zhao et al., 2016). Therefore, the establishment of an appropriate cell synchronization protocol in G₀/G₁ is a crucial step in the development of SCNT usage in collared peccaries.

In recent years, our group has developed steps to implement the SCNT technique in collared peccaries. This path has been developed since the establishment of the ideal environment for receptor oocytes (cytoplasts, Borges et al., 2018b), and the development of an efficient oocyte artificial activation protocol to provide the development of the reconstructed embryo (Borges et al., 2020b). In parallel, we established somatic tissue banks (Borges et al., 2017, 2018a; Queiroz Neta et al., 2018), somatic cell banks (Lira et al., 2020), and, recently, we established skin-derived fibroblasts lines (Borges et al., 2020a). Therefore, now, we have evaluated the different cell cycle synchronization protocols of these lines, aiming at a higher percentage of cells in G₀/G₁.

Several cell cycle synchronization protocols in G₀/G₁ stages have been proposed. However, the cellular response to these protocols has been variable in domestic (Goissis et al., 2007) and wild mammals (Mahesh et al., 2012), necessitating the evaluation of each protocol for a given cell type and species (Wittayarat et al., 2013; Veraguas et al., 2017). Three main approaches have been employed for cell synchronization: serum starvation (SS, Kues et al., 2000), contact inhibition (CI, Boquest et al., 1999), and using cell cycle regulatory inhibitors (Goissis et al., 2007). While SS and CI act on the checkpoints by depriving the cells of adequate environmental or nutritional conditions (Kues et al., 2000), cell cycle inhibitors regulate specific biosynthetic processes such as repression or induction of cyclins (dimethyl sulfoxide, DMSO), protein synthesis inhibition (cycloheximide, CHX), cytoskeleton inhibition (cytochalasin B, CB), and protein kinase inhibition (6-dimethylaminopurine, 6-DMAP) to cause prolongation of the G₁ phase (Koo et al., 2009; Kretz et al., 2019).

The aim of this study was to evaluate SS, CI, and cell cycle regulatory inhibitors (DMSO, CHX, CB, or 6-DMAP) on synchronization in G₀/G₁ phases as well as the viability of collared peccary skin fibroblasts, in order to develop efficient cell cycle synchronization protocols for this species. We want to achieve this goal by defining appropriate methodologies for the last stage of the preparation of karyoplast to be used in the cloning of this species.

Materials and methods

Chemicals and bioethics

All chemicals were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Dulbecco's modified Eagle's medium (DMEM), fetal bovine serum (FBS), trypsin-EDTA, and antibiotic-antimycotic solution were obtained from Gibco-BRL (Carlsbad, CA, USA). The experiments were conducted in accordance with the Animal Ethics Committee (CEUA/UFERSA, no. 23091.001072/2015-92) and Chico Mendes Institute for Biodiversity Conservation (ICMBio, no. 48633-2).

Establishment and culture of fibroblasts

Skin tissue samples were obtained from the ear notch of four adult male collared peccaries at the Wild Animal Multiplication Center (CEMAS/UFERSA, no. 1478912, BR). The skin samples were cultured, and four fibroblast lines were previously established (Borges et al., 2020a). Subsequently, cells from four lines frozen in 10% DMSO, 10% FBS and 0.2 M sucrose were thawed, and 4th and 5th passage cells were used for this study.

Initially, cells were cultured in DMEM supplemented with 10% FBS and 2% antibiotic-antimycotic solution in a humid atmosphere containing 5% CO₂ at 38.5 °C. Prior to initiating cell cycle synchronization protocols, the cells were evaluated for their proliferative activity by obtaining the growth curve and determining the population doubling time (PDT). Cells were seeded in 12-well plates at a concentration of 1.0 x 10⁴ cells/mL. Cells from each well were counted at 24 h intervals for up to 216 h of culture. After each interval, the mean cell count was recorded; finally, the cell growth curve was generated, and the PDT was estimated based on these measurements (Roth, 2006).

1 Additionally, the percentage of living cells was determined by trypan blue exclusion
 2 assay (Borges et al., 2018a). Briefly, the cells were stained with 0.4% trypan blue in
 3 phosphate buffered saline (PBS) and counted using a hemocytometer.

4 5 *Cell treatments*

6 Evaluating three cell cycle synchronization methods, we examined the effects of
 7 serum starvation (SS), contact inhibition (CI), and cell cycle regulatory inhibitors (DMSO,
 8 CHX, CB, and 6-DMAP) treatments on the cell cycle stages of collared peccary fibroblasts in
 9 different incubation times. In each treatment group, cells without any treatment and with 70%
 10 confluence were used as a control (growing cells, GC). All treatments were performed in
 11 duplicate for each animal, producing a total of eight repetitions for each treatment and each
 12 incubation time.

13 14 *Experiment 1 – Effect of serum starvation*

15 SS was performed by replacing the culture medium with DMEM containing 0.5%
 16 FBS after the cell confluence had reached 70% in DMEM with 10% FBS (Caamaño et al.,
 17 2008). After the start of the SS treatment (day = 0), the stage of fibroblasts from each animal
 18 was analyzed on days 1, 2, 3 and 4. The culture medium was changed every 2 days.

19 20 *Experiment 2 – Effect of cell contact inhibition*

21 The effect of CI was studied for 3 days in culture after confluence had reached 90-
 22 100%. During the treatment of CI, the culture medium composed of DMEM and 10% FBS
 23 was changed every 2 days (Gómez et al., 2018).

24 25 *Experiment 3 – Effect of cell cycle regulatory inhibitors*

26 DMSO (0.5%, inhibitor of cytokinesis), CHX (7.5 µg/mL, protein synthesis inhibitor),
 27 CB (7.5 µg/mL, inhibitor of cytokinesis), and 6-DMAP (2 mM, protein kinase inhibitor)
 28 treatments were performed for 2 days of cell culture after the cell confluence reached 70%.
 29 After the start of treatment (day = 0), the stage of fibroblasts from each animal was analyzed
 30 on days 1 and 2 (Mahesh et al., 2012).

31 32 *Cell cycle synchronization analysis*

After the treatments, cells were trypsinized, centrifuged at 600×g for 10 min and resuspended in 1.0 mL of cold 70% ethanol for fixation. The cells were then maintained at -20 °C for 5 days. The fixed were washed in PBS for ethanol removal and each sample was centrifuged at 400×g for 10 min. Subsequently, cells were stained with 20 µg/mL of propidium iodide, 50 µg/mL of RNase were added, and samples were incubated at 4 °C for 50 min. After that, the samples were analyzed using Guava Easycyte flow cytometer (Guava Technologies, Stamford, Lincolnshire, United Kingdom). Data were obtained from 15,000 events from each sample. Histograms of red fluorescence vs counts were generated to evaluate the percentages of cells for each cell cycle phase (G₀/G₁, S, G₂/M) as well as the levels of sub G₀/G₁. The proportion of cells in each phase of the cell cycle and sub G₀/G₁ levels was assessed using MODFIT software version 5.0 (Verity, <https://www.vsh.com/products/mflt/index.asp>).

Cellular viability assessment

A cell aliquot was resuspended in 2 µg/mL of acridine orange and 10 µg/mL of ethidium bromide. Subsequently, the cells were evaluated using fluorescence microscopy at 480 nm, where 300 cells were counted at 200×. The cells were then classified into (i) viable cells, with a uniform light green nucleus; (ii) cells in initial apoptosis, with a non-uniform green nucleus; (iii) cells in late apoptosis, with a non-uniform bright orange nucleus, and (iv) necrotic cells, with a uniform orange nucleus (Kosmider et al., 2004). A fluorescence microscope was used to observe apoptotic changes in the stained cells, which were quantified using ImageJ software (National Institutes of Health, Bethesda, Maryland, USA).

Statistical analysis

Data were expressed as mean ± standard error (one fibroblast line/ one repetition) and analyzed using the GraphPad software (Graph-Pad Software Incorporation, La Jolla, CA, USA). All results were verified for normality by the Shapiro-Wilk test and homoscedasticity by Levene's test. Since data did not show a normal distribution, they were arcsine transformed and analyzed by ANOVA followed by Tukey test. Significance was set at P<0.05.

Results

Initially, thawed cells were morphologically normal (Figure 1A), with a percentage of $91.2\% \pm 1.2$ live cells after the thawing. Moreover, the cells showed a sigmoidal curve with the lag phase of adaptation up to day 2 followed by exponential and stationary growth, indicating that these cells were going through various phases of growth. Additionally, the decreasing phase was not observed (Figure 1B). The growth kinetics of the cells was demonstrated by counting total cells during 216 h of culture, resulting in an estimated PDT of 15.3 ± 1.4 h.





Figure 1. Culture of collared peccary fibroblast cells. (A) Fibroblasts at 70% confluency showing normal morphology. (B) Growth curve of fibroblasts.

Experiment 1 – Effect of serum starvation

After 4 days of SS, the percentage of G₀/G₁ phase cells was significantly higher ($P < 0.05$) when compared to GC, which was the control (Figure 2A–B, Table 1). Moreover, there was no change in the other phases of the cell cycle. Although the SS did not cause differences in sub G₀/G₁, when the levels of viability were evaluated by differential staining, 3 and 4 days of SS caused cell damage (Figure 3A).

1



2

3 Figure 2. Representative histograms of the flow cytometry analysis of collared peccary fibroblasts. (A) growing cells (GC, control). (B) serum
 4 starvation (SS). (C) contact inhibition (CI). (D) dimethylsulfoxide (DMSO). (E) cycloheximide (CHX). (F) cytochalasin B (CB). (G) 6-
 5 dimethylaminopurine (6-DMAP).

1
2 Table 1. Effect of serum starvation (SS) on the percentage of collared peccary fibroblasts in the G₀/G₁, S and G₂/M phases of the cycle

<i>Conditions</i>	<i>Cell cycle phase (%)</i>			
	<i>G₀/G₁</i>	<i>S</i>	<i>G₂/M</i>	<i>Sub G₀/G₁</i>
Growing cells (GC)	68.1 ± 8.5 ^a	17.6 ± 10.3 ^a	14.3 ± 1.8 ^a	15.9 ± 7.2 ^a
1 day SS	78.7 ± 2.3 ^{ab}	6.7 ± 1.5 ^a	14.6 ± 0.9 ^a	7.0 ± 1.8 ^a
2 day SS	78.1 ± 1.7 ^{ab}	7.2 ± 0.9 ^a	14.7 ± 1.5 ^a	10.5 ± 3.0 ^a
3 day SS	75.8 ± 2.9 ^{ab}	6.8 ± 2.2 ^a	17.4 ± 0.7 ^a	19.0 ± 11.3 ^a
4 day SS	79.0 ± 1.6 ^b	5.0 ± 0.8 ^a	16.0 ± 1.1 ^a	5.9 ± 2.2 ^a

3 Within a column, values with different superscripts differ (P<0.05).

4

5

6

7

8

9

10




Figure 3. Effect of different synchronization methods on the viability of collared peccary fibroblasts. (A) and (A') Effect of SS on fibroblast viability. (B) and (B') Evaluation of the viability of fibroblasts subjected to CI. (C) and (C') Effect of cell cycle inhibitors on the viability of synchronized fibroblasts. Within a bar at the same time, the values of a,b differ ($P < 0.05$). Within a bar at different times, A,B differ ($P < 0.05$). Fine arrow: viable cell; arrowhead: cell in initial apoptosis; Fat arrow: cell in late apoptosis; triangle: cell in necrosis.

1 *Experiment 2 – Effect of cell contact inhibition*

2 CI for 3 days significantly increased the proportion of fibroblasts in G₀/G₁ phase when
3 compared to GC (Figure 2C, Table 2) and decreased the proportion of cells in G₂/M phase
4 when compared to GC (P<0.05). However, neither evaluated time interval promoted
5 modifications in S cell cycle phase or levels of sub G₀/G₁ (P>0.05). Additionally, CI did not
6 affect the cell viability at any of the analyzed intervals (P>0.05, Figure 3B).

7
8 *Experiment 3 – Effect of cell cycle regulatory inhibitors*

9 Cell cycle inhibitors did not promote cell synchronization in G₀/G₁ at the time tested
10 when compared to GC for 1 and 2 days (Figure 2D-G, Table 3). While DMSO did not change
11 any of the evaluated parameters, CHX increased sub G₀/G₁ levels on both days and reduced
12 the percentage of cells in S phase after 2 days when compared to CG (P<0.05). After 2 days
13 of treatment with CHX, the proportion of fibroblasts in G₀/G₁ phase was higher when
14 compared to that after 1 day (P<0.05).

15
16 Moreover, CB reduced the percentage of cells in G₀/G₁ after 1 day, increasing the
17 percentage of cells in S phase when compared to GC (P<0.05). Additionally, treatment with
18 6-DMAP for 1 and 2 days significantly reduced the proportion of fibroblasts in S phase and
19 caused an increase in the proportion of cells in G₂/M phase. 6-DMAP also caused an increase
20 in sub G₀/G₁ levels when compared to GC (Table 3). Finally, after 1 day of treatment of 6-
21 DMAP, a reduction of viable cells was observed when compared to GC, although no
22 reduction was observed after 2 days of treatment (P<0.05, Figure 3C).

1 Table 2. Effect of contact inhibition (CI) on the percentage of collared peccary fibroblasts in the G₀/G₁, S and G₂/M phases of the cycle

<i>Conditions</i>	<i>Cell cycle phase (%)</i>			
	<i>G₀/G₁</i>	<i>S</i>	<i>G₂/M</i>	<i>Sub G₀/G₁</i>
Growing cells (GC)	70.8 ± 1.4 ^a	5.9 ± 0.8 ^a	23.3 ± 1.3 ^a	1.1 ± 0.2 ^a
1 day CI	75.6 ± 3.2 ^{ab}	7.3 ± 1.0 ^a	17.1 ± 2.2 ^{ab}	2.0 ± 0.7 ^a
2 day CI	75.5 ± 1.8 ^{ab}	6.9 ± 0.5 ^a	17.6 ± 1.5 ^{ab}	5.5 ± 3.5 ^a
3 day CI	78.0 ± 1.4 ^b	6.9 ± 1.2 ^a	15.1 ± 1.8 ^b	7.6 ± 5.3 ^a

2 Within a column, values with different superscripts differ (P<0.05).

1 Table 3. Effect of treatment with dimethyl sulfoxide (DMSO), cycloheximide (CHX), cytochalasin B (CB), or 6-dimethylaminopurin (6-DMAP)
 2 on the percentage of collared peccary fibroblasts in the G₀/G₁, S and G₂/M phases of the cycle

<i>Conditions</i>	<i>Cell cycle phase (%)</i>							
	<i>Treated for 1 day</i>				<i>Treated for 2 days</i>			
	<i>G₀/G₁</i>	<i>S</i>	<i>G₂/M</i>	<i>Sub G₀/G₁</i>	<i>G₀/G₁</i>	<i>S</i>	<i>G₂/M</i>	<i>Sub G₀/G₁</i>
Growing cells (GC)	75.4 ± 0.8 ^a	7.1 ± 0.7 ^b	17.5 ± 0.3 ^b	10.9 ± 2.9 ^b	75.4 ± 0.8 ^a	7.1 ± 0.7 ^a	17.5 ± 0.3 ^b	10.9 ± 2.9 ^b
0.5% DMSO	75.7 ± 0.9 ^a	5.9 ± 1.5 ^b	18.4 ± 1.0 ^b	10.9 ± 0.5 ^b	74.1 ± 0.5 ^a	8.0 ± 0.6 ^a	17.9 ± 0.3 ^b	17.7 ± 0.7 ^b
7.5 µg/mL CHX	72.7 ± 0.6 ^{abB}	9.1 ± 0.5 ^{abA}	18.2 ± 1.0 ^b	24.3 ± 2.7 ^c	77.4 ± 1.0 ^{aA}	5.4 ± 0.6 ^{bB}	17.2 ± 0.9 ^b	25.7 ± 3.3 ^a
7.5 µg/mL CB	68.3 ± 1.6 ^{bB}	12.5 ± 1.2 ^a	19.2 ± 0.7 ^{bA}	5.8 ± 2.9 ^d	74.4 ± 0.1 ^{aA}	9.8 ± 0.5 ^a	15.8 ± 0.5 ^{bB}	17.2 ± 1.3 ^b
2 mM 6-DMAP	73.5 ± 1.5 ^a	0.0 ± 0.0 ^{cA}	26.5 ± 1.5 ^{aA}	46.7 ± 1.1 ^{aA}	74.6 ± 1.5 ^a	3.7 ± 0.6 ^{bB}	21.7 ± 1.2 ^{aB}	27.3 ± 5.7 ^{aB}

3 Within a column, ^{a,b} values with different superscripts differ (P<0.05). Within a row, ^{A,B} values with different superscripts differ (P<0.05).

4

1 Discussion

2 An appropriate cell synchronization protocol in G_0/G_1 is a crucial step in the
3 development of cloning by SCNT in collared peccaries. Previously, our group developed
4 somatic resource banks aiming at their application in the conservation of this species (Borges
5 et al., 2017, 2018a, 2020a,b; Lira et al., 2020; Queiroz Neta et al., 2018a). With this new
6 study, we have developed a suitable protocol for somatic cell synchronization, the last step in
7 the preparation of karyoplasts, i.e., somatic nucleus donor cells. To our knowledge, this is the
8 first work to elucidate nuclear reprogramming in cells of this species.

9
10 Here, we identified that using CI for 3 days was more efficient for cell cycle
11 synchronization when compared to SS for 1 to 4 days and treatment with cell cycle inhibitors
12 (DMSO, CHX, CB, or 6-DMAP) for 1 and 2 days. This result may be related to a high cell
13 density where CI favors the regulation of reactive oxygen species (ROS) as well as activates
14 the coactivator-1 α (PGC1 α), which functions as a key regulator of energy expenditure,
15 involved in reduction of ROS and protection of cells from oxidative stress (Yang et al., 2018).
16 In studies with the Asian golden cat (*Catopuma temminckii*), leopard (*Panthera pardus*), and
17 Siamese cat, researchers observed that CI for 5 days promoted an increase of more than 80%
18 in the percentage of cells in G_0/G_1 when compared to non-synchronized cells, without
19 increase in apoptotic cells (Wittayarat et al., 2013). In swine, domestic species
20 phylogenetically close to the collared peccaries, cells were efficiently synchronized by CI for
21 3 to 4 days and resulted in 20.4% of zygotes developing into blastocysts after cloning by
22 SCNT (Lee et al., 2019). Moreover, for wood bison (*Bison bison*), the CI was adequate for
23 cell synchronization in G_0/G_1 phase, which resulted in 19.2% of developed blastocysts (Seaby
24 et al., 2013). Therefore, the CI has been efficient in producing clone embryos.

25
26 Although SS promoted cell synchronization after 4 days in collared peccaries, this
27 treatment had a negative effect on cell viability after 3 and 4 days. In general, the
28 synchronization of the G_0/G_1 cell cycle of mammalian cells via SS is due to the response to
29 absence or presence of mitogens to continue the cell cycle during the onset of G_1 phase, so
30 when these cells are in absence of mitochondrial growth factors that would be offered by the
31 serum, they accumulate in a state of a $2n$ DNA content (Coller et al., 2007). In porcine cells,
32 the SS for 1 to 3 days increased the proportion of cells at G_0/G_1 phase up to 77.9-80.2% (Kues
33 et al., 2000). However, these authors also observed cell damage by visualization of DNA

fragmentation. Similar to our study, goat fibroblasts were found to be efficiently synchronized in G₀/G₁ by both CI and SS (Dalman et al., 2010); the number of cells in apoptosis increased by 39.8% in 3 days in SS whereas in cells at CI there was no increased apoptosis (8.7%).

The negative effect of SS was also observed on embryonic development. The cell synchronization using SS resulted in embryos with a higher degree of apoptosis and increased ROS when compared to CI (Lee et al., 2016). Likewise, observed that SS led to high Bak (pro-apoptotic) and low Bcl-xL (anti-apoptotic) expression while the CI showed equilibrium in the increase in Bak and Bcl-xL expression (Park et al., 2010). Hence, the increase in apoptosis in cells synchronized in G₀/G₁ by 4 days with SS may have a negative effect on the later embryonic development in collared peccaries.

In the present study, no cell cycle inhibitors promoted synchronization in G₀/G₁ in collared peccaries and 6-DMAP had negative effects on cell viability with increase in sub G₀/G₁ levels and reduction of viable cells. A study with goral (*Naemorhedus caudatus*) fibroblasts also indicated no cell cycle synchronization in G₀/G₁ using CHX (60.1%), 6-DMAP (3.9%) and CB (10.8%) in cultures for 1 day (Hashem et al., 2006). Porcine fibroblasts treated with CHX after 2 days showed an increased percentage of dead cells when compared to that after 1 day (Goissis et al., 2007). In our study, we observed an increase in cells in sub G₀/G₁ at both times. Also, was evaluated the synchronization in G₀/G₁ for porcine fibroblasts in different passages (5th, 10th and 15th) and observed that CHX when used in cells from the 5th passage was unable to synchronize cells when compared to that observed with cells from the 10th and 15th passages, presenting higher number of cells in G₀/G₁ only in 10th passage fibroblasts (Goissis et al., 2007). Possibly, 10th passage presented a higher proportion of senescent cells, becoming more susceptible to synchronization by this inhibitor because of the lower levels of cyclin D, the target mechanism of CHX in the cell cycle (Goissis et al., 2007). The treated cells in this study were from the 4th and 5th passage; this was probably why the cells did not present a positive synchronization result using CHX.

Furthermore, DMSO did not change any of the evaluated parameters in collared peccary cells. Studies in porcine, brown bear (*Ursus arctos pyrenaicus*), and goral cells have shown that cell cycle synchronization with DMSO is concentration dependent. While for porcine cells, 1% DMSO (88.7%) was more efficient than 0.5% (86.5%) and 2.5% (86.0%)

DMSO after 2 days (Boquest et al., 1999), brown bear cells were efficiently synchronized with DMSO concentrations of 1-3% (79.0-79.1%) after 2 days (Caamaño et al., 2008). Nevertheless, in goral cells, while 0.5% (74.8%) and 1% DMSO (75.2%) caused synchronization in G₀/G₁ after 1 day, no effect was observed using 2.5% DMSO (Hashem et al., 2006). Therefore, these results show that the effect of DMSO is linked to the concentration used and the sensitivity of the species to this chemical. Additionally, although DMSO may be efficient for synchronization in G₀/G₁, the use of high concentrations of this agent may cause negative effects on cells (Caamaño et al., 2008).

In summary, this study identified that both CI for 3 days and SS for 4 days promoted the synchronization of somatic cells in collared peccaries; however, the percentage of viable cells was maintained only in cells synchronized by CI. Therefore, CI for 3 days was considered the most efficient method. With the cells being subjected to less chemical exposure and metabolic stress, CI can be efficiently applied to cell cycle synchronization in collared peccary fibroblasts. Thus, we established the last step of the preparation for the use of these fibroblast as karyoplast for application in somatic cell nuclear transfer of collared peccary, with potential application to conservation of the species.

Acknowledgments

The authors thank the CEMAS/UFERSA for providing biological material.

References

- Boquest, A.C., Day, B.N., Prather, R.S. (1999). Flow cytometric cell cycle analysis of cultured porcine fetal fibroblast cells. *Biol. Reprod.*, 60: 1013–1019.
- Borges, A.A., Lima, G.L., Queiroz Neta, L.B., Santos, M.V.O., Oliveira, M.F., Silva, A.R., Pereira, A.F. (2017). Conservation of somatic tissue derived from collared peccaries (*Pecari tajacu* Linnaeus, 1758) using direct or solid-surface vitrification techniques. *Cytotechnology*, 69: 643–654.
- Borges, A.A., Lira, G.P.O., Nascimento, L.E., Queiroz Neta, L.B., Santos, M.V.O., Oliveira, M.F., Pereira, A.F. (2018a). Influence of cryopreservation solution on the *in vitro* culture of skin tissues derived from collared peccary (*Pecari tajacu* Linnaeus, 1758). *Biopreserv. Biobank.*, 16: 77–81.
- Borges, A.A., Lira, G.P.O., Nascimento, L.E., Santos, M.V.O., Oliveira, M.F., Silva, A.R.,

- 1 Pereira, A.F. (2020a). Isolation, characterization, and cryopreservation of collared peccary
2 skin-derived fibroblast cell lines. *PeerJ*, 8, e9136.
- 3 Borges, A.A., Pereira, A.F. (2019). Potential role of intraspecific and interspecific cloning in
4 the conservation of wild mammals. *Zygote*, 27: 111–117.
- 5 Borges, A.A., Santos, M.V.O., Queiroz Neta, L.B., Oliveira, M.F., Silva, A.R., Pereira, A.F.
6 (2018b). *In vitro* maturation of collared peccary (*Pecari tajacu*) oocytes after different
7 incubation times. *Pesq. Vet. Bras.*, 38: 1863–1868.
- 8 Borges, A.A., Santos, M.V.O., Nascimento, L.E., Lira, G.P.O., Praxedes, E.A., Oliveira, M. F.,
9 Silva, A.R., Pereira, A.F. (2020b). Production of collared peccary (*Pecari tajacu* Linnaeus,
10 1758) parthenogenic embryos following different oocyte chemical activation and *in vitro*
11 maturation conditions. *Theriogenology*, 142: 320–327.
- 12 Briceño-Méndez, M., Naranjo, E., Mandujano, S., Altricher, M., Reyna-Hurtado, R. (2016).
13 Responses of two sympatric species of peccaries (*Tayassu pecari* and *Pecari tajacu*) to
14 hunting in Calakmul, Mexico. *Trop. Conserv. Sci.*, 9: 1940082916667331.
- 15 Caamaño, J.N., Rodriguez, A., Salas, A., Munoz, M., Diez, C., Prather, R., Gomez, E. (2008).
16 Flow cytometric cell cycle analysis of cultured brown bear fibroblast cells. *Cell Biol.*
17 *Int.*, 32: 855–859.
- 18 Coller, H.A. (2007). What's taking so long? S-phase entry from quiescence versus proliferation.
19 *Nat. Rev. Mol. Cell Biol.*, 8: 667–670.
- 20 Dalman, A., Eftekhari-Yazdi, P., Valojerdi, M.R., Shahverdi, A., Gourabi, H., Janzamin, E.,
21 Fakheri, R., Sadeghian, F., Hasani, F. (2010). Synchronizing cell cycle of goat fibroblasts by
22 serum starvation causes apoptosis. *Reprod. Domest. Anim.*, 45: 46–53.
- 23 Desbiez, A.L., Keuroghlian, A. (2009). Can bite force be used as a basis for niche separation
24 between native peccaries and introduced feral pigs in the Brazilian Pantanal? *Mammalia*, 73:
25 369–372.
- 26 Desbiez, A.L.J., Keuroghlian, A., Beisiegel, B.M., Medici, E.P., Gatti, A., Pontes, A.R. M.,
27 Pinho, G.M. (2012). Avaliação do risco de extinção do cateto *Pecari tajacu* Linnaeus, 1758,
28 no Brasil. *Biodivers. Bras.*, 3: 74–83.
- 29 Goissis, M.D., Caetano, H.V.A., Marques, M.G., Barros, F.R.O., Feitosa, W.B., Milazzotto,
30 M.P., Binelli, M., Assumpção, M.E.O.A., Visintin, J.A. (2007). Effects of serum deprivation
31 and cycloheximide on cell cycle of low and high passage porcine fetal fibroblasts. *Reprod.*
32 *Domest. Anim.*, 42: 660–663.
- 33 Gómez, N.A., Ramírez, M.M., Ruiz-Cortés, Z.T. (2018). Primary fibroblast cell cycle

- 1 synchronization and effects on handmade cloned (HMC) bovine embryos. Ciênc. Anim.
2 Bras., 19: 1–17.
- 3 Gongora, J., Biondo, C., Cooper, J.D., Taber, A., Keuroghlian, A., Altrichter, M., Nascimento,
4 F.F., Chong, A.Y., Miyaki, C.Y., Bodmer, R., Mayor, P., González, S. (2011). Revisiting the
5 species status of *Pecari maximus* van Roosmalen et al., 2007 (Mammalia) from the Brazilian
6 Amazon. Bonn Zoolog. Bull., 60: 95–101.
- 7 Hashem, M.A., Bhandari, D.P., Kang, S.K., Lee, B.C., Suk, H.W. (2006). Cell cycle analysis of
8 *in vitro* cultured goral (*Naemoredus caudatus*) adult skin fibroblasts. Cell Biol. Int., 30:
9 698–703.
- 10 Koo, O.J., Hossein, M.S., Hong, S.G., Martinez-Conejero, J.A., Lee, B.C. (2009). Cell cycle
11 synchronization of canine ear fibroblasts for somatic cell nuclear transfer. Zygote, 17: 37–
12 43.
- 13 Kosmider, B., Zyner, E., Osiecka, R., Ochocki, J. (2004). Induction of apoptosis and necrosis in
14 A549 cells by the cis-Pt (II) complex of 3-aminoflavone in comparison with cis-DDP. Mutat.
15 Res., Genet. Toxicol. Environ. Mutagen., 563: 61–70.
- 16 Kretz, R., Wendt, L., Wongkanoun, S., Luangsa-Ard, J.J., Surup, F., Helaly, S.E., Noumeur,
17 S.R., Stadler, M., Stradal, T.E. (2019). The effect of cytochalasins on the actin cytoskeleton
18 of eukaryotic cells and preliminary structure–activity relationships. Biomolecules, 9: 73–87.
- 19 Kues, W., Anger, M., Carnwath, J., Paul, D., Motlik, J., Niemann, H. (2000). Cell cycle
20 synchronization of porcine fetal fibroblasts: effects of serum deprivation and reversible cell
21 cycle inhibitors. Biol. Reprod., 62: 412–419.
- 22 Lee, J.H., Chun, J.L., Kim, K.J., Kim, E.Y., Kim, D.H., Lee, B.M., Han, K.W., Park, K.S., Lee,
23 K.B., Kim, M.K. (2016). Effect of acteoside as a cell protector to produce a cloned dog. PloS
24 One, 11: e0159330.
- 25 Lee, Y., Shim, J., Ko, N., Kim, H.J., Park, J.K., Kwak, K., Choi, K. (2019). Effect of alanine
26 supplementation during *in vitro* maturation on oocyte maturation and embryonic
27 development after parthenogenesis and somatic cell nuclear transfer in pigs. Theriogenology,
28 127: 80–87.
- 29 Lira, G.P.O., Borges, A.A., Nascimento, M.B., Aquino, L.V.C., Oliveira, M.F., Silva, A.R.,
30 Pereira, A.F. (2020). Cryopreservation of collared peccary (*Pecari tajacu* Linnaeus, 1758)
31 somatic cells is improved by sucrose and high concentrations of fetal bovine serum.
32 CryoLetters, 41: 271–279.
- 33 Loi, P., Modlinski, J.A., Ptak, G. (2011). Interspecies somatic cell nuclear transfer: a salvage

- 1 tool seeking first aid. *Theriogenology*, 76: 217–228.
- 2 Mahesh, Y.U., Rao, B.S., Katari, V.C., Komjeti, S., Christo, D., Lakshmikantan, U., Pawar,
3 R.M., Shivaji, S. (2012). Cell cycle synchronization of Bison (*Bos gaurus*) fibroblasts
4 derived from ear piece collected *post-mortem*. *Reprod. Domest. Anim.*, 47: 799–805.
- 5 Park, H., Koo, O., Kwon, D., Kang, J., Jang, G., Lee, B. (2010). Effect of roscovitine-treated
6 donor cells on development of porcine cloned embryos. *Reprod. Domest. Anim.*, 45: 1082–
7 1088.
- 8 Queiroz Neta, L.B., Lira, G.P.O., Borges, A.A., Santos, M.V.O., Silva, M.B., Oliveira, L.R.M.,
9 Silva, A.R., Oliveira, M.F., Pereira A.F. (2018). Influence of storage time and nutrient
10 medium on recovery of fibroblast-like cells from refrigerated collared peccary (*Pecari tajacu*
11 Linnaeus, 1758) skin. *In Vitro Cell. Dev. Biol. Anim.*, 54: 486–495.
- 12 Roth, V. (2006). Doubling time computing [Program for calculation]. Retrieved from
13 www.doubling-time.com/compute.php. Accessed 08 September 2019
- 14 Seaby, R., Alexander, B., King, W., Mastromonaco, G. (2013). *In vitro* development of bison
15 embryos using interspecies somatic cell nuclear transfer. *Reprod. Domest. Anim.*, 48: 881–
16 887.
- 17 Veraguas, D., Gallegos, P., Castro, F., Alvarez, L.R. (2017). Cell cycle synchronization and
18 analysis of apoptosis-related gene in skin fibroblasts from domestic cat (*Felis silvestris*
19 *catus*) and kodkod (*Leopardus guigna*). *Reprod. Domest. Anim.*, 52: 881–889.
- 20 Wittayarat, M., Thongphakdee, A., Saikhun, K., Chatdarong, K., Otoi, T., Techakumphu, M.
21 (2013). Cell cycle synchronization of skin fibroblast cells in four species of family Felidae.
22 *Reprod. Domest. Anim.*, 48: 305–310.
- 23 Yang, S., Hwang, S., Jang, J., Kim, M., Gwak, J., Jeong, S.M. (2018). PGC1 α is required for
24 the induction of contact inhibition by suppressing ROS. *Biochem. Biophys. Res. Commun.*,
25 501: 739–744.
- 26 Zhao, Q., Wu, Y., Shan, Z., Bai, G., Wang, Z., Hu, J., Liu, L., Li, T., Shen, J., Lei, L. (2016).
27 Serum starvation-induced cell cycle synchronization stimulated mouse rDNA transcription
28 reactivation during somatic cell reprogramming into iPSCs. *Stem Cell Res. Ther.*, 7: 112–
29 122.
- 30
- 31
- 32
- 33
- 34

**CAPÍTULO 6 – *IN VITRO* MATURATION OF COLLARED PECCARY (*Pecari tajacu*
LINNAEUS, 1758) OOCYTES AFTER DIFFERENT INCUBATION TIMES
(Artigo experimental)**

PUBLICADO NA PESQUISA VETERINÁRIA BRASILEIRA

QUALIS: Qualis 2013-2016: A2/ Novo Qualis: A4

FATOR DE IMPACTO: 0,1474

DOI: <https://doi.org/10.1590/1678-5150-pvb-5471>

***In vitro* maturation of collared peccary (*Pecari tajacu* Linnaeus, 1758) oocytes after different incubation times¹**

Alana A. Borges², Maria V. O. Santos², Luiza B. Queiroz Neta², Moacir F. Oliveira³, Alexandre R. Silva⁴, Alexsandra F. Pereira^{2*}

ABSTRACT.- Borges A.A., Santos M.V.O, Queiroz Neta L.B, Oliveira M.F, Silva A.R & Pereira A.F. 2017. [*In vitro* maturation of collared peccary (*Pecari tajacu* Linnaeus, 1758) oocytes after different incubation times.] *Pesquisa Veterinária Brasileira* 00(0):00-00. Federal Rural University of Semi-Arid, Av. Francisco Mota, 572, Mossoró, RN, 59625-900, Brazil. E-mail: alexandra.pereira@ufersa.edu.br

Oocyte *in vitro* maturation (IVM) is the first step of the *in vitro* reproductive technologies that enables mature oocytes to be generated *ex vivo* and after used for embryo production. In this sense, the establishment of culture environment, as oocyte incubation time, is essential for the success of the IVM. Therefore, the study was carried out to investigate the relationship between the meiotic potential and the IVM times of collared peccary oocytes, wild mammals of great commercial and ecological interest. Thus, ovaries were collected of females derived from captivity and transported to the laboratory within 1 h of slaughtering. The oocytes derived from follicles (3–6 mm in diameter) were recovered by aspirated and sliced. Good quality oocytes (evenly granulated cytoplasm with a least one layer of surrounding *cumulus* cells) were selected and subjected to culture in TCM 199 supplemented with 10 µg/mL FSH, 10% FBS and 100 µM cysteamine at 38.5°C, 5% CO₂ and maximum humidity for 24 or 48 h. After the incubation period, the nuclear status, the presence of first polar body and the expansion of *cumulus* cells of oocytes were assessed. The data obtained were analyzed by Fisher exact test ($P < 0.05$). A total of four sessions (2–3 females per session) were performed, resulting in eighteen aspirated and sliced ovaries with normal morphological characteristics. An oocyte recovery rate of approximately 83.1% (59/71) was obtained with 3.3 oocytes/ovary and 2.3 viable oocytes/ovary. After different incubation times, differences ($P < 0.05$) were observed between 24 h and 48 h for expansion of the *cumulus* cells (38.1% vs. 100%), presence of first polar body (52.4% vs. 90.5%) and nuclear status in second metaphase (19.0% vs. 76.2%), respectively. In conclusion, 48 h is suitable time for the *in vitro* maturation of oocytes derived from collared peccaries when compared to the time of 24 h,

according to the meiotic potential observed. Additional studies should be conducted to improve the quality of the oocyte culture environment, as medium composition, aiming to obtain viable mature oocytes for other *in vitro* biotechnologies.

INDEX TERMS: Assisted Reproductive Technologies, meiotic competence, nuclear maturation, wild mammals.

¹ Received on

Accepted for publication on

²Laboratory of Animal Biotechnology. Federal Rural University of Semi-Arid (UFERSA), Av. Francisco Mota, 572, Mossoró, RN, 59625-900, Brazil. *Corresponding author: alexsandra.pereira@ufersa.edu.br

³Laboratory of Applied Animal Morphophysiology. Federal Rural University of Semi-Arid, Av. Francisco Mota, 572, Mossoró, RN, 59625-900, Brazil.

⁴Laboratory of Animal Germplasm Conservation. Federal Rural University of Semi-Arid, Av. Francisco Mota, 572, Mossoró, RN, 59625-900, Brazil.

RESUMO.- [Maturação *in vitro* de oócitos de cateto (*Pecari tajacu* Linnaeus, 1758) após diferentes períodos de incubação.] A maturação *in vitro* (MIV) oocitária é a primeira etapa das tecnologias reprodutivas *in vitro* que permite que oócitos maturados sejam gerados *ex vivo* e depois usados para a produção de embriões. Nesse sentido, o estabelecimento do ambiente de cultivo, como o período de incubação de oócitos, é essencial para o sucesso da MIV. Portanto, o estudo foi realizado para investigar a relação entre o potencial meiótico e os períodos de MIV de oócitos derivados de catetos, mamíferos selvagens de grande interesse comercial e ecológico. Para tanto, os ovários foram coletados de fêmeas derivadas de cativeiro e transportados ao laboratório dentro de 1 h após o abate. Os oócitos derivados de folículos (3–6 mm de diâmetro) foram recuperados por aspiração e fatiados. Oócitos de boa qualidade (citoplasma uniformemente granuloso com pelo menos uma camada circundante de células *cumulus*) foram selecionados e submetidos ao cultivo em TCM 199 suplementado

com 10 µg/mL de FSH, 10% de SFB e 100 µM de cisteamina a 38,5°C, 5% de CO₂ e umidade máxima por 24 ou 48 h. Após o período de incubação, o estado nuclear, a presença do primeiro corpúsculo polar e a expansão das células do *cumulus* dos oócitos foram avaliados. Os dados obtidos foram analisados pelo teste exato de Fisher ($P < 0,05$). Um total de quatro sessões (2–3 fêmeas por sessão) foi realizado, resultando em dezoito ovários aspirados e fatiados com características morfológicas normais. Uma taxa de recuperação oocitária de aproximadamente 83,1% (59/71) foi obtida com 3,3 oócitos/ovário e 2,3 oócitos viáveis/ovário. Após diferentes períodos de incubação, diferenças ($P < 0,05$) foram observadas entre 24 e 48 h para a expansão das células *cumulus* (38,1% vs. 100%), presença de primeiro corpúsculo polar (52,4% vs. 90,5%) e estado nuclear na segunda metáfase (19,0% vs. 76,2%), respectivamente. Em conclusão, 48 h é o período adequado para a maturação *in vitro* de oócitos derivados de catetos quando comparado ao tempo de 24 h, de acordo com o potencial meiótico observado. Estudos adicionais devem ser conduzidos para melhorar a qualidade do ambiente de cultivo oocitário, como a composição de meio, objetivando obter oócitos maturados viáveis para outras biotecnologias *in vitro*.

TERMOS DE INDEXAÇÃO: Tecnologias Reprodutivas Assistidas, competência meiótica, maturação nuclear, mamíferos silvestres.

Introduction

Oocyte *in vitro* maturation (IVM) is the first step of the *in vitro* reproductive technologies that enables mature oocytes to be generated *ex vivo* (Rahman et al. 2008) and further destined to systems of *in vitro* embryo production (IVEP), as *in vitro* fertilization (IVF), intracytoplasmic sperm injection (ICSI) and somatic cell nuclear transfer (SCNT). All these biotechniques are interesting tools for applying the reproductive management (Kumar et al. 2016) and biodiversity conservation (Moulavi et al. 2017).

The capability of embryonic development depends on successful IVM, since it stimulates the oocyte to achieve the meiotic competence required (Lee et al. 2014) through the nuclear and cytoplasmic events that promote the synthesis of RNA and proteins (Sirard et al. 2006). Therefore, the IVM quality is a determinant factor for the success of the IVEP (Zhang et al. 2015, Park et al. 2017). In this sense, the establishment of culture environment, as

oocyte incubation time, should be clarified, once all IVM events require a specific time that could vary according to the peculiarities of each species (Higaki et al. 2016).

The optimization of IVM conditions for collared peccaries is interesting under various concerns. Although this species is not threatened (IUCN, 2017), their phylogenetic proximity to endangered species as the *Catagonus wagneri* (Keuroghlian et al. 2014), as well as the commercial importance on their meat and leather (Nogueira-Filho et al. 2004), have been leading to the development of studies focused on their reproductive physiology (Maia et al. 2014, Miranda-Moura et al. 2016). Based on these studies, it was possible to know that specifically for females, they have symmetrical and oval ovaries with a relatively smooth surface, which can present irregular appearance due to corpus luteum or follicles (Garcia et al. 2009). Moreover, the ovaries are enveloped by an ovarian bursa (Guimarães et al. 2012), as also observed in swine (Męczyński 1974). Additionally, collared peccaries' estrous cycle lasts 21 days, with a follicular phase of 6 days and 15 days for the luteal phase (Maia et al. 2014). On the other hand, these individuals have a lower ovulation rate (2.3 follicles) when compared to swine (Sowls 1997) and usually produce 1.7–1.9 offspring per female after 141–151 days of gestation (Mayor et al. 2005).

Therefore, in view of the need for studies related to the reproductive biotechniques in collared peccaries, the initial point has been to observe reproductive relationship with close domestic phylogenetic species, as swine and others artiodactyls. Thus, similarities were observed for folliculogenesis (Guimarães et al. 2012), hormonal profiles (Ahuja-Aguirre et al. 2017) and follicular histological characteristics (Lima et al. 2013) between collared peccaries and swine and/or artiodactyla species, as buffalos.

In this context, IVM time used for buffalo oocytes is of 24 h (Kadoom et al. 2014), while maturation time of 48 h is used for porcine oocytes (Pereyra-Bonnet et al. 2008). Moreover, it had observed that a suboptimal culture environment for porcine oocytes may promote nuclear maturation without the structures reaching cytoplasmic maturation (Abeydeera 2002), confirming that the IVM time to be precise and defined (Sirard et al. 2006). Therefore, this is first study was carried out to investigate the relationship between the meiotic potential and the IVM times of collared peccary oocytes. This information will be the first about the IVM of oocytes derived from collared peccary, aiming to direct pathways to

obtain an embryo *in vitro* of this species. This step can serve as a basis for both enlightening studies of embryogenesis and physiological aspects of the animal. In addition, it may be the starting point for advanced biotechnology that can increase zootechnical potential and ways to ensure effective tools that can be used for conservation of phylogenetically near species.

Materials and Methods

Animal ethics and care

The experiment was conducted in accordance with the Committee for Ethics in Animal Use of Federal Rural University of Semi-Arid (CEUA/UFERSA; no. 23091.001072/2015-92) and the Chico Mendes Institute for Biodiversity Conservation (ICMBio; no. 48633-2). The animals used in this research belong to the Centre of Multiplication of Wild Animals (CEMAS/UFERSA, Mossoró, RN, Brazil; 5°10'S, 37°10'W), registered at the Brazilian Institute of Environment and Renewable Natural Resources (IBAMA) as a scientific breeding site (no. 1478912). The breeder contains 200 collared peccaries on average. Annually, a programmed slaughter is conducted for population control and the biological material is destined for several experiments.

Reagents and media

All reagents and media used were obtained from Sigma-Aldrich (St. Louis, MO, USA), but fetal bovine serum (FBS) was obtained from Gibco-BRL (Carlsbad, CA, USA). When necessary, the pH was adjusted to 7.2–7.4 for all media. Media were filtered using 0.22 µm system (Corning, NY, USA).

Oocyte collection, selection and in vitro maturation

Ovaries were obtained from nine adult female collared peccaries (age 4–10 years). Eighteen ovaries were removed from the ovarian bursa and transported to the laboratory in 0.9% (w/v) NaCl solution at 37°C for up to 1 h. Antral follicles (3–6 mm) were aspirated with 21 gauge needle attached to a 5.0 mL syringe. After the aspiration, for a greater oocyte recovery, ovaries were sliced with a scalpel blade in 100x20 mm plates. All follicular fluid was pooled in conical tubes for ten minutes. The sediment was diluted in the oocyte collection medium composed of TCM 199 culture medium (M2520) supplemented with 2.2 g/L bicarbonate (S5761), 0.2 mM sodium pyruvate (P45622), 1% antibiotic-antimycotic solution (A5955) and 10% FBS (12657-029). Thus, *cumulus*-oocyte complexes (COCs) were selected

according to the number of *cumulus* cells layers and homogeneity of the cytoplasm, being classified in four grades. Grade 1 (3-5 *cumulus* cells layers and homogeneous cytoplasm); Grade 2 (1-2 *cumulus* cells layers and homogeneous cytoplasm); Grade 3 (partially denuded cytoplasm and slightly heterogeneous); Grade 4 (denuded and heterogeneous cytoplasm), according to Kumar et al. (2016). Only grade 1 and 2 COCs were considered viable to be matured.

After three washes in oocyte collection medium, the COCs were incubated in 100 μ L drops of maturation medium composed of TCM 199 with 2.2 g/L bicarbonate, 0.2 mM sodium pyruvate, 1% antibiotic-antimycotic solution and supplemented 10 μ g/mL follicle stimulating hormone (Folltropin[®]-V, Bioniche, Canada), 100 μ M cysteamine (M9768) and 10% FBS. All drops were covered with mineral oil and cultured at 38.5°C in humidified atmosphere of 5% CO₂ and divided in two groups, 24 or 48 h.

Assessment of in vitro maturation

After the IVM, COCs were evaluated by the analysis of *cumulus* cells expansion using a stereomicroscope at 20–40X magnification. Thus, oocytes with *cumulus* cells were considered matured (Appeltant et al. 2015). Subsequently, *cumulus* cells were removed by pipetting the oocytes in oocyte collection medium and denuded structures were verified for the presence of the first polar body under stereomicroscope at 20–40X magnification. Thus, oocytes with first polar body were considered matured (Pereyra-Bonnet et al. 2008).

The nuclear status of oocytes was determined by marking it with Hoechst 33342 (B2261). For this, denuded oocytes were fixed in paraformaldehyde buffered in 4% phosphate buffered solution (PBS) for 30 min and then washed in PBS with 0.4% bovine serum albumin (BSA). Afterward, Hoechst 33342 (10 μ g/mL) labeling was performed for 15 min and exposed under UV light in fluorescence microscope. Finally, oocytes presenting nucleus in metaphase II (metaphase plate, MII) were considered matured and those in other nuclear phases (prophase I, metaphase I, anaphase I and telophase I) were considered immature (Appeltant et al. 2015).

Data analysis

Data were obtained from four sessions, with two a three females per session, totaling in nine experimental animals. For statistical analysis, the Graphpad Instat 3.06 software (GraphPad Software Inc., La Jolla, CA, USA) was used. Mean numbers in percentage $\pm$ standard error of the matured oocytes in two different times (24 vs. 48 h) were compared using Fisher exact test ($P < 0.05$) for expansion of *cumulus* cells, presence of the first polar body and nuclear status.

Results

Under gross evaluation, the bursa surrounded all peccaries' ovaries, which were symmetric, presenting an oval shape. Moreover, they had a uniform appearance on most of its surface (Fig. 1 A); nevertheless, some elevations were observed derived from large follicles or the presence of corpus luteum (Fig. 1 B).




Fig. 1. Macroscopic aspects of collared peccary ovaries. **(A)** Uniform appearance with small follicles. **(B)** Presence of large follicles and corpus luteum. The arrowhead indicates a small follicle and arrow shows a large follicle.

Afterward all sessions, a total of 18 ovaries resulted in 59 immature oocytes from follicles aspirated and sliced follicles with an estimated recovery rate of 83.1% (approximately, 59/71; oocytes recovered/follicles aspirated) and 3.3 oocytes per ovary. Of these, 42 structures (71.2%, 42 structures viable/59 recovered structures) were classified as viable according to the number of *cumulus* cells layers and homogeneity of the cytoplasm, resulting in 2.3 viable oocytes per ovary, (Fig. 2 A).




Fig. 2. *In vitro* maturation of oocytes of collared peccary. **(A)** Immature oocytes before IVM (100X); **(B)** *Cumulus* cell expansion after 24 h IVM (100X); **(C)** *Cumulus* cell expansion after 48 h IVM (100X); **(D)** Oocyte with extrusion of first polar body (400X); **(E)** Oocyte in MII stage (400X).

After the IVM, the time of 48 h was more adequate than 24 h for the IVM in all evaluated parameters, as *cumulus* cells expansion ($P = 0.0002$), presence of the first polar body ($P = 0.0148$) and nuclear status in second metaphase ($P = 0.0003$) (Table 1). Moreover, a 100% *cumulus* cells expansion was observed for oocytes cultured for 48 h (Fig. 2 C) when compared to 24 h (Fig. 2 B). Additionally, oocytes with first polar body (Fig. 2 D) and MII presence (Fig. 2 E) were evidenced in more of 50% and 19% of the oocytes, respectively.

1 **Table 1.** *In vitro* maturation of collared peccary oocytes after different incubation times.

Evaluation of <i>in vitro</i> maturation	Incubation times (in hours)	
	24	48
Expansion of <i>cumulus</i> cells	38.1 ± 1.4 (8/21) ^b	100.0 ± 0.0 (21/21) ^a
Presence of the first polar body	52.4 ± 2.2 (11/21) ^b	90.5 ± 2.0 (19/21) ^a
Nuclear status in second metaphase	19.0 ± 1.4 (4/21) ^b	76.2 ± 1.3 (16/21) ^a

2 Mean in % ± standard error (*n*). ^{a,b} in same row differ (P < 0.05).

3

4

5

1 Discussion

2 The results indicated that the most appropriate incubation period for collared
3 peccaries' oocytes achieve meiotic competence was 48 h. This time is similar to that observed
4 for porcine oocytes (Pereyra-Bonnet et al. 2008). In this sense, we can infer that similar
5 follicular characteristics described between collared peccaries and swine (Lima et al. 2013)
6 justify this relationship, since the *in vitro* conditions for maturation should mimic the *in vivo*
7 conditions that occur in the females.

8
9 Moreover, other peculiar feature of collared peccary oocytes similar to the porcine
10 oocytes is the presence of cytoplasmic lipid droplets (Lima et al. 2013). This aspect is crucial
11 for the establishment of the appropriate requirements for IVM, evidencing that further IVM
12 protocols should be adapted from porcine to peccaries' oocytes. Nevertheless, a factor that
13 still needs to be further investigated by the similarity that collared peccary oocytes had in
14 relation to porcine oocytes is that the IVM in swine oocytes have their altered medium to
15 adapt to the variable requirements of IVM (Gruppen 2014) and this step may improve the
16 quality of matured oocytes derived from collared peccaries. Thus, the tracks of such
17 similarities between peccaries and porcine oocytes indicate a starting point, but not the direct
18 application of protocols without further experimentation, considering that differences occur
19 between these species for follicular morphology (Lima et al. 2013).

20
21 On the other hand, the follicular development in collared peccaries was also similar to
22 that found for other artiodactyl species regarding the stages of folliculogenesis, i.e.
23 recruitment, selection and dominance (Guimarães et al. 2012) when compared to buffalos
24 (Manik et al. 2002) and camelids (Bausiouni 2007). Nevertheless, despite this similarity, the
25 24 h IVM time that is frequently used for buffaloes (Kadoom et al. 2014) was not suitable for
26 oocyte maturation in the peccaries. The IVM time to guarantee the formation of the
27 metaphase plate is essential to ensure that the oocytes possess the ability to develop quality
28 embryos (Gruppen 2014). In this sense, the 24 h of IVM for collared peccary oocytes was not
29 long enough for the organization of the metaphase plate (19.0%), while the majority of
30 oocytes were able to reach metaphase II in 48 h of IVM (76.2%). As developmental
31 competence is also reached in cytoplasmic maturation, which occurs after meiotic maturation
32 and refers to the ability of the oocyte to be fertilized and to block polyspermy, this type of
33 maturation can be visualized by migration of the cortical granules to the oocyte periphery and

by the expansion of *cumulus* cells (Milakovic et al. 2015). The analysis of *cumulus* cells expansion at 24 and 48 h showed different degrees of expansion (38.1% vs. 100.0%, respectively); these may be related to cytoplasmic maturation, being an indicator of successful maturation (Abeydeera 2002). Therefore, 24 h of IVM is not sufficient for IVM of collared peccary oocytes (Fig. 2B).

Several important factors for the IVM success are dependent of the incubation time, which can influence the final quality of the embryonic development (Miyoshi et al. 2002). The signaling pathway of 3',5' cyclic adenosine monophosphate (cAMP) is highly time-dependent and restricted to time mechanisms, which is what triggers germinal vesicle breakdown, so nuclear maturation depends on cAMP so the oocyte can reach metaphase II (Sirard et al. 2006). Likewise, another time dependent factor is the secretion of steroid hormones by COCs derived small and large follicles during IVM. Topfer et al. (2016) observed that oocytes derived from small follicles have a lesser ability to support estradiol synthesis in the first half of IVM (28 h) and to suppress progesterone synthesis at the end of IVM (48 h) compared with oocytes from large follicles. Thus, the steroid hormone secretion by COCs is time-dependent and therefore, the meiotic maturation time should to be precise and defined (Sirard et al. 2006; Topfer et al. 2016).

Moreover, the *cumulus* cells play a bidirectional role for oocyte development. One of the factors that contribute to this role is their mitochondria and mitochondrial DNA, besides contributing to the passage of nutrients and growth factors to the oocyte (Pawlak et al. 2016). Additionally, it is known that the *cumulus* cells play a regulatory role for cAMP, showing great relevance for oocyte maturation (Gruppen 2014). Therefore, the clear expansion of *cumulus* cells with 48 h (Fig. 2C) in this study ensured the optimal performance of the cells during IVM. Further adjustments of cAMP levels during maturation may improve fertilization capacity and oocyte development (Appeltant et al. 2015).

Finally, other factors, as follicles size and technique employed for oocyte collection, may influence the IVM conditions (Sun et al. 2001). In this study, 3–6 mm follicles aspirated and matured for 48 h allowed the obtaining of oocytes to reach meiotic competence. Moreover, the recovery rate obtained can be due to the use of two methods of oocytes collection, first the follicular aspiration and then the slicing technique. The slicing technique

allows the increase oocyte recovery, especially when there is limited availability of ovaries (Ferraz et al. 2016). Moreover, viability rate of 71.2% can be attributed to the good development of immature oocytes in the follicular environment (Grupen 2014) and the efficiency of the methods for oocyte recovery.

In conclusion, 48 h IVM for collared peccary is the most suitable incubation period for oocytes reaching the meiotic competence when compared to 24 h time. This was the first study with *in vitro* maturation, in an attempt to improve a protocol of IVEP for this species. Although this time has been defined in the present work, further details for the improvement of IVM, as medium composition, should be considered not only to achieve meiotic competence but also to guarantee appropriate embryonic development.

Acknowledgement.- This study was supported by National Counsel of Technological and Scientific Development (CNPq) and Coordination for the Improvement of Higher Education Personnel (CAPES). The authors thank the CEMAS/UFERSA for providing the animals. AR Silva and MF Oliveira were recipients of CNPq grants.

References

- Abeydeera L.R. 2002. *In vitro* production of embryos in swine. *Theriogenology*. 57:257–273.
- Ahuja-Aguirre C., López-deBuen L., Rojas-Maya S. & Hernández-Cruz, B.C. 2017. Progesterone and estradiol profiles in different reproductive stages of captive collared peccary (*Pecari tajacu*) females assessed by fecal metabolites. *Anim. Reprod. Sci.* *in press*.
- Appeltant R., Beek J., Vandenbergh L., Maes D. & Van Soom A. 2015. Increasing the cAMP concentration during *in vitro* maturation of pig oocytes improves *cumulus* maturation and subsequent fertilization *in vitro*. *Theriogenology*, 83:344–352.
- Basiouni G.F. 2007. Follicular wave pattern, folliculogenesis and assisted reproductive techniques in the non-pregnant female dromedary camel (*Camelus dromedarius*): A review. *J. Biol. Sci.* 7:1038–1045.
- Ferraz M.S., Moraes Júnior F.D.J., Feitosa M.L.T., Almeida H.M.D., Bezerra D.D.O., Pessoa G.T., Albuquerque D.M.N & Carvalho, M.A.M.D. 2016. Ovarian slicing technique to obtain oocytes from agoutis (*Dasyprocta prymnolopha*). *Pesq. Vet. Bras.* 36:204–208.
- Garcia A.R., Kahwage P.R. & Ohashi O.M. 2009. Aspectos reprodutivos de caititus (*Tayassu tajacu*). *Rev. Bras. Reprod. Anim.* 33:71–81.

- 1 Grupen C.G. 2014. The evolution of porcine embryo *in vitro* production. *Theriogenology*.
2 81:24–37.
- 3 Guimarães D.A., Garcia S.C.G., Pantoja M.A.F., Silva S.D.S.B., Albuquerque N.I. & Le Pendu
4 Y. 2012. Ovarian folliculogenesis in collared peccary: *Pecari tajacu* (Artiodactyla:
5 Tayassuidae). *Rev. Biol. Trop.* 60: 437–455.
- 6 Higaki S., Kishi M., Koyama K., Nagano M., Katagiri S., Takada T. & Takahashi Y. 2016.
7 Early germinal vesicle breakdown is a predictor of high preimplantation developmental
8 competent oocytes in mice. *Zygote*. 25:41–48.
- 9 IUCN (2017) IUCN Red List of Threatened Species. Version 2015.4. www.iucnredlist.org
10 <http://dx.doi.org/10.2305/IUCN.UK.2015-2.RLTS.T4015A72587993.en>. Accessed 13 April
11 2017.
- 12 Kadoom A.K., Abdel-Khalek A.E., Shamiah S.M., El-Sharawy M.E. & El-Razek I.A. 2014. *In*
13 *vitro* maturation, fertilization and development of prepubertal and mature buffalo oocytes.
14 *Egyptian J. Anim. Prod.* 51:65–69.
- 15 Keuroghlian A., Andrade Santos M.D.C. & Eaton D.P. 2014. The effects of deforestation on
16 white-lipped peccary (*Tayassu pecari*) home range in the southern Pantanal. *Mammalia*. 1–
17 7.
- 18 Kumar S., Devi B., Purkayastha A., Bharti P.K., Doley S. & Kadirvel G. 2016. Porcine ovarian
19 biometry, oocyte retrieval and quality of oocytes under different seasons in north east hill
20 region. *Indian J. Anim. Sci.* 86:300–303.
- 21 Lee J., Lee Y., Park B., Elahi F., Jeon Y., Hyun S.H. & Lee E. 2014. Developmental
22 competence of IVM pig oocytes after SCNT in relation to the shrinkage pattern induced by
23 hyperosmotic treatment. *Theriogenology*. 81:974–981.
- 24 Lima G.L., Santos E.A.A., Luz V.B., Rodrigues A.P.R. & Silva A.R. 2013. Morphological
25 characterization of the ovarian preantral follicle population of collared peccaries (*Tayassu*
26 *tajacu* Linnaeus, 1758). *Anat. Histol. Embryol.* 42:304–311.
- 27 Maia K.M., Peixoto G.C., Campos L.B., Bezerra J.A.B., Ricarte A.R., Moreira N., Oliveira
28 M.F. & Silva A.R. 2014. Estrus cycle monitoring of captive collared peccaries (*Pecari*
29 *tajacu*) in semiarid conditions. *Pesq. Vet. Bras.* 34:1115–1120.
- 30 Manik R.S., Palta P., Singla S.K. & Sharma V. 2002. Folliculogenesis in buffalo (*Bubalus*
31 *bubalis*): a review. *Reprod. Fertil. Dev.* 14:315–325.

- 1 Mayor P., López-Gatius F. & López-Béjar, M. 2005. Integrating ultrasonography within the
2 reproductive management of the collared peccary (*Tayassu tajacu*). Theriogenology.
3 63:1832–1843.
- 4 Męczyński S. 1974. Morphohistological structure of female genital organs in sousliks. Acta
5 theriologica. 19:91–106.
- 6 Milakovic I., Jeseta M., Hanulakova S., Knitlova D., Hanzalova K., Hulinska P., Machal L.,
7 Kempisty B., Antosik P. & Machatkova, M. 2015. Energy status characteristics of porcine
8 oocytes during *in vitro* maturation is influenced by their meiotic competence. Reprod. Dom.
9 Anim. 50:812–819.
- 10 Miranda-Moura M.T.M., Oliveira G.B., Peixoto G.C.X., Pessoa J.M. & Papa P.C. 2016.
11 Morphology and vascularization of the corpus luteum of peccaries (*Pecari tajacu*, Linnaeus,
12 1758) throughout the estrous cycle. Arq. Bras. Med. Vet. Zootec. 68:87–96.
- 13 Miyoshi K., Rzucidlo S.J., Pratt S.L. & Stice S.L. 2002. Utility of rapidly matured oocytes as
14 recipients for production of cloned embryos from somatic cells in the pig. Biol. Reprod.
15 67:540–545.
- 16 Moulavi F., Hosseini S.M., Tahhaie-Vash N., Ostadhosseini S., Hosseini S.H., Hajinasrollah
17 M., Asghari M.H., Gourabi H., Shahverdi A., Vosough A.D. & Nasr-Esfahani M.H. 2017.
18 Interspecies somatic cell nuclear transfer in Asiatic cheetah using nuclei derived from *post-*
19 *mortem* frozen tissue in absence of cryo-protectant and *in vitro* matured domestic cat
20 oocytes. Theriogenology. 90:197–203.
- 21 Nogueira-Filho S., Nogueira S., Mendes A. & Jori F. 2004. A large-scale commercial farming
22 of collared peccary (*Tayassu tajacu*) in north-eastern Brazil. Game and Wildlife Science.
23 21:413–420.
- 24 Park S.J., So K.H. & Hyun, S.H. 2017. Effect of zeaxanthin on porcine embryonic development
25 during *in vitro* maturation. J. Biomed. Res. 31:1–8.
- 26 Pawlak P., Chabowska A., Malyszka N. & Lechniak D. 2016. Mitochondria and mitochondrial
27 DNA in porcine oocytes and *cumulus* cells - A search for developmental competence
28 marker. Mitochondrion. 27:48–55.
- 29 Pereyra-Bonnet F., Fernández-Martín R., Olivera R., Jarazo J., Vichera G., Gibbons A. &
30 Salamone D. 2008. A unique method to produce transgenic embryos in ovine, porcine,
31 feline, bovine and equine species. Reprod. Fertil. Dev. 20:741–749.
- 32 Rahman A.N.M.A., Abdullah R.B. & Wan-Khadajah W.E. 2008. *In vitro* maturation of oocytes
33 with special reference to goat: a review. Biotechnology. 7:599–611.

- 1 Sirard M.A., Richard F., Blondin P. & Robert C. 2006. Contribution of the oocyte to embryo
2 quality. *Theriogenology*. 65:126–136.
- 3 Sowls L.K. 1997. Javelinas and other peccaries: The Biology, Management and Use. Texas
4 A&M University Press, College Station, Texas, pp. 86–104.
- 5 Sun Q.Y., Lai L., Bonk A., Prather R.S. & Schatten H. 2001. Cytoplasmic changes in relation
6 to nuclear maturation and early embryo developmental potential of porcine oocytes: effects
7 of gonadotropins, *cumulus* cells, follicular size, and protein synthesis inhibition. *Mol.*
8 *Reprod. Dev.* 59:192–198.
- 9 Topfer D., Ebeling S., Weitzel J.M. & Spannbrucker A.C. 2016. Effect of follicle size on *in*
10 *vitro* maturation of pre-pubertal porcine *cumulus* oocyte complexes. *Reprod. Dom. Anim.*
11 51:370–377.
- 12 Zhang J.Y., Jiang Y., Lin T., Kang J.W., Lee J.E. & Jin D.I. 2015. Lysophosphatidic acid
13 improves porcine oocyte maturation and embryo development *in vitro*. *Mol. Reprod. Dev.*
14 82:66–77.
- 15

**CAPÍTULO 7 – PRODUCTION OF COLLARED PECCARY (*Pecari tajacu*
LINNAEUS, 1758) PARTHENOGENIC EMBRYOS FOLLOWING DIFFERENT
OOCYTE CHEMICAL ACTIVATION AND *IN VITRO* MATURATION
CONDITIONS**

(Artigo experimental)

PUBLICADO NA THERIOGENOLOGY

QUALIS: Quadriênio 2013-2016: A2/ Novo Qualis: A1

FATOR DE IMPACTO: 2,299

DOI: <https://doi.org/10.1016/j.theriogenology.2019.10.016>

Production of collared peccary (*Pecari tajacu* Linnaeus, 1758) parthenogenic embryos following different oocyte chemical activation and *in vitro* maturation conditions

Alana Azevedo Borges^a, Maria Valéria de Oliveira Santos^a, Lucas Emanuel Nascimento^a, Gabriela Pereira de Oliveira Lira^a, Érika Almeida Praxedes^a, Moacir Franco de Oliveira^b, Alexandre Rodrigues Silva^c, Alexsandra Fernandes Pereira^{a*}

^aLaboratory of Animal Biotechnology, Federal Rural University of Semi-Arid, Mossoro, RN, Brazil. ^bLaboratory of Applied Animal Morphophysiology, Federal Rural University of Semi-Arid, Mossoro, RN, Brazil. ^cLaboratory of Animal Germplasm Conservation, Federal Rural University of Semi-Arid, Mossoro, RN, Brazil.

*Corresponding author: Alexsandra Fernandes Pereira

Laboratory of Animal Biotechnology

Federal Rural University of Semi-Arid

Av. Francisco Mota, 572, Mossoró, RN, 59625-900, Brazil

Phone: +55 84 3317 8361

E-mail address: alexsandra.pereira@ufersa.edu.br

Abstract

To optimize the protocols for assisted reproductive techniques (ARTs) in collared peccary (*Pecari tajacu* Linnaeus, 1758), we evaluated various conditions for oocyte *in vitro* maturation (IVM) and chemical activation. Initially, we assessed the IVM rates, *cumulus*-oocyte complex (COC) quality, and oocyte morphometry in the absence or presence of epidermal growth factor (EGF). There was no difference between the COCs matured in absence or presence of EGF for the expansion of *cumulus* cells ($97.6\% \pm 1.2$ vs. $100\% \pm 0.0$), presence of first polar body ($65.9\% \pm 1.2$ vs. $70.5\% \pm 1.8$), nuclear status in second metaphase ($62.5\% \pm 11.6$ vs. $68.4\% \pm 4.9$), cytoplasmic maturation ($100.0\% \pm 0.7$ vs. $75.0\% \pm 0.7$), reactive oxygen species levels (0.5 ± 0.2 vs. 0.3 ± 0.1), and mitochondrial membrane potential (1.1 ± 0.2 vs. 1.1 ± 0.1). However, the zona pellucida thickness of matured COCs was reduced in the presence of EGF. Thus, the EGF group was used for further experiments. The oocytes were artificially activated with ionomycin and four secondary activator combinations [6-dimethylaminopurine (6D), 6D and cytochalasin B (6D + CB), cycloheximide (CHX), and CHX and CB (CHX + CB)]. The effect of immature COCs based on *cumulus* cell layers and cytoplasm homogeneity (GI and GII or GIII COCs) on embryonic development and quality was evaluated. There was no difference in the cleavage rates among the groups of secondary activators. The cleavage rates of embryos derived from GI/GII and GIII COCs were greater than 72.2% and 25.0%, respectively. Moreover, treatment with CHX showed a reduction in the cleavage rate of embryos derived from GIII COCs when compared to the cleavage rate of embryos derived from GI/GII COCs ($P < 0.05$). Nevertheless, higher rates of blastocyst/total GI and GII COCs were observed in the 6D group ($27.6\% \pm 0.3$) compared to CHX group ($6.9\% \pm 0.3$). Additionally, only 6D treatment resulted in the production of embryos derived from GIII COCs ($25.0\% \pm 0.2$). The percentage of the ICM/total cell ratio was also greater in blastocysts derived from 6D ($42.5\% \pm 19.0$), 6D + CB ($37.9\% \pm 21.9$), and CHX + CB ($43.8\% \pm 19.6$) groups when compared to CHX ($3.6\% \pm 0.1$) group. Thus, the combination of ionomycin and 6D could produce collared peccary embryos by activation of both GI/GII COCs and GIII COCs. These optimized IVM conditions using EGF and chemical activation using ionomycin and 6D in collared peccaries form the first steps for establishing ARTs to conserve this species.

Keywords: Wildlife; Somatic cell nuclear transfer; Artificial activation; Embryo development; Epidermal growth factor.

1. Introduction

In the last decade, a rapid decline in the population of various mammalian species was reported in the South American countries [1]. The collared peccary (*Pecari tajacu* Linnaeus, 1758), a wild pig, is found only in the Americas and is classified as the Least Concern species [2]. However, the species is already extinct in eastern and southern Argentina and is declining in some biomes, such as Caatinga and the Atlantic forest [1]. Generally, this species plays an important role in the ecosystem as a seed dispersing agent. Thus, the maintenance of these animals in captivity is essential for their sustainable use and conservation [3].

The collared peccary populations can be conserved through assisted reproductive techniques (ARTs), such as cloning by somatic cell nuclear transfer (SCNT) and *in vitro* fertilization (IVF). Cloning by SCNT enables the conservation of species as it can be used for increasing the species population [4], basic research [5], and obtaining induced pluripotent cells [6]. Therefore, establishment of all the stages involved in SCNT can aid in the optimal use of this technology for species conservation [5].

There are various steps involved in cloning via SCNT. *In vitro* maturation (IVM) and artificial oocyte activation are the fundamental steps for the development of SCNT technique to obtain cytoplasmic donor cells (or cytoplasm) and evaluate their developmental competence. Recently, we had demonstrated that the collared peccary *cumulus*-oocyte complexes (COCs) require maturation times of greater than 24 h for complete meiotic development, which resulted in an IVM rate of 76.2% [7]. However, developmental competence was not evaluated in this study. It is important to understand the development of *in vitro*-matured oocytes to blastocysts.

Moreover, the optimization of IVM medium composition can potentially result in higher IVM rates. In some mammals (swine [8]; sheep [9]; canine [10]), supplementation of IVM medium with 10 ng/mL of epidermal growth factor (EGF) increases the meiotic development and subsequent embryonic development. In pigs, EGF along with gonadotropins enhances oocyte nuclear and cytoplasmic maturation [11]. Although this domestic species is phylogenetically close to collared peccary [12], there are differences in the reproductive features between these species, such as ovulation rate [13, 14] and litter size [14]. Although swine and collared peccary are phylogenetically close, they exhibit substantial phylogenetic

1 distance. Thus, direct extrapolations between the collared peccary and swine were not
2 possible. Therefore, we aimed to specifically evaluate the effects of EGF on the collared
3 peccary oocyte development. Additionally, different species may exhibit differential
4 responses, as observed in the collared peccary semen cryopreservation. Treatment with 20%
5 low-density lipoproteins (LDL) improved post-thaw sperm motility and plasma membrane
6 integrity of collared peccaries [15], whereas treatment with LDL at concentrations above 10%
7 did not result in a beneficial effect on porcine semen cryopreservation [16].

8
9 To the best of our knowledge, we are not aware of any published reports on the
10 establishment of oocyte artificial activation protocols in collared peccaries. Artificial
11 activation of oocytes is also a critical step of SCNT [5] and the activation procedures vary
12 among species (bovine and equine [17]; goat [18]; swine [19]). This indicates that the
13 response to these artificial activation protocols also may vary depending on the species.
14 Previously, Campos Junior et al. [20] artificially activated the collared peccary oocytes using
15 ionomycin and 6-dimethylaminopurine (6D), which resulted in 10% four-cell embryo stage.
16 Increasing the number of oocytes and optimizing the artificial activation protocols may
17 potentially promote higher rates of embryonic development.

18
19 An efficient artificial activation protocol promotes *in vitro* embryonic development
20 competencies by decreasing the level of the maturation promoter factor (MPF) through Ca^{2+}
21 oscillation [18]. Thus, different chemical activation methods may be employed using
22 ionomycin as the primary activator, which allows Ca^{2+} transport and induces calcium influx
23 by activation of endogenous entry pathways [21]. Moreover, primary activation can be used in
24 combination with secondary activators, such as protein phosphorylation inhibitors (6D) that
25 inhibit the protein kinase activity and promote mitosis, and protein synthesis inhibitors
26 (cycloheximide or CHX) that inhibit the production of cyclin B, which is a regulatory
27 component of MPF [22]. Additionally, cytochalasin B (CB) has been used to prevent the
28 fragmentation of embryos [23] and increase the rate of diploid embryos as well as to
29 effectively inhibit the secondary polar body extrusion [24].

30
31 Generally, COCs used for IVM and chemical activation are selected based on *cumulus* cell
32 layers and cytoplasm homogeneity (Grade I to IV). Only GI and GII COCs are used for IVM,
33 which are considered more appropriate according to morphological classification described

above. However, due to the reduced availability of oocytes in wild mammals, evaluation of development in GIII COCs is important to increase the quantities of oocytes in the subsequent stages of SCNT. In some mammals, similar molecular patterns were observed between GI/GII and GIII COCs [25, 26]. Thus, this study aimed to evaluate various conditions for IVM and chemical activation of collared peccary oocytes.

2. Materials and methods

All experiments were conducted in accordance with the Animal Ethics Committee of the Federal Rural University of Semi-Arid (Opinion N° 23091.001072/2015-92) and Chico Mendes Institute for Biodiversity Conservation (ICMBio, N° 48633-2). Unless otherwise stated, all chemicals were obtained from Sigma Chemical Co. (St. Louis, MO, USA).

2.1. Cumulus-oocyte complex (COC) collection

Twenty collared peccaries (n = 20) aged 25.1 ± 2.5 months housed at the Centre of Multiplication of Wild Animals (CEMAS/UFERSA, Mossoró, RN, Brazil; 5°10'S, 37°10'W) and registered at the Brazilian Institute of Environment and Renewable Natural Resources (IBAMA, N° 1478912) were used in this study. The animals were stimulated with 600 IU PG600® (400 IU eCG and 200 IU hCG, MSD Saúde Animal, São Paulo, SP, Brazil) before COC collection, following the methods of Peixoto et al. [27] with modifications. At day 4 post-hormone administration, the animals were euthanized and the ovaries were excised. The ovaries were transported to the laboratory within 1 h and maintained in saline solution (NaCl, 0.9%) supplemented with 0.05 mg/mL penicillin at 35–37°C.

At the laboratory, all visible follicles were classified as small (< 0.3 cm), medium (0.3–0.5 cm), or large (> 0.5 cm) [28]. The follicles with 0.2–0.6 cm diameter were aspirated for COC collection using a 21 G needle attached to a 5.0 mL syringe containing COC collection medium (CCM; TCM199 (Gibco-BRL, Carlsbad, CA, USA) supplemented with 0.2 mM sodium pyruvate, 10% fetal bovine serum (FBS, serum (FBS, Gibco-BRL), and 1% antibiotic-antimycotic solution).

2.2. COC classification and IVM

The COCs were observed under a stereomicroscope and classified into four grades following the classification system proposed by Kumar et al. [29], which was based on the

number of *cumulus* cell layers and homogeneity of the cytoplasm: Grade I (3–5 *cumulus* cells layers and homogeneous cytoplasm); Grade II (1–2 *cumulus* cells layers and homogeneous cytoplasm); Grade III (partially denuded and slightly heterogeneous cytoplasm); Grade IV (denuded and heterogeneous cytoplasm). Only Grade I/II and Grade III COCs were used for IVM in separate drops, according to the experimental design.

The COCs were divided into drops (15–30 COCs per 100 μ L) and the drops were covered with mineral oil. The COCs were incubated for 44 h at 38.5°C and 5% CO₂ in a humidified atmosphere. The IVM medium comprised CCM supplemented with 20 μ g/mL FSH/LH (Pluset[®], Hertape Calier, Juatuba, MG, Brazil), 100 μ M cysteamine, and an absence or presence of 10 ng/mL EGF.

2.3. Evaluation of *cumulus* cells

After IVM, *cumulus* cell expansion was evaluated using a stereomicroscope. The COCs exhibiting *cumulus* cell expansion were considered mature. The *cumulus* cell expansion was scored as follows: score 4, total expansion; score 3, partial expansion; score 2, slight expansion; score 1, minimum expansion; score 0, no expansion [30].

The *cumulus* cells were removed by successive pipetting. The viability of *cumulus* cells was evaluated by staining the cells with trypan blue (0.2%) for 2 min. The viable cells were unstained, whereas the non-viable cells were stained blue. All cells were counted in the 4 outer quadrants of the Neubauer chamber [31].

2.4. Assessment of nuclear maturation

The denuded oocytes were observed under a stereomicroscope to assess the presence of first polar body (1PB). The oocytes exhibiting 1PB were considered mature. To assess the nuclear stage, denuded oocytes were fixed in 4% paraformaldehyde prepared in phosphate buffer saline (PBS) for 30 min. Next, the oocytes were washed with PBS containing 0.4% bovine serum albumin (BSA) fraction V. The cells were then stained with Hoechst 33342 (10 μ g/mL) for 15 min and observed under a fluorescent microscope. The oocytes with nucleus at metaphase II (metaphase plate, MII) stage and 1PB were considered mature, while those in other nuclear phases were considered immature.

2.5. Quantification of intracellular ROS levels by a dichlorofluorescein assay

For evaluation of oocyte quality after IVM, reactive oxygen species (ROS) levels in the oocytes were determined using the fluorescent probe, 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA; Invitrogen, Carlsbad, CA, USA), according to Santos et al. [31]. Briefly, denuded oocytes were washed twice with PBS and incubated in 500 μ L of 5 μ M H₂DCFDA at 38.5°C and 5% CO₂ for 30 min. The stained oocytes were washed twice with PBS and imaged using a fluorescence microscope (Olympus BX51TF, Tokyo, Japan). The fluorescence signal intensity (pixels) was quantified using the ImageJ software (version 1.49v, Java 1.8.0_201, Wayne Rasband, U.S. National Institutes of Health, Bethesda, MD, USA; website: <http://rsb.info.nih.gov/ij/download.html>). The background signal intensity was subtracted from the fluorescent intensity values of the treated oocytes. Immature oocytes were assessed as the calibrator. The relative expression levels were calculated by dividing the measured value of each treatment micrograph with the mean value of the calibrator.

2.6. Assessment of mitochondrial membrane potential ($\Delta\Psi_m$) and cytoplasmic maturation

The $\Delta\Psi_m$ was measured by incubating the oocytes with MitoTracker Red® (CMXRos, Invitrogen) at 500 nM for 30 min [31]. The staining procedures and evaluation of images were similar to those used for the quantification of ROS levels. Additionally, the cytoplasmic maturation of the labeled oocytes was evaluated based on the mitochondrial distribution as follows: peripheral (immature oocytes), dispersed (mature oocytes), and transitional (between peripheral and dispersed distributions; partially mature oocytes).

2.7. Morphometric evaluation of oocytes after maturation

The denuded oocytes were observed under an inverted microscope for assessing the oocyte morphometric parameters. The images were captured using an inverted microscope (Nikon TS100, Tokyo, Japan) equipped with a camera and image processing software. The morphometric parameters of the captured images were analyzed in the ImageJ software using the scale bar as an arbitrary scale for pixel analysis. The measured morphometric parameters included outer oocyte diameter (ZPO), zona pellucida thickness (ZPT), inner oocyte diameter (ZPI), and ooplasm diameter (OD), as described by Saadeldin et al. [32]. Thus, perivitelline space (PVS) diameter, inner oocyte area (IA), ooplasm area (OA), and PVS area were mathematically calculated based on previous parameters using the following formulae: IA =

3.14 x (ZPI/2)²; OA = 3.14 x (OD/2)², PVS diameter = ZPI – OD and PVS area = IA – OA.
Diameters were measured in µm and areas in µm².

2.8. Artificial activation and in vitro development (IVD) of embryos

After IVM, the *cumulus* cells were removed by pipetting the oocytes in the presence of hyaluronidase for up to 2 min. The oocytes with 1PB were activated using 5 µM ionomycin (I24222; Gibco-BRL) prepared in CCM for 4 min at 37°C. Next, the oocytes were washed and incubated with drops of secondary activators at 38.5°C and 5% CO₂ for 3 h, according to the experimental design. Subsequently, the oocytes were washed and incubated in 50 µL drops of IVD medium covered with mineral oil. Synthetic oviductal fluid (SOF) supplemented with 0.2 mM sodium pyruvate, 0.2 mM L-glutamine, 0.34 mM sodium citrate, 2.8 mM myo-inositol, 2% essential amino acid solution, 1% non-essential amino acid solution, 1% antibiotic-antimycotic solution, 5.0 mg/mL BSA, and 2.5% FBS was employed as the IVD medium. The day of activation was considered as D0 and the cells were cultured until D7. On D3, 50% of the culture medium was replaced with fresh medium.

2.9. Evaluation of blastocysts and differential staining of inner cell mass and trophectoderm cells

After seven days of IVD, the blastocyst cells were fixed in ethanol and labeled with Hoechst 33342 (10 µg/mL) and propidium iodide (PI) (10 µg/mL) for 15 min at 38.5°C. Next, the cells were washed twice with PBS and placed on glass slides in glycerol droplets. A coverslip was placed over the droplet and the images were captured under UV light at 330–385 nm using a fluorescent microscope. The embryos were visualized individually and the nuclei were counted in the captured fluorescent images. The nuclei of the inner cellular mass (ICM) appeared blue as they were labeled only with Hoechst 33342, while those of the trophectoderm (TE) appeared red or pink as they were labeled with both PI and Hoechst 33342. The numbers of ICM, TE, and total cells were quantified using the ImageJ software [33].

2.10. Experimental design

To evaluate various conditions for IVM and chemical activation of collared peccary oocytes, two experiments were performed. Initially, we assessed the IVM rates, COC quality, and oocyte morphometric parameters after IVM in the absence or presence of 10 ng/mL of

EGF (experiment 1). Furthermore, we activated the oocytes with ionomycin and four secondary activator combinations (6D, 6D and CB, CHX, and CHX and CB). The effect of immature COC grades (GI and GII or GIII COCs) on the embryonic development and quality was evaluated (experiment 2). For the second experiment, only the oocytes grown under optimal IVM conditions were used.

2.10.1. Experiment 1: effects of EGF supplementation on IVM of collared peccary COCs

The COCs were washed with the medium and four replicates (two females per replicate) were used to assess the effect of EGF on IVM. GI and GII COCs derived from eight females were matured in the absence (EGF negative group) or presence of 10 ng/mL of EGF (EGF positive group). Meiotic competence was measured by visualization of the 1PB and nuclear stage (MII). Moreover, oocytes were evaluated for cytoplasmic maturation based on mitochondrial distribution using MitoTracker Red[®]. Additionally, the *cumulus* cells were assessed and the ROS levels and $\Delta\Psi_m$ were quantified. The matured oocytes were compared by morphometric evaluation according to the ZPO, ZPT, OD, PVS area, and PVS diameter.

2.10.2. Experiment 2: effects of secondary activators and COC quality on the development of collared peccary parthenogenetic embryos

After 44 h of IVM, matured GI/GII and GIII COCs derived from twelve females distributed in three replicates were artificially activated and cultured for seven days. The COCs were artificially activated with ionomycin and four secondary activator combinations: 1.9 mM 6D (6D group), 1.9 mM 6D and 7.5 μ g/mL CB (6D + CB group), 10 μ g/mL CHX (CHX group), and 10 μ g/mL CHX and 7.5 μ g/mL CB (CHX + CB group). On day 3 of IVD, the total cleavage rate and number of embryos containing 2 cells, 3–7 cells, or 8 or more cells were quantified. On day 7 of IVD, the total blastocyst formation rate and the developmental phase [initial blastocyst (iB), blastocyst (B), expanded blastocyst (eB), and blastocyst hatching/hatched (hB)] were quantified. To evaluate the quality, the total numbers of cells were counted by differentiating the ICM and TE.

2.11. Statistical analysis

All data are expressed as the mean $\pm$ standard error and were analyzed using StatView 5.0 (SAS Institute Inc., Cary, NC, USA). The normality of the data was verified using the Shapiro-Wilk test, while the homoscedasticity was verified with Levene's test. The

morphometric oocyte data were examined by variance analysis (ANOVA) and Tukey's test. The levels of ROS and $\Delta\Psi_m$, as well as the number of blastomers were transformed with arcsine and analyzed by ANOVA followed by Tukey's test. All other data were compared with Kruskal-Wallis test and Dunn (multiple comparisons) test. The difference was considered statistically significant when the P value was less than 0.05.

3. Results

In total, 630 visible follicles (31.5 follicles per female on average) were obtained upon stimulation of 40 ovaries (Fig. 1A) derived from 20 collared peccaries with PG600®. The number of medium follicles (385 ± 14.1) was higher than that of small follicles (113 ± 1.9 , $P = 0.03$) and large follicles (132 ± 5.4 , $P = 0.04$). The number of small and large follicles was similar ($P > 0.05$).




Fig. 1. Representative images of (A) ovaries of collared peccary obtained after hormonal induction with PG 600®; (B) *Cumulus*-oocyte complex recovered with homogeneous cytoplasm and more one layer of *cumulus* cells; (C) COCs matured after 44 h IVM; (D) Denuded oocyte with presence of 1PB; (E) Visualization of the metaphasic plate (MII).

Moreover, all ovaries were used to acquire 333 ± 10.7 immature oocytes (Fig. 1B) resulting in a recovery rate of $52.9\% \pm 4.2$ ($333/630$) and 16.2 ± 2.8 oocytes per female. Of these, 253 ± 8.5 (76.0%) structures were classified as GI and GII (12.1 ± 2.2 COCs per female), 37.0 ± 3.3 (11.1%) structures as GIII (3.1 ± 0.8 COCs per female), and 43.0 ± 1.8 (12.9%) structures as GIV (2.8 ± 1.0 COCs per female), according to the cytoplasm homogeneity and the number of *cumulus* cell layers.

Moreover, 333 ± 10.7 immature oocytes (Fig. 1B) were obtained from all ovaries with a recovery rate of $52.9\% \pm 4.2$ (333/630) and 16.2 ± 2.8 oocytes per female. Of these, 253 ± 8.5 (76.0%) oocytes were classified as GI and GII (12.1 ± 2.2 COCs per female), 37.0 ± 3.3 (11.1%) as GIII (3.1 ± 0.8 COCs per female), and 43.0 ± 1.8 (12.9%) as GIV (2.8 ± 1.0 COCs per female) based on the cytoplasm homogeneity and the number of *cumulus* cell layers.

3.1. Experiment 1: effects of EGF supplementation on IVM of collared peccary COCs

The effect of EGF on the IVM of collared peccary COCs is shown in Table 1 and 2. Initially, there was no difference in the IVM rates of COCs exhibiting *cumulus* cell expansion (Fig. 1C) between EGF positive and negative groups. All COCs exhibited *cumulus* cell expansion scores of 3 and 4 and the *cumulus* cell viability was greater than 76.0% for both EGF positive and negative groups. Moreover, there was no difference in the IVM rates of COCs selected on the basis of 1PB presence (Fig. 1D), MII assessments (Fig. 1E), and cytoplasmic maturation evaluated based on mitochondrial distribution. Additionally, the intracellular ROS levels (Fig. 2A, C, and D) and $\Delta\Psi_m$ in the matured COCs were similar between the EGF positive and EGF negative groups.

There was no difference for most of the evaluated oocyte morphometric assessment parameters. However, the thickness of the zona pellucida in the matured oocytes reduced in the presence of EGF (Table 2). Thus, the IVM medium supplemented with EGF was used for experiment 2.

1 **Table 1.** Effects of EGF supplementation on *in vitro* maturation of collared peccary *cumulus*-oocyte complex (COC).

Group	Evaluation of <i>cumulus</i> cells				Nuclear maturation		Cytoplasmic maturation mitochondrial distribution		
	Expansion and grade of expansion, %			Viability, %	1PB, %	MII, %	Peripheral, %	Transition, %	Dispersed, %
	Total	Grade 3	Grade 4						
With EGF	100.0 ± 0.0 (44/44)	63.6 ± 3.0 (28/44)	29.5 ± 0.3 (13/44)	76.0 ± 3.1 (784/1031)	70.5 ± 1.8 (31/44)	68.4 ± 4.9 (13/19)	0.0 ± 0.0 (0/9)	25.0 ± 0.0 (2/8)	75.0 ± 0.7 (6/8)
Without EGF	97.6 ± 1.2 (40/41)	70.7 ± 2.5 (29/41)	29.3 ± 2.1 (12/41)	79.4 ± 1.4 (765/963)	65.9 ± 1.2 (27/41)	62.5 ± 11.6 (10/16)	0.0 ± 0.0 (0/9)	0.0 ± 0.0 (0/9)	100.0 ± 0.0 (9/9)

2 No differences were observed among groups ($P > 0.05$). 1PB, first polar body, MII, metaphase II.

3

1 **Table 2.** Morphometric assessment (in μm and μm^2) in of matured COCs derived from collared peccaries (mean $\pm$ standard error).

Groups	Outer oocyte diameter (ZPO)	Zona pellucida thickness (ZPT)	Ooplasm diameter (OD)	Perivitelline space (PVS) area	Perivitelline space (PVS) diameter
With EGF	373.9 ± 4.4 (37)	33.3 ± 1.2 (37) ^a	284.3 ± 4.2 (37)	7312.6 ± 823.1 (35)	16.2 ± 2.1 (35)
Without EGF	382.2 ± 5.3 (34)	38.6 ± 1.5 (34) ^b	285.6 ± 3.0 (34)	6284.0 ± 390.9 (27)	13.8 ± 0.8 (27)

2 ^{a,b}: Values with different superscript letters within columns are significantly different ($P < 0.05$).

3




Fig. 2. Oocytes of collared peccaries after maturation evaluated as intracellular concentration of reactive oxygen species (ROS) levels and mitochondrial membrane potential ($\Delta\Psi_m$). **(A)** Measurement of fluorescence of ROS levels in matured oocytes in the presence of EGF (EGF positive group) or in the absence (EGF negative group) ($P > 0.05$). **(B)** Quantification of mitochondrial membrane potential in oocytes matured in the presence of EGF (EGF positive group) or in the absence (EGF negative group) ($P > 0.05$). **(C)** Oocytes labelled with fluorescent probe H₂DCFDA for ROS quantification with EGF (EGF positive group) and **(D)** without EGF (EGF negative group). **(E)** Oocytes labelled with fluorescent probe MitoTracker Red[®] for $\Delta\Psi_m$ with EGF (EGF positive group) and **(F)** without EGF (EGF negative group).

3.2. Experiment 2: effects of secondary activators and COC quality on the development of collared peccary parthenogenetic embryos

Assessment of embryonic development is described in Table 3 and 4. The cleavage rate (Fig. 3A–C) after 3 days of IVD was similar in the evaluated groups. The cleavage rates were greater than 72.2% and 25.0% for the embryos derived from GI/GII and GIII COCs, respectively. Moreover, there was no reduction in the cleavage rates of embryos derived from GI/GII and GIII COCs among the 6D and CHX + CB groups (Table 3).

Table 3. Embryonic development of collared peccary GI/GII and GIII *cumulus*-oocyte complex activated with ionomycin in combination with different secondary activators.

Groups	No. COCs		Cleavage, %		Blastocyst/Oocyte, %		Blastocyst /Cleaved, %	
	GI/GII	GIII	GI/GII	GIII	GI/GII	GIII	GI/GII	GIII
			COCs	COCs	COCs	COCs	COCs	COCs
6D	34	4	82.9 ± 2.4 (29)	75.0 ± 0.4 (3)	23.5 ± 0.3 (8) ^a	25.0 ± 0.2 (1) ^a	27.6 ± 0.3 (8) ^a	33.3 ± 0.3 ^a (1)
6D + CB	36	9	72.2 ± 1.4 (26)	77.8 ± 0.7 (7)	16.7 ± 0.5 (6) ^{a,b}	0.0 ± 0.0 (0) ^b	23.1 ± 0.5 (6) ^{a,b}	0.0 ± 0.0 (0) ^b
CHX	37	4	78.4 ± 2.2 (29) ^A	25.0 ± 0.2 (1) ^B	5.4 ± 0.2 (2) ^b	0.0 ± 0.0 (0) ^b	6.9 ± 0.3 (2) ^b	0.0 ± 0.0 (0) ^b
CHX + CB	34	7	76.5 ± 2.2 (26)	57.1 ± 0.4 (4)	17.6 ± 0.3 (6) ^{a,b}	0.0 ± 0.0 (0) ^b	23.1 ± 0.2 (6) ^{a,b}	0.0 ± 0.0 (0) ^b

^{a,b}: Values with different superscript letters within columns are significantly different ($P < 0.05$). ^{A,B}: Values with different superscript letters within rows are significantly different ($P < 0.05$). 6D group: 6-DMAP; 6D + CHX group: 6-DMAP and cytochalasin B; CHX group: cycloheximide; CHX + CB group: cycloheximide and cytochalasin B.




Fig. 3. Kinetics of embryonic development in collared peccaries. After 3 days of IVD different embryos with (A) 2 cells; (B) 4 cells, (C) more than 8 cells. Evaluation of embryonic development on day 7 of IVD (D) blastocyst; (E) blastocyst hatching; (F) blastocyst hatched already without zona pellucida. Counting the total number of blastocyst (G) cells inner mass labelled with Hoechst 33342. (H) trophoctoderm cells stained with propidium iodide. (I) merge total cells.

Nevertheless, the 6D group exhibited higher rates of blastocyst/total cell in the embryos derived from GI/GII COCs than CHX group (Table 3). Moreover, only 6D treatment resulted

1 in the production of embryos derived from GIII COCs ($25.0\% \pm 0.2$). Further, the percentages
2 of ICM/total cell ratio in blastocysts derived from 6D, 6D + CB, and CHX + CB group COCs
3 were higher than those in blastocysts derived from CHX group COCs. Interestingly, the total
4 number of blastocyst cells (Fig. 3G–H) derived from 6D + CB and CHX + CB oocyte groups
5 was similar (Table 4).


6

1 **Table 4.** Cell count of *cumulus-oocyte complexes* (COCs) as viable and activated with ionomycin in combination with different secondary
 2 activators.

Secondary activators	No. of embryos	ICM, %	TE, %	Total	%ICM/Total
6D	5	78.8 ± 35.2	93.2 ± 41.7	172 ± 76.9 ^{a,b}	42.5 ± 19.0 ^a
6D + CB	3	33.7 ± 19.4	65.0 ± 37.5	98.7 ± 57.0 ^a	37.9 ± 21.9 ^a
CHX	2	7.0 ± 0.1	188.0 ± 0.1	195 ± 0.1 ^b	3.6 ± 0.1 ^b
CHX + CB	5	40.2 ± 18.0	57.6 ± 25.8	97.8 ± 43.7 ^a	43.8 ± 19.6 ^a

3 ^{a,b}: Values with different superscript letters within columns are significantly different ($P < 0.05$). ICM: internal cellular mass; TE: trophectoderm;
 4 %ICM/Total: internal cellular mass/ total cell ratio.

1 In relation to embryo development kinetics (Fig. 3D–F), or the proportion of embryos at
2 different stages of development, there was no significant difference among the groups in the
3 percentage of embryos classified with 2, 3–7, or ≥ 8 cells (Fig. 4A–B). On D7, the
4 percentages of iB, B, eB, or hB embryos were similar among the treatment groups (Fig. 4C, P
5 > 0.05).



6
7 **Fig. 4.** Embryonic development of classified activated with ionomycin in combination with
8 different secondary activators. Embryonic development in 3 days with 2 cells, 3–7 cells and 8
9 cells after activation with ionomycin in combination with secondary activators in (A) GI/GII

(P > 0.05) and **(B)** GIII COCs (P > 0.05). **(C)** Percentage of embryos in D7 classified in different stages of blastocyst after activation of GI/GII COCs (P > 0.05).

4. Discussion

Establishment of SCNT is an important strategy to develop this technology as a conservation tool for collared peccary. In this study, we optimized the protocols for improving matured oocyte quality and artificial oocyte activation, which are important for the success of SCNT. Recently [34, 35, 36, 37], our group has developed strategies for the establishment of donor nuclei (or karyoplast), which form a database of somatic resources for the species. We have established the conditions for obtaining cytoplasts and evaluated their competence after artificial oocyte activation using different protocols.

Initially, we evaluated the effects of EGF on the IVM of collared peccary oocytes. We, for the first time, demonstrated that the zona pellucida thickness in oocytes matured with EGF was lower than that in oocytes matured without EGF. The thickness of the zona pellucida is an indicator of the success of hatching and implantation of embryos [38]. The elasticity and thinning of zona pellucida are essential for the hatching process. Thick zona pellucida is associated with low-quality embryos. Increased thickness of zona pellucida may be caused by an inappropriate *in vitro* environment [38]. Zhou et al. [39] demonstrated that thinning solutions enhance the nuclear maturation of oocytes that previously had thick zona pellucida. The study hypothesized that thinner zona pellucida could facilitate first polar body extrusion. Khanmohammadi et al. [38] observed that a suitable culture medium promotes optimal environment that allows zona pellucida thinning and subsequently facilitates embryonic implantation. Furthermore, morphometric evaluation of matured oocytes has been used to estimate the oocyte developmental competence [32].

The expansion of *cumulus* cells was observed in almost all oocytes derived from both EGF positive and negative groups (> 97.8%). A similar response was also observed in swine oocytes [8]. Further, both groups exhibited high rates of *cumulus* cell expansion after maturation. There was no difference in the viability of *cumulus* cells, with values higher than 76% for all groups. This response may have occurred due to the presence of gonadotrophins and/or serum in the IVM medium, which can interfere with the stimulation induced by EGF [40]. Additionally, FSH may act through EGF receptors to exert a synergistic effect on

cytoplasmic maturation [11] and may interfere with the effects of EGF on *cumulus* cell expansion.

The quality of matured oocytes is important for the evaluation of meiotic competence. Thus, we evaluated the ROS levels and $\Delta\Psi_m$ in matured oocytes and observed no difference between the groups for both parameters. Generally, the decrease in ROS levels is accompanied with an increase in glutathione (GSH) levels, a natural antioxidant that reduces ROS in oocytes [41], where ROS generation depends on $\Delta\Psi_m$ [31]. Fan et al. [41] observed that treating the goat oocytes with EGF and cysteamine resulted in lower ROS levels in matured oocytes. In our study, both groups were cultured in a medium containing cysteamine, and EGF probably did not enhance its activity.

Similarly, EGF did not increase the IVM rates of oocytes at MII phase and have 1PB. Although Kishida et al. [42] observed a higher rate of nuclear maturation in porcine oocytes when EGF was present in the medium (10 ng/mL), another study demonstrated that EGF does not influence the nuclear maturation [43]. This may be due to the interaction of EGF with different supplements of the IVM medium, which are highly variable in each study. Further, EGF was involved primarily in the maintenance of embryonic quality where EGF-activated pathways promote greater competence of the oocyte to support embryonic development. Therefore, we suggest the use of EGF in the IVM medium of collared peccary oocytes for future experiments.

In this study, we observed that 6D was the most optimal secondary activator for collared peccary oocytes. However, the combination of 6D and CB did not have a beneficial effect on embryonic development. Further, CHX did not exhibit good potential for oocyte activation, which was improved only in combination with CB. Several studies have demonstrated the ability of 6D, CHX, and CB to inhibit meiotic resumption and second polar body extrusion in porcine oocytes [44, 45]. Moreover, the higher efficiency of 6D compared to CHX was also observed in porcine oocytes [46]. Although 6D and CHX exerted positive effect on oocyte activation and morula-blastocyst formation rates, 6D was more effective than CHX on both matured and immature oocytes [46]. This may be due to the mechanism of action of CHX, which may result in the inhibition of proteins involved in embryonic development [47], whereas 6D specifically inhibits protein kinases [48]. Additionally, Khadijah et al. [49] and

1 Zhang et al. [50] demonstrated that the combination of ionomycin and 6D was better than the
2 combination of ionomycin and CHX for embryonic development in cattle (7.1% vs. 2.2%)
3 and rabbit oocytes (8.6% vs. 1.2%).

4
5 Moreover, the embryonic development rates in the presence of ionomycin and 6D for
6 collared peccary oocytes were higher than those observed for swine oocytes [51] evaluated
7 based on the cleavage percentage (82.9% vs. 47.2%), blastocyst percentage (23.5% vs.
8 11.3%), and ICM/total cell ratio (42.5% vs. 24.8%). This indicates that despite similarities in
9 the responses of protocols used between species, some differences are observed, suggesting
10 the importance of establishing a species-specific protocol. Additionally, Campos Junior et al.
11 [20] activated the collared peccary oocyte with the combination of ionomycin and 6D and
12 obtained up to two-cell (40%) and 4-cell (10%) embryos. Thus, this is the first study that
13 achieved the blastocyst stage of embryonic development in collared peccaries through
14 artificial oocyte activation.

15
16 The low performance of CHX was observed in both artificial oocyte activation of collared
17 peccaries and blastocyst quality. Although our results demonstrated a higher total number of
18 cells in the 6D and CHX groups, CHX group exhibited a higher number of trophectoderm
19 cells and a very low number of cells of the internal mass, which would later impair the
20 embryonic development of the CHX group that had a low blastocyst rate compared to the
21 other groups. The CHX + CB group exhibited higher ICM/total cell rates compared to the
22 CHX group. This may be because CHX has no effect on blocking the extrusion of the second
23 polar body. Thus, parthenotes treated with CHX were largely haploid and exhibited low
24 blastocyst development, which does not improve blastocyst formation further [52]. The
25 combination of 6D and CB was more effective in inhibiting the extrusion of the second
26 polar body, which promotes the development of diploid embryos with higher developmental
27 competence compared to haploid embryos [53].

28
29 The combination of ionomycin and 6D exhibited optimal performance as this was the only
30 combination that produced embryos from GIII oocytes. As CB enhances the effect of CHX on
31 GI and GII oocytes, its mechanism cannot be as efficient as 6D because the combination of
32 CHX and CB could not promote the development of GIII oocytes to blastocysts. However,
33 matured GIII oocytes were activated and developed into embryonic blastocyst only in the 6D

group. Therefore, embryonic development depends on the efficiency of the activation protocol [53]. Thus, use of oocytes with heterogeneous cytoplasm and less than one layer of *cumulus* cells may be suitable when there is low availability of oocytes.

In conclusion, EGF can be used to supplement the maturation medium to obtain a greater quality of matured oocytes and to improve embryonic development. Moreover, the combination of ionomycin and 6D could produce collared peccary embryos from the activation of both GI/GII and GIII COCs. These optimized IVM conditions and artificial activation in collared peccaries represent the first steps for cloning this species through ARTs.

Acknowledgments

The authors thank the CEMAS/UFERSA for providing biological material for the experiments carried out in this research and MSD Animal Health Brazil for the donation of hormones. This study was supported by Brazilian Council of Scientific Development – CNPq and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES, Financial Code 001). AR Silva, MF Oliveira and AF Pereira are recipients of CNPq grants.

References

- [1] Desbiez ALJ, Keuroghlian A, Beisiegel BM, Medici EP, Gatti A, Pontes ARM, et al. Avaliação do risco de extinção do cateto *Pecari tajacu* Linnaeus, 1758, no Brasil. Biodiversidade Brasileira 2012: 74–83.
- [2] Gongora J, Reyna-Hurtado R, Beck H, Taber A, Altrichter M, Keuroghlian A. *Pecari tajacu*. The IUCN Red List of Threatened Species 2011: e.T41777A10562361.
- [3] Albuquerque N, Dias H, Guimarães DDA, Pendu YI, Garcia A, Kahwage P, et al. Criação de caititus em cativeiro: sistema intensivo de produção na Amazônia Oriental. Embrapa Amazônia Oriental-Livro Técnico (INFOTECA-E) n.d. 2016.
- [4] Wani NA, Vettical BS, Hong SB. First cloned Bactrian camel (*Camelus bactrianus*) calf produced by interspecies somatic cell nuclear transfer: A step towards preserving the critically endangered wild Bactrian camels. PloS One 2017; 12:e0177800.
- [5] Borges AA, Pereira AF. Potential role of intraspecific and interspecific cloning in the conservation of wild mammals. Zygote 2019; 27:111–7.
- [6] Honda A, Choijookhuu N, Izu H, Kawano Y, Inokuchi M, Honsho K, et al. Flexible adaptation of male germ cells from female iPSCs of endangered *Tokudaia osimensis*. Sci Adv

- 1 2017; 3:e1602179.
- 2 [7] Borges AA, Santos, MVO, Queiroz Neta LB, Oliveira MF, Silva AR, Pereira AF. *In vitro*
3 maturation of collared peccary (*Pecari tajacu* Linnaeus, 1758) oocytes after different
4 incubation times. Pesq Vet Bras 2018; 38: 1863–8.
- 5 [8] Abeydeera LR, Wang W, Cantley TC, Rieke A, Prather RS, Day BN. Presence of
6 epidermal growth factor during *in vitro* maturation of pig oocytes and embryo culture can
7 modulate blastocyst development after *in vitro* fertilization. Mol Reprod Dev 1998; 51: 395–
8 401.
- 9 [9] Guler A, Poulin N, Mermillod P, Terqui M, Cognie Y. Effect of growth factors, EGF and
10 IGF-I, and estradiol on *in vitro* maturation of sheep oocytes. Theriogenology 2000; 54: 209–
11 18.
- 12 [10] Song H, Kang E, Maeng G, Ock S, Lee S, Yoo J, et al. Influence of epidermal growth
13 factor supplementation during *in vitro* maturation on nuclear status and gene expression of
14 canine oocytes. Res Vet Sci 2011; 91: 439–45.
- 15 [11] Uhm S, Gupta M, Yang J, Chung H-J, Min T, Lee H. Epidermal growth factor can be used
16 in lieu of follicle-stimulating hormone for nuclear maturation of porcine oocytes *in vitro*.
17 Theriogenology 2010; 73:1024–36.
- 18 [12] Orliac MJ, Pierre-Olivier A, Ducrocq S. Phylogenetic relationships of the Suidae
19 (Mammalia, Cetartiodactyla): new insights on the relationships within Suoidea. Zool Scr
20 2010; 39: 315–30.
- 21 [13] Hellgren EC, Syntzke DR, Oldenberg PW, Guthery FS. Demography of a collared peccary
22 population in South Texas. J Wild Management 1995, 59: 153–63.
- 23 [14] Gottdenker N, Bodmer RE. Reproduction and productivity of white-lipped and collared
24 peccaries in the Peruvian Amazon. J Zool 1998; 245: 423–30.
- 25 [15] Souza ALP, Lima GL, Peixoto GCX, Castelo TS, Oliveira MGC, Paula VV, et al. Sperm
26 characteristics following freezing in extenders supplemented with whole egg yolk and
27 different concentrations of low-density lipoproteins in the collared peccary (*Pecari tajacu*).
28 Reprod Biol 2015;15:223e8.
- 29 [16] Hu J, Li OW, Li G, Chen XY, Yang H, Zhang SS, et al. The cryoprotective effect on
30 frozen-thawed boar semen of egg yolk low density lipoproteins. Asian Aust J Anim Sci
31 2006;19:486–94.
- 32 [17] Fernandes CB, Devito LG, Martins LR, Blanco IDP, Lima Neto JF, Tsuribe PM, et al.
33 Artificial activation of bovine and equine oocytes with cycloheximide, roscovitine, strontium,

- 1 or 6-dimethylaminopurine in low or high calcium concentrations. *Zygote* 2014; 22: 387–94.
- 2 [18] Pathak J, Kharche S, Goel A. Effects of different activation protocols on cleavage rate and
3 blastocyst production of caprine oocytes. *Iran J Vet Res* 2017; 18: 243.
- 4 [19] Macedo MP, Glanzner WG, Rissi VB, Gutierrez K, Currin L, Baldassarre H, et al. A fast
5 and reliable protocol for activation of porcine oocytes. *Theriogenology* 2019; 123: 22–9.
- 6 [20] Campos Junior PHA, Costa GMJ, Avelar GF, Lacerda SMSN, Costa NN, Ohashi OM, et
7 al. Derivation of sperm from xenografted testis cells and tissues of the peccary (*Tayassu*
8 *tajacu*). *Reproduction* 2014; 147:291–9.
- 9 [21] Cervera R, Silvestre M, Martí N, García-Mengual E, Moreno R, Stojkovic M. Effects of
10 different oocyte activation procedures on development and gene expression of porcine pre-
11 implantation embryos. *Reprod Domest Anim* 2010; 45: 12–20.
- 12 [22] Ock S, Kwack D, Mohana Kumar B, Han J, Kim S, Rho G. Effects of activation methods
13 on DNA synthesis and development of parthenogenetic porcine embryos. *Reprod Domest*
14 *Anim* 2011; 46:1082–9.
- 15 [23] Collas P, Robl JM. Factors affecting the efficiency of nuclear transplantation in the rabbit
16 embryo. *Biol Reprod* 1990; 43:877–884.
- 17 [24] Qiu X, Li N, Xiao X, Li Y. Aggregation of a parthenogenetic diploid embryo and a male
18 embryo improves the blastocyst development and parthenogenetic embryonic stem cell
19 derivation. *Turk J Biol* 2017; 41: 629–639.
- 20 [25] Assidi M, Dufort I, Ali A, Hamel M, Algriany O, Dielemann S, et al. Identification of
21 potential markers of oocyte competence expressed in bovine *cumulus* cells matured with
22 follicle-stimulating hormone and/or phorbol myristate acetate *in vitro*. *Biol Reprod* 2008; 79:
23 209–22.
- 24 [26] Almeida KC, Pereira AF, Alcantara Neto AS, Avelar SRG, Bertolini LR, Bertolini M, et
25 al. Real-time qRT-PCR analysis of EGF receptor in *cumulus*-oocyte complexes recovered by
26 laparoscopy in hormonally treated goats. *Zygote* 2011; 19:127–36.
- 27 [27] Peixoto GCX, Lima GL, Maia KM, Souza ALP, Castelo TS, Paiva ALC, Paula VV, et al.
28 Single injection of eCG/hCG leads to successful estrous synchronization in the collared
29 peccary (*Pecari tajacu* Linnaeus, 1758). *Anim Reprod Sci* 2019; 298: 106112.
- 30 [28] Viana ACS, Freitas JLS, Campos LB, Silva AR, Oliveira MF, Freitas VJF, et al. Analysis
31 of ovarian response to superstimulation treatment in collared peccary (*Pecari tajacu*
32 Linnaeus, 1758). *Anim. Reprod* 2017; 4:804.
- 33 [29] Kumar S, Devi B, Purkayastha A, Bharti PK, Doley S, Kadirvel G. Porcine ovarian

- 1 biometry, oocyte retrieval and quality of oocytes under different seasons in north east hill
2 region. Indian J Anim Sci 2016; 86: 300–3.
- 3 [30] Kubo N, Cayo-Colca IS, Miyano T. Effect of estradiol-17 β during *in vitro* growth culture
4 on the growth, maturation, *cumulus* expansion and development of porcine oocytes from early
5 antral follicles. Anim Sci J 2015; 86: 251–9.
- 6 [31] Santos MVO, Nascimento LE, Praxedes ÉA, Borges AA, Silva AR, Bertini LM, et al.
7 *Syzygium aromaticum* essential oil supplementation during *in vitro* bovine oocyte maturation
8 improves parthenogenetic embryonic development. Theriogenology 2019; 128:74–80.
- 9 [32] Saadeldin IM, Swelum AAA, Yaqoob SH, Alowaimer AN. Morphometric assessment of
10 *in vitro* matured dromedary camel oocytes determines the developmental competence after
11 parthenogenetic activation. Theriogenology 2017; 95:141–8.
- 12 [33] Paul AK, Liang Y, Srirattana K, Nagai T, Parnpai R. Vitrification of bovine matured
13 oocytes and blastocysts in a paper container. Anim Sci J 2018; 89: 307–15.
- 14 [34] Borges AA, Lima GL, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, et al.
15 Conservation of somatic tissue derived from collared peccaries (*Pecari tajacu* Linnaeus,
16 1758) using direct or solid-surface vitrification techniques. Cytotechnology 2017; 69: 643–54.
- 17 [35] Borges AA, Lira GPO, Nascimento LE, Queiroz Neta LB, Santos MVO, Oliveira MF, et
18 al. Influence of cryopreservation solution on the *in vitro* culture of skin tissues derived from
19 collared peccary (*Pecari tajacu* Linnaeus, 1758). Biopreserv Biobank 2018; 16: 77–81.
- 20 [36] Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, Pereira AF.
21 Combination of ethylene glycol with sucrose increases survival rate after vitrification of
22 somatic tissue of collared peccaries (*Pecari tajacu* Linnaeus, 1758). Pesq Vet Bras 2018; 38:
23 77–81.
- 24 [37] Santos MLT, Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, et al.
25 *In vitro* culture of somatic cells derived from ear tissue of collared peccary (*Pecari tajacu*
26 Linnaeus, 1758) in medium with different requirements. Pesq Vet Bras 2016; 36: 1194–202.
- 27 [38] Khanmohammadi N, Movahedin M, Safari M, Sameni HR, Yousefi B, Jafari B, et al.
28 Effect of L-carnitine on *in vitro* developmental rate, the zona pellucida and hatching of
29 blastocysts and their cell numbers in mouse embryos. Int J Reprod Biomed 2016; 14:649.
- 30 [39] Zhou HX, Ma YZ, Liu YL, Chen Y, Zhou CJ. Assessment of mouse germinal vesicle
31 stage oocyte quality by evaluating the *cumulus* layer, zona pellucida, and perivitelline space.
32 Plos One 2014; 9: e105812.
- 33 [40] Lorenzo P, Illera M, Illera J, Illera M. Enhancement of *cumulus* expansion and nuclear

- 1 maturation during bovine oocyte maturation *in vitro* by the addition of epidermal growth
- 2 factor and insulin-like growth factor I. *Reproduction* 1994; 101:697–701.
- 3 [41] Fan Z, Yang M, Regouski M, Polejaeva IA. Effects of three different media on *in vitro*
- 4 maturation and development, intracellular glutathione and reactive oxygen species levels, and
- 5 maternal gene expression of abattoir-derived goat oocytes. *Small Rumin Res* 2017; 147: 106–
- 6 14.
- 7 [42] Kishida R, Lee E, Fukui Y. *In vitro* maturation of porcine oocytes using a defined medium
- 8 and developmental capacity after intracytoplasmic sperm injection. *Theriogenology* 2004; 62:
- 9 1663–76.
- 10 [43] Abeydeera L, Wang W, Cantley T, Rieke A, Murphy C, Prather R, et al. Development and
- 11 viability of pig oocytes matured in a protein-free medium containing epidermal growth factor.
- 12 *Theriogenology* 2000; 54: 787–97.
- 13 [44] Beux G, Richard FJ, Sirard MA. Effect of cycloheximide, 6-DMAP, roscovitine and
- 14 butyrolactone I on resumption of meiosis in porcine oocytes. *Theriogenology* 2003; 60: 1049–
- 15 58.
- 16 [45] Yi Y, Park C. Parthenogenetic development of porcine oocytes treated by ethanol,
- 17 cycloheximide, cytochalasin B and 6-dimethylaminopurine. *Anim Reprod Sci* 2005; 86: 297–
- 18 304.
- 19 [46] Demir K, Can A, Ertürk E, Özdemirci S, Karacam H, Sandal AI, et al. Effect of different
- 20 activation techniques on immature and *in vitro* matured cat oocytes. *Kafkas Univ Vet Fak*
- 21 *Derg* 2014; 20: 565–70.
- 22 [47] Che L, Lalonde A, Bordignon V. Chemical activation of parthenogenetic and nuclear
- 23 transfer porcine oocytes using ionomycin and strontium chloride. *Theriogenology* 2007; 67:
- 24 1297–304.
- 25 [48] Alexander B, Coppola G, Berardino D, Rho G, St John E, Betts D, et al. The effect of 6-
- 26 dimethylaminopurine (6-DMAP) and cycloheximide (CHX) on the development and
- 27 chromosomal complement of sheep parthenogenetic and nuclear transfer embryos. *Mol*
- 28 *Reprod Dev* 2006; 73: 20–30.
- 29 [49] Khadijah S, Rahman M, Abdullah R, Wan K. Comparison of bovine embryonic
- 30 development of parthenogenetically activated oocytes by different combinations of chemical
- 31 treatments. *Mal J Anim Sci* 2014; 17: 85–94.
- 32 [50] Zhang S, Xiang S, Yang J, Shi J, Guan X, Jiang J, et al. Optimization of parthenogenetic
- 33 activation of rabbit oocytes and development of rabbit embryo by somatic cell nuclear

- 1 transfer. *Reprod Dom Anim* 2019; 54: 258–69.
- 2 [51] García-Mengual E, Alfonso J, Salvador I, Duque C, Silvestre M. Oocyte activation
3 procedures and influence of serum on porcine oocyte maturation and subsequent
4 parthenogenetic and nuclear transfer embryo development. *Zygote* 2008; 16:279–84.
- 5 [52] Lee J, Tian XC, Yang X. Optimization of parthenogenetic activation protocol in porcine.
6 *Mol Reprod Dev* 2004; 68: 51–7.
- 7 [53] Kim Y, Lee S, Ock S, Balasubramanian S, Choe S, Rho G. Development of cloned pig
8 embryos by nuclear transfer following different activation treatments. *Mol Reprod Dev* 2005;
9 70: 308–13.
- 10 [54] Felmer R, Arias M. Activation treatment of recipient oocytes affects the subsequent
11 development and ploidy of bovine parthenogenetic and somatic cell nuclear transfer (SCNT)
12 embryos. *Mol Reprod Dev* 2015; 82: 441–9.

6 CONSIDERAÇÕES FINAIS E PERSPECTIVAS

A presente tese obteve resultados significativos no desenvolvimento da clonagem por transferência nuclear de célula somática (TNCS) em catetos, uma vez que nós estabelecemos a obtenção de células doadoras de núcleo ou carioplastos e células doadoras de citoplasma ou citoplastos. No que se refere aos estudos dos carioplastos, inicialmente, nós descrevemos o primeiro trabalho de estabelecimento e caracterização de fibroblastos de catetos adultos durante o cultivo *in vitro* e criopreservação. Posteriormente, nós estabelecemos que a combinação de 10% de dimetilsulfóxido associado com 0,2 M de sacarose e 50% de soro fetal bovino foi a mais eficiente solução de criopreservação para a formação de bancos de células somáticas de catetos. Adicionalmente, nós definimos como melhor protocolo de sincronização do ciclo dessas células em G₀/G₁ o método de inibição por contato por três dias. Portanto, nós estabelecemos a etapa de carioplastos da TNCS de catetos, obtendo células de qualidade e aptas a serem usadas como doadoras de núcleo.

Já no que se refere aos estudos dos citoplastos, inicialmente, nós otimizamos as condições de maturação *in vitro* de oócitos, observando o tempo de 48 h e a presença de 10 ng/mL de fator de crescimento epidermal (EGF) como adequados para uma maior competência meiótica destas estruturas. Além disso, nós verificamos que a ativação química usando ionomicina e 6-dimetilaminopurina (6-DMAP) foi a mais eficiente combinação, tendo esta tese alcançado como resultado significativo, uma taxa de 27,6% de blastocistos de catetos derivados da ativação oocitária artificial.

Em síntese, nós obtivemos carioplastos e citoplastos que poderão ser empregados na TNCS de catetos, deixando a ponto as etapas fundamentais para a clonagem desta espécie. Ainda, destaca-se que os conhecimentos aqui gerados poderão ser aplicados em estudos para produção de células induzidas à pluripotência; ensaios de toxicidade e farmacológicos; fecundação *in vitro*; injeção intracitoplasmática de espermatozoides; compreensão da biologia reprodutiva e do desenvolvimento embrionário; obtenção de células embrionárias totipotentes; avaliações de criotolerância do oócito e do embrião. Portanto, este trabalho foi um grande passo para a conservação de taiassuídeos, especialmente dos catetos.

ANEXO A – VERSÃO DO ARTIGO PUBLICADO NA ZYGOTE

**POTENTIAL ROLE OF INTRASPECIFIC AND INTERSPECIFIC CLONING IN
THE CONSERVATION OF WILD MAMMALS**

QUALIS: Qualis 2013-2016: B1/ Novo Qualis: B2

FATOR DE IMPACTO: 1,278

DOI: <https://doi.org/10.1017/S0967199419000170>

Potential role of intraspecific and interspecific cloning in the conservation of wild mammals

Alana Azevedo Borges and Alessandra Fernandes Pereira

Laboratory of Animal Biotechnology, Federal Rural University of Semi-Arid, Mossoró, RN, Brazil

Review

Cite this article: A.A. Borges and A.F. Pereira (2019) Potential role of intraspecific and interspecific cloning in the conservation of wild mammals. *Zygote* page 1 of 7. doi: 10.1017/S0967199419000170

Received: 11 December 2018

Revised: 8 February 2019

Accepted: 25 February 2019

Keywords:

Biodiversity; Biotechnology; Embryo development; SCNT; Somatic cells

Address for correspondence:

Alessandra Fernandes Pereira. Laboratory of Animal Biotechnology, Federal Rural University of Semi-Arid, Av. Francisco Mota, 572, Mossoró, RN, 59625-900, Brazil. Tel: +55 84 3317 8361. E-mail address: alexsandra.pereira@ufersa.edu.br

Summary

Intraspecific and interspecific cloning via somatic cell nuclear transfer (iSCNT) is a biotechnique with great possibilities for wild mammals because it allows the maintenance of biodiversity by recovering species, nuclear reprogramming for the production of pluripotency-induced cells, and studies related to embryonic development. Nevertheless, many areas in cloning, especially those associated with wild mammals, are still in question because of the difficulty in obtaining cytoplasmic donor cells (or cytoplasts). Conversely, donor cell nuclei (or karyoplasts) are widely obtained from the skin of living or post-mortem individuals and often maintained in somatic cell banks. Moreover, the creation of karyoplast–cytoplast complexes by fusion followed by activation and embryo development is one of the most difficult steps that requires further clarification to avoid genetic failures. Although difficult, cloning different species, such as wild carnivores and ungulates, can be successful via iSCNT with embryo development and the birth of offspring. Thus, novel research in the area that contributes to the conservation of biodiversity and knowledge of the physiology of species continues. The present review presents the failures and successes that occurred with the application of the technique in wild mammals, with the goal of helping future work on cloning via iSCNT.

Introduction

The decrease in the biodiversity of wild mammals has been caused mainly by human activity, resulting in an increase in research aimed at the development of conservation strategies (Pereira *et al.*, 2016). In general, several techniques to help with the conservation of wild animals are available, including the formation of biobanks (León-Quinto *et al.*, 2009), artificial insemination (Howard *et al.*, 2016), embryo transfer (Goeritz *et al.*, 2012), *in vitro* fertilization (Herrick *et al.*, 2010), and cloning using somatic cell nuclear transfer (SCNT, Folch *et al.*, 2009). Because of the low availability of oocytes for SCNT, interspecific cloning using intraspecific and interspecific nuclear transfer techniques (iSCNT) has been shown to be an important tool in conservation (Wani *et al.*, 2017).

The main argument for the application of iSCNT is the rapid decrease in the number of species. Any tool that can avoid this decrease is important. iSCNT preserves and even expands genetic variability when somatic cells of different individuals representative of the original biodiversity of a population are collected for its use (Loi *et al.*, 2001). In addition, interest in cloning has increased not only for the conservation of endangered species, but also for the multiplication of reproducers with better genetic characteristics (Saini *et al.*, 2015), basic research on cell epigenetic status (Saragusty *et al.*, 2016), embryonic development (González-Grajales *et al.*, 2016), and the production of induced pluripotent cells (Sukparangsi *et al.*, 2018).

Therefore, in all applications of cloning, studies related to the improvement of iSCNT, as well as its wide use in different individuals, are important.

Overview of the iSCNT technique and its limitations

The iSCNT technique involves embryo reconstruction by fusing a nucleus of a donor cell (karyoplast) derived from a wild mammal with an enucleated oocyte (cytoplast) from a domestic mammal of a different species, family, order, or class (Do & Taylor-Robinson, 2014). The nucleus in G0/G1 is exposed to reprogramming by the oocyte, followed by the fusion and activation of the reconstructed embryo (Loi *et al.*, 2011). Subsequently, the resultant embryo can be transplanted into the uterus of a recipient for term development (Pereira & Freitas, 2009).

Different steps are involved in the production of clones via iSCNT. Therefore, it is interesting to highlight the steps of the technique and its peculiarities that can define the success of cloning by iSCNT.

Preparation of cytoplasts

Whether using the oocyte from a domestic or a wild mammal, some fundamental criteria must be met to obtain a cytoplast suitable for cloning, such as oocyte selection, *in vitro* maturation, and enucleation systems (Loi *et al.*, 2011). In general, follicular size, the oocyte collection method, and the culture environment are factors that can affect the quality of mature oocytes, and different responses to these factors can be observed in wild mammals. In some cervid species, Brahmasani *et al.* (2013) observed that low maturation rates could probably be caused by slicing. In this method, non-competent oocytes can be recovered, as the technique can result in the recovery of structures of small diameter follicles (Rho *et al.*, 2001). Additionally, the quality of ovaries obtained post-mortem may have been one of the factors that reduce the quality of oocytes in these species.

Therefore, studies have shown that enriched culture medium and ovarian transport conditions may result in good results using ovaries from post-mortem animals for recovery of immature oocytes. Macías-García *et al.* (2018) verified that oocytes of Iberian red deer (*Cervus elaphus hispanicus*) obtained from ovaries maintained for 16 h in a holding medium increased the oocyte meiotic competence. Moreover, these authors observed that the epidermal growth factor (EGF) demonstrated a differential effect depending upon oocyte grading and conditions of ovary transportation. Additionally, for ovaries derived from Hokkaido sika deer (*Cervus nippon yesoensis*), maturation rates of oocytes were highest when ovaries were kept for 12 h at 20–25°C, when compared with 24 h (Tulake *et al.*, 2014).

Specifically, with respect to the culture environment, the requirements for both composition and maturation time should be established for the *in vitro* maturation of each species. In the Indian blackbuck (*Antelope cervicapra*), oocytes cultured in the presence of gonadotropins (follicle-stimulating hormone, FSH and luteinizing hormone, LH) showed higher rates of expansion of the cumulus oophorus (79.3%) and extrusion of the first polar body (46.1%) compared with oocytes cultured without gonadotropins (60.4% and 33.3%, respectively) (Rao *et al.*, 2010). In the sika deer (*Cervus nippon hortulorum*), oocytes cultured in medium supplemented with fetal bovine serum (FBS), FSH, LH, cysteamine and EGF resulted in a higher maturation rate (75.4%) compared with medium without supplementation (30.1%; Yin *et al.*, 2013). Already, different hormonal combinations of FSH, LH and 17 β -estradiol did not alter the maturation rates in oocytes derived from lions (*Panthera leo*; Fernandez-Gonzalez *et al.*, 2015). In the collared peccary (*Pecari tajacu*), we proved that oocytes need 48 h to achieve maturation instead of 24 h, according to the expansion of the cumulus cells (100% vs. 38.1%), the presence of first polar body (90.5% vs. 52.4%), and the status of the nucleus in the second metaphase (76.2% vs. 52.4%), respectively (Borges *et al.*, 2018c).

In addition to obtaining mature oocytes, the preparation of cytoplasts depends on the method of enucleation of these structures. The amount of ooplasm present in the reconstructed embryo is related to the enucleation technique that removes the nucleus from the oocyte. Matured oocytes can be enucleated in different ways, including squeezing the first polar body and the surrounding cytoplasm through a cleft in the zona pellucida of the oocyte (Lee *et al.*, 2003). Another method is manual removal in which zona-free oocytes

are enucleated with a bisection blade that hand bisect the metaphase II chromosomes along with a small volume of the surrounding cytoplasm. Oocytes can also be aspirated using a micromanipulator at the location of the metaphase II chromosomes and the polar body via brief exposure to ultraviolet light (Pereira *et al.*, 2015).

Selection of karyoplasts

To obtain karyoplasts appropriate for cloning, their type and age and the manipulation techniques used are important for their future reprogramming (Kim *et al.*, 2007). Karyoplasts can be obtained from fresh or cryopreserved somatic tissues (Folch *et al.*, 2009, Pan *et al.*, 2014), from an adult (Moulavi *et al.*, 2017) or a fetus (Liu *et al.*, 2018), and *in vivo* or post-mortem (Pereira *et al.*, 2014). Although the recovery of these cells is not a difficult task, their processing and preservation until use in iSCNT require attention (Pereira *et al.*, 2014). In general, skin cells have been the most used cell type for karyoplasts (Song *et al.*, 2007). The skin has an abundance of cells of interest that may have different efficiencies in cloning, as observed in wild buffalo (*Bubalus arnee*). Saini *et al.* (2015) detected that fibroblasts of this species are easier to reprogram than epithelial cells.

After harvest, cells used as nuclei donors need to be characterized with respect to their culture conditions, cryopreservation, and cell cycle synchronization (Pereira *et al.*, 2014). For these steps, cells are evaluated for the number of passages, nutritional requirements during *in vitro* culture (Santos *et al.*, 2016), and the damage done during cryopreservation (Song *et al.*, 2007). Thus, karyoplasts have been established *in vitro* in Dulbecco's modified Eagle's medium (DMEM) supplemented with FBS and growth factors (Santos *et al.*, 2016).

Karyoplasts have been routinely cryopreserved by slow freezing (Sharma *et al.*, 2018) using a combination of dimethyl sulfoxide (DMSO), FBS, and sucrose as the cryoprotectant, as observed with Iberian lynx (*Lynx pardinus*, León-Quinto *et al.*, 2014). Although it is more desirable to use a somatic cell bank after tissue culture, the absence of *in vitro* culture conditions sometimes makes these banks unfeasible, resulting in the immediate formation of the targets for those somatic tissues (Borges *et al.*, 2017a,b; Queiroz Neta *et al.*, 2018). The three somatic tissue conservation techniques used for wild animals are slow-freezing cryopreservation (Mestre-Citrinovit *et al.*, 2016), vitrification (Borges *et al.*, 2018a,b), and cooling at 4–6°C (Queiroz Neta *et al.*, 2018). In collared peccaries, we compared two techniques of vitrification and we observed that solid-surface vitrification was found to be a more efficient method for vitrifying skin tissue when compared with direct vitrification in cryovials, probably due to tissues not being involved in large amounts of cryoprotectants before passing through a drastic change in temperature during the solid-surface vitrification (Borges *et al.*, 2017b).

Finally, the third step in the preparation of the karyoplasts is cell synchronization in the G0/G1 stage (Gómez *et al.*, 2003; Yelisetti *et al.*, 2016). In general, nuclear reprogramming is controlled by epigenetic modification. For this to occur, the somatic cells must be in G0/G1 to allow the removal of reversible epigenetic changes acquired during cell differentiation (Song *et al.*, 2007). Therefore, cells can be subjected to different treatments for synchronization during culture. Inhibition by contact (Moulavi *et al.*, 2017), serum deprivation (Wani *et al.*, 2017), and chemicals that inhibit the cell cycle (Gómez *et al.*, 2003) are methods used for synchronization. Serum deprivation and inhibition by contact are the most commonly used (Moulavi *et al.*, 2017).

Under high confluence or serum privation, fibroblast cells derived from the skin of adult argali (*Ovis ammon*) were efficiently

Table 1. iSCNT in some wild mammals

Scientific name	IUCN Category*	Country	Karyoplast	Cytoplast	Outcome	Authors
Carnivores						
<i>Canis lupus</i>	Least Concern	Seoul, Korea	Fibroblast	<i>Canis lupus familiaris</i>	17% pregnancies, 2 pups	Kim <i>et al.</i> (2007)
<i>Canis lupus</i>	Least Concern	Seoul, Korea	Fibroblast	<i>Canis lupus familiaris</i>	23.5% pregnancies, 4 pups	Oh <i>et al.</i> (2008)
<i>Felis margarita</i>	Least Concern	USA	Fibroblast	<i>Felis catus</i>	3 pups	Gómez <i>et al.</i> (2008)
<i>Felis silvestris lybica</i>	Least Concern	USA	Fibroblast	<i>Felis catus</i>	28% blastocyst	Gómez <i>et al.</i> (2003)
<i>Felis silvestris lybica</i>	Least Concern	USA	Fibroblast	<i>Felis catus</i>	75% pregnancies, 17 pups	Gómez <i>et al.</i> (2004)
Ungulates						
<i>Acinonyx jubatus</i>	Vulnerable	Argentina	Fibroblast	<i>Felis catus</i>	27.4% blastocyst	Moro <i>et al.</i> (2015)
<i>Acinonyx jubatus venaticus</i>	Vulnerable	Iran	Fibroblast	<i>Felis catus</i>	5.9% morula	Moulavi <i>et al.</i> (2017)
<i>Bos gaurus</i>	Vulnerable	USA	Fibroblast	<i>Bos taurus</i>	25% pregnancies, none term	Lanza <i>et al.</i> (2000)
<i>Bos javanicus</i>	Endangered	USA	Fibroblast	<i>Bos taurus</i>	17% pregnancies, none term	Sansinena <i>et al.</i> (2005)
<i>Bubalus arnee</i>	Endangered	India	Fibroblast	<i>Bubalus bubalis</i>	38.7% blastocyst	Priya <i>et al.</i> (2014)
<i>Bubalus arnee</i>	Endangered	India	Fibroblast/Epithelial cell	<i>Bubalus bubalis</i>	50.6 vs 20.5% blastocyst	Saini <i>et al.</i> (2015)
<i>Capra ibex</i>	Least Concern	China	Fibroblast	<i>Capra hircus</i>	11% blastocyst	Wang <i>et al.</i> (2007)
<i>Capra pyrenaica</i>	Absente	Spain	Fibroblast	<i>Capra pyrenaica</i>	1 born	Folch <i>et al.</i> (2009)
<i>Ovis ammon</i>	Near Threatened	China	Fibroblast/Cumulus cell	<i>Ovis aries</i>	22.1% blastocyst	Pan <i>et al.</i> (2014)
<i>Ovis orientalis musimon</i>	Vulnerable	Italy	Granulosa cell	<i>Ovis aries</i>	1 pup	Loi <i>et al.</i> (2001)
<i>Tragelaphus eurycerus isaaci</i>	Critically Endangered	USA	Fibroblast	<i>Bos taurus</i>	24% blastocyst	Lee <i>et al.</i> (2003)
Others species						
<i>Balaenoptera bonaerensis</i>	Data deficient	Japan	Cumulus cells	<i>Bos taurus/Sus scrofa domestica</i>	27.5–52.8% cleavage	Ikumi <i>et al.</i> (2004)
<i>Macaca fascicularis</i>	Least Concern	Thailand	Fibroblast	<i>Bos taurus</i>	33% blastocyst	Lorthongpanich <i>et al.</i> (2008)

*IUCN: International Union for Conservation of Nature and Natural Resources. USA: United States of America.

synchronized at G0/G1; nevertheless, cells were in lower proportion in the growing stage (Pan *et al.*, 2014). Authors observed that the highest proportion of cells from the African wild cat (*Felis silvestris lybica*) at G0/G1 was obtained by serum deprivation compared with that obtained by inhibition by contact and the inhibitor roscovitine (Gómez *et al.*, 2003). Leopard (*Panthera pardus*) skin cells treated with chemical inhibitors such as sodium butyrate have a greater propensity to undergo alterations (Yelisetti *et al.*, 2016).

Embryonic reconstruction stages

After the transfer of the nucleus into the enucleated oocyte, the cytoplasm–karyoplast complex is subjected to an electric pulse that not only induces the fusion of the somatic cell nucleus with the enucleated oocyte to form a new complex, but also promotes the release of intracellular calcium that initiates cellular activation

(Pereira & Freitas, 2009). In general, the successful development of a reconstructed embryo depends on the complex interactions between the cytoplasm and the nuclear structure during embryonic development; failures in this interaction can cause problems during early cleavage and embryonic development (González-Grajales *et al.*, 2016).

The activation of the cytoplasm–karyoplast complexes guarantees adequate embryonic reconstruction (Yamochi *et al.*, 2013). Because the iSCNT technique reprograms the nucleus of a somatic cell of one species using the oocyte cytoplasm of another species, it is essential that the activation protocol be able to activate the reconstructed embryo (Zhao *et al.*, 2006). Physiologically, a mammalian oocyte is activated during fusion with a sperm, releasing meiotic cell cycle arrest and enabling the resumption of the oocyte meiotic cell cycle (Sparman *et al.*, 2010). Therefore, a well developed protocol allows a high rate of blastocyst formation by promoting good embryonic development through activation.

Activation protocols, including physical methods such as electrical pulses and alteration of osmolarity, and chemical methods such as calcium-mobilizing compounds like strontium chloride, ionomycin, and ethanol, to promote the initial release of calcium have been evaluated in different species, as sika deer (Yin *et al.*, 2013), alpaca (*Vicugna pacos*) and llama (*Lama glama*, Ruiz *et al.*, 2015), with blastocyst rates of 32.4%, 22.5% and 18.7%, respectively. In general, calcium mobilizers are used in combination with kinase protein inhibitors or protein synthesizers such as cycloheximide and 6-dimethylaminopurine (6-DMAP). In addition, a cytostatic factor inactivator and microfilament inhibitor such as cytochalasin B are used to prevent extrusion of the second polar body and maintain the diploidy of the presumed embryo (Ruiz *et al.*, 2015).

For red deer (*Cervus elaphus*), electrical activation before chemical activation with ionomycin and 6-DMAP was efficient for the production of clone embryos (32–44%), obtaining genetically healthy calves (Berg *et al.*, 2007). Nevertheless, the same protocol resulted in a low developmental rate (5.7%) of activated oocytes in swamp deer and 0.0% embryos in spotted deer, sambar deer, and brow-antlered deer after oocyte parthenogenetic activation (Brahmasani *et al.*, 2013). Blackbuck (*Antelope cervicapra*) oocytes activated with ionomycin and 6-DMAP resulted in 58% cleaved embryos and 13% blastocysts (Rao *et al.*, 2010). Therefore, the artificial activation method (chemical, electrical protocols or your combination) can result in different responses among species. In this sense, it is necessary to evaluate the type of artificial activation that promotes the best rates of embryonic development in the species of interest.

In vitro culture systems are essential for early embryonic development and nuclear reprogramming (Gómez *et al.*, 2008; Pereira *et al.*, 2013). Choosing the appropriate culture medium for each species is considered the initial step in proper embryonic development (Zhao *et al.*, 2006). Lee *et al.* (2003) used somatic cells of the mountain bongo (*Tragelaphus eurycerus isaaci*) and domestic cow (*Bos taurus*) oocytes and observed that a chemically defined, protein-free medium of TCM199 supplemented with FBS supported embryonic development. Nonetheless, there is no one culture medium suitable for all species that allows better embryonic development for a given species under study.

Finally, the effect of epigenetic reprogramming is a very relevant factor in the success of iSCNT (Gómez *et al.*, 2008). Some epigenetic markers were characterized with respect to their function during embryonic reprogramming and their influence on the chromatin structure from post-translational modifications (Song *et al.*, 2007). The overall level of the acetylation of histone H3 at lysine 18 (H3K18ac) and trimethylation of histone H3 at lysine 27 (H3K27me3), and the expression level of some important apoptosis proteins (caspase 3 and caspase 7), and p53 were evaluated. The hyperacetylated state of histones is associated with transcriptionally active domains, while the hypoacetylated state is associated mainly with silenced chromatin regions of histone acetyl transferases and histone deacetylases. The methylation pattern of the DNA is determined by DNA methyltransferases. OCT3/4, NANOG, and CDX2 are very important because of their close association with pluripotency and early embryonic development (Saini *et al.*, 2015).

Advances and perspectives of iSCNT in wild mammals

Several works aimed at cloning different wild mammals have been conducted (Table 1). Among these studies, those that obtained offspring were on wild bovine (Lanza *et al.*, 2000), sheep

(Loi *et al.*, 2001), felid (Gómez *et al.*, 2004; Li *et al.*, 2007), canid (Kim *et al.*, 2007; Oh *et al.*, 2008), and goat (Folch *et al.*, 2009). Therefore, several families have proven the success of using iSCNT for the recovery and reintroduction of wild mammals.

An important point to remember is that as the taxonomic distance between donor and recipient species increases, the production of blastocysts decreases because of the decreased ability of somatic cells to be reprogrammed (Priya *et al.*, 2014). In general, enucleated oocytes are from a domestic species that is phylogenetically close to the wild species that donates the nucleus. For example, domestic sheep cytoplasts were able to reprogram me argali fibroblast nuclei (Pan *et al.*, 2014) and domestic buffalo cytoplast was able to reprogram me wild buffalo karyoplast (Priya *et al.*, 2014).

Carnivores

Some works have shown the advances achieved by iSCNT in wild canine species. These species, including the grey wolf (*Canis lupus*), have gradually become endangered or extinct. Therefore, in 2007, with the goal of canid conservation, Kim *et al.* (2007) cultured fibroblasts derived from the ear of an adult female grey wolf that were then used as donor cells of nuclei. Using domestic canine oocytes, the authors produced a pregnancy with cloned embryos of two genetic identities of the cloned wolves, but there were no births. In 2008, Oh *et al.* (2008) obtained three wolf pups from cloned embryos using cells obtained from a male grey wolf 6 h after death and domestic canine oocytes. These studies demonstrated the successful cloning of endangered wild canines.

In felid species, the main oocyte source has been the domestic cat. In species from the Felidae subfamily, some progress has been achieved. Therefore, synchronized nuclei were donated by the African wild cat and transferred to enucleated domestic cat oocytes resulting in a high rate of blastocyst formation but no pregnancies (Gómez *et al.*, 2003). In another study in which embryos were constructed using somatic cells derived from the African wild cat and domestic cat oocytes, 75% of the embryos developed to term and 25% underwent fetal resorption or abortion (Gómez *et al.*, 2004). Of the 17 cloned kittens born, seven were stillborn, eight died within hours of delivery or up to 6 weeks of age, and two are currently alive and healthy. Additionally, some studies on wild felids have shown the establishment of somatic resource banks. There is a bank with somatic samples of 69 individual Iberian lynx, considered the most endangered felid in the world, with the aim of future cloning (León-Quinto *et al.*, 2009; 2014).

In addition, works on the cheetah (*Acinonyx jubatus*), a species of the Pantherinae subfamily, have been performed in South America and Asia. Somatic cells from a cheetah raised in South America were transferred to domestic cat oocytes, and, after embryo aggregation during *in vitro* culture, high blastocyst formation rates were obtained (16.7–28.3%) (Moro *et al.*, 2015). Moulavi *et al.* (2017) used non-viable frozen cells derived from frozen tissue from an Asiatic cheetah (*Acinonyx jubatus venaticus*) and *in vitro*-matured domestic cat oocytes and obtained morula rates of 5.9%. Although no blastocyst was obtained, this study demonstrated that enucleated cat oocytes can partially remodel and reactivate nonviable nuclei of the Asiatic cheetah and support its reprogramming back to the embryonic stage.

Ungulates

Some studies with ungulates have been performed with significant success, especially for species already extinct. The first animal

derived from an extinct subspecies was obtained using fibroblasts from skin biopsies collected before the death of the last female *Capra pyrenaica pyrenaica*. After a year under cryopreservation, these cells were used as karyoplasts and fused with the cytoplasts of a domestic goat to reconstruct embryos. The rate of cleaved embryos after 36 h was 47.3%, of which 65.5% were transferred. Five recipients were pregnant at 45 days but only one pregnancy went to term. Unfortunately, a few minutes after birth the animal died from pulmonary complications (Folch *et al.*, 2009).

Experiments were carried out with wild yak (*Bros grunniens*) with the goal of evaluating the parameters that affect the success of iSCNT (Li *et al.*, 2007). Fibroblasts and cumulus cells were used as donor cells, but the cell type and different ages were found to have no significant effect on iSCNT.

In 2017, the birth of a Bactrian camel cloned by iSCNT was first reported (Wani *et al.*, 2017). The fibroblasts used to donate nuclei were obtained from ear skin biopsy samples from an adult male Bactrian camel (*Camelus bactrianus*) and the cytoplasm of dromedary camel (*Camelus dromedaries*) was the oocyte recipient. Twenty-six blastocysts were transferred to 23 synchronized dromedary recipients yielding five pregnancies with one going to term. This work has great importance because the Bactrian camel is the eighth most endangered large mammal on Earth.

Finally, the woolly mammoth (*Mammuthus primigenius*) is perhaps the one wild mammal of the ungulates whose cloning arouses the greatest interest. This animal became extinct about 10,000 years ago. However, epithelial and muscular cells from 14,000–15,000-year-old mammoth tissues were cryopreserved, with the goal of producing embryos of this species (Kato *et al.*, 2009). In this study, the authors injected cell nucleus-like structures into mature mouse enucleated oocytes; however, the oocytes did not form pronuclear-like structures at 7 h after injection.

Other species

The ability of bovine enucleated oocytes to support dedifferentiation of nuclei from monkey fibroblasts in interspecies cloned monkey embryos has been observed (Lorthongpanich *et al.*, 2008). These embryos were cultured in conditions different from the medium used for cattle with monkey-specific alterations, but the embryos were not able to develop past 16 cells under any culture condition. Nevertheless, OCT-4 was detected, demonstrating the ability of bovine ooplasm to support dedifferentiation but not embryonic development. Therefore, the culture medium promotes dedifferentiation but is not able to support complete embryonic development (Lorthongpanich *et al.*, 2008). In another work that used porcine cytoplasts and donor cells from a rhesus monkey, it was possible to obtain blastocysts despite the low rate (2.04%) (Zhu *et al.*, 2014). Although being a SCNT study, the cloning of cynomolgus monkeys (*Macaca fascicularis*) is cited here because of recent advances in this species. Thus, in a study on cynomolgus monkeys using SCNT, Liu *et al.* (2018) applied histone demethylase Kdm4d mRNA and histone deacetylase inhibitor trichostatin A after activation. Embryonic development improved followed by a greater number of pregnancies, which resulted in the birth of two monkeys via the SCNT technique using fetal fibroblasts and oocytes of cynomolgus monkeys.

With respect to aquatic mammals, a study performed on the minke whale (*Balaenoptera bonaerensis*) compared different conditions of iSCNT, including the ability of porcine and bovine ooplasm to produce reconstructed embryos and the effects of different donor cell types (viable or nonviable cells) on whale

SCNT embryos (Ikumi *et al.*, 2004). The authors concluded that whale iSCNT embryos can develop to at least the four-cell stage, regardless of the survivability of the donor cells and the porcine or bovine ooplasm.

Final considerations

Although cloning has several technical limitations that require greater attention to improve the technique, iSCNT has been applied to numerous species of wild mammals and has achieved positive results with respect to embryonic stages in pregnancies and offspring born. The works cited in this paper have made it possible to analyze the state of the art and to perform specific studies the problems in the technique that can be fixed according to the species being studied.

This review has shown that there is no rule that says several species should be cloned following the same protocol, but that each species has different needs at each stage of the technique. In addition, all the papers referred to in this review point to the need for improvement and study at a certain stage, which will lead to improvement of the technique. Thus, to achieve a satisfactory result with iSCNT, each step involved in cloning must be suitable for the species being studied.

Although iSCNT is not the main tool for the reestablishment of endangered wild mammals, its use to increase the possibilities of reproduction and multiplication of individuals has been proposed. It should be refined so that it can be an alternative when traditional techniques cannot be applied. In addition, cloning helps elucidate the embryonic development of a wild species and the subsequent application of this knowledge.

Finally, this biotechnology can help generate more ways to maintain individual species. Therefore, the improvement of protocols to potentiate this technique is of interest because although it has low efficiency rates, iSCNT shows promise because of the pups of different species that have been born.

Financial support. Alana Azevedo Borges is a recipient of a grant from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES, Financial Code 001). Alessandra Fernandes Pereira is recipient of a grant from CNPq (no. 306963/2017-5).

Conflicts of interest. None of the authors has any conflict of interest to declare.

Ethical standards. Not applicable

References

- Berg DK, Li C, Asher G, Wells DN and Oback B (2007) Red deer cloned from antler stem cells and their differentiated progeny. *Biol Reprod* 77, 384–94.
- Borges AA, Bezerra FVB, Costa FN, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR and Pereira AF (2017a) Histomorphological characterization of collared peccary (*Pecari tajacu* Linnaeus, 1758) ear integumentary system. *Arq Bras Med Vet Zootec* 69, 948–54.
- Borges AA, Lima GL, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR and Pereira AF (2017b) Conservation of somatic tissue derived from collared peccaries (*Pecari tajacu* Linnaeus, 1758) using direct or solid-surface vitrification techniques. *Cytotechnology* 69, 643–54.
- Borges AA, Lira GPO, Nascimento LE, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR and Pereira AF (2018a) Influence of

- cryopreservation solution on the *in vitro* culture of skin tissues derived from collared peccary (*Pecari tajacu* Linnaeus, 1758) *Biopreserv Biobank* 16, 77–81.
- Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR and Pereira AF (2018b) Combination of ethylene glycol with sucrose increases survival rate after vitrification of somatic tissue of collared peccaries (*Pecari tajacu* Linnaeus, 1758) *Pesq Vet Bras* 38, 350–6.
- Borges AA, Santos MVO, Queiroz Neta LB, Oliveira MF, Silva AR and Pereira AF (2018c) *In vitro* maturation of collared peccary (*Pecari tajacu* Linnaeus, 1758) oocytes after different incubation times. *Pesq Vet Bras* 38, 1863–8.
- Brahmasani SR, Yelisetti UM, Katari V, Komjeti S, Lakshmikantham U, Pawar RM and Sisinthy S (2013) Developmental ability after parthenogenetic activation of *in vitro* matured oocytes collected postmortem from deers. *Small Rumin Res* 113, 128–35.
- Do VH and Taylor-Robinson A (2014) Somatic cell nuclear transfer in mammals reprogramming mechanism and factors affecting success. *Clon Transgen* 3, 129.
- Fernandez-Gonzalez L, Hribal R, Stagegaard J, Zahmel J and Jewgenow K (2015) Production of lion (*Panthera leo*) blastocysts after *in vitro* maturation of oocytes and intracytoplasmic sperm injection. *Theriogenology* 83, 995–9.
- Folch J, Cocero M, Chesné P, Alabart J, Domínguez V, Cognié Y, Roche A, Fernández-Arias A, Martí J and Sánchez P (2009) First birth of an animal from an extinct subspecies (*Capra pyrenaica pyrenaica*) by cloning. *Theriogenology* 71, 1026–34.
- Goeritz F, Painer J, Jewgenow K, Hermes R, Rasmussen K, Dehnhard M and Hildebrandt T (2012) Embryo retrieval after hormonal treatment to control ovarian function and non-surgical artificial insemination in African lions (*Panthera leo*) *Reprod Domest Anim* 47, 156–60.
- Gómez MC, Jenkins JA, Giraldo A, Harris RF, King A, Dresser BL and Pope CE (2003) Nuclear transfer of synchronized African wild cat somatic cells into enucleated domestic cat oocytes. *Biol Reprod* 69, 1032–41.
- Gómez MC, Pope CE, Giraldo A, Lyons LA, Harris RF, King AL, Cole A, Godke RA and Dresser BL (2004) Birth of African wild cat cloned kittens born from domestic cats. *Cloning Stem Cells* 6, 247–58.
- Gómez MC, Pope CE, Kutner RH, Ricks DM, Lyons LA, Ruhe M, Dumas C, Lyons J, López M and Dresser BL (2008) Nuclear transfer of sand cat cells into enucleated domestic cat oocytes is affected by cryopreservation of donor cells. *Cloning Stem Cells* 10, 469–84.
- González-Grajales LA, Favetta LA, King WA and Mastromonaco GF (2016) Lack of effects of ooplasm transfer on early development of interspecies somatic cell nuclear transfer bison embryos. *BMC Dev Biol* 16, 36.
- Herrick J, Campbell M, Levens G, Moore T, Benson K, D'Agostino J, West G, Okeson D, Coke R and Portacio S (2010) *In vitro* fertilization and sperm cryopreservation in the black-footed cat (*Felis nigripes*) and sand cat (*Felis margarita*) *Biol Reprod* 82, 552–62.
- Howard J, Lynch C, Santymire R, Marinari P and Wildt D (2016) Recovery of gene diversity using long-term cryopreserved spermatozoa and artificial insemination in the endangered black-footed ferret. *Anim Conserv* 19, 102–11.
- Ikumi S, Sawai K, Takeuchi Y, Iwayama H, Ishikawa H, Ohsumi S and Fukui Y (2004) Interspecies somatic cell nuclear transfer for *in vitro* production of Antarctic minke whale (*Balaenoptera bonaerensis*) embryos. *Cloning Stem Cells* 6, 284–93.
- Kato H, Anzai M, Mitani T, Morita M, Nishiyama Y, Nakao A, Kondo K, Lazarev PA, Ohtani T and Shibata Y (2009) Recovery of cell nuclei from 15,000 years old mammoth tissues and its injection into mouse enucleated matured oocytes. *Proc Jpn Acad Ser B Phys Biol Sci* 85, 240–7.
- Kim MK, Jang G, Oh HJ, Yuda F, Kim HJ, Hwang WS, Hossein MS, Kim JJ, Shin NS and Kang SK (2007) Endangered wolves cloned from adult somatic cells. *Cloning Stem Cells* 9, 130–7.
- Lanza RP, Cibelli JB, Diaz F, Moraes CT, Farin PW, Farin CE, Hammer CJ, West MD and Damiani P (2000) Cloning of an endangered species (*Bos gaurus*) using interspecies nuclear transfer. *Cloning* 2, 79–90.
- Lee B, Wirtu GG, Damiani P, Pope E, Dresser BL, Hwang W and Bavister BD (2003) Blastocyst development after intergeneric nuclear transfer of mountain bongo antelope somatic cells into bovine oocytes. *Cloning Stem Cells* 5, 25–33.
- León-Quinto T, Simon MA, Cadenas R, Jones J, Martínez-Hernández FJ, Moreno JM, Vargas A, Martínez-Hernández FJ and Soria B (2009) Developing biological resource banks as a supporting tool for wildlife reproduction and conservation: the Iberian lynx bank as a model for other endangered species. *Anim Reprod Sci* 112, 347–61.
- León-Quinto T, Simón MA, Cadenas R, Martínez Á and Serna A (2014) Different cryopreservation requirements in foetal versus adult skin cells from an endangered mammal, the Iberian lynx (*Lynx pardinus*) *Cryobiology* 68, 227–33.
- Li Y, Dai Y, Du W, Zhao C, Wang L, Wang H, Liu Y, Li R and Li N (2007) *In vitro* development of yak (*Bos grunniens*) embryos generated by interspecies nuclear transfer. *Anim Reprod Sci* 101, 45–59.
- Liu Z, Cai Y, Wang Y, Nie Y, Zhang C, Xu Y, Zhang X, Lu Y, Wang Z and Poo M (2018) Cloning of macaque monkeys by somatic cell nuclear transfer. *Cell* 172, 881–7.
- Loi P, Modlinski J and Ptak G (2011) Interspecies somatic cell nuclear transfer: a salvage tool seeking first aid. *Theriogenology* 76, 217–28.
- Loi P, Ptak G, Barboni B, Fulka J Jr, Cappai P and Clinton M (2001) Genetic rescue of an endangered mammal by cross-species nuclear transfer using *post-mortem* somatic cells. *Nat Biotech* 19, 962.
- Lorthongpanich C, Laowtammathron C, Chan AWS, Ketudat-Cairns M and Parnpai R (2008) Development of interspecies cloned monkey embryos reconstructed with bovine enucleated oocytes. *J Reprod Dev* 54, 306–13.
- Macías-García B, González-Fernández L, Matilla E, Hernández N, Mijares J, and Sánchez-Margallo FM (2018) Oocyte holding in the Iberian red deer (*Cervus elaphus hispanicus*): Effect of initial oocyte quality and epidermal growth factor addition on *in vitro* maturation. *Reprod Domest Anim* 53, 243–8.
- Mestre-Citrinovit AC, Sestelo AJ, Ceballos MB, Baranao JL and Saraguet P (2016) Isolation of primary fibroblast culture from wildlife: the *Panthera onca* case to preserve a South American endangered species. *Curr Protoc Mol Biol* 116, 28.7.1–14.
- Moro LN, Hiriart MI, Buemo C, Jarazo J, Sestelo A, Veraguas D, Rodríguez-Alvarez L and Salamone DF (2015) Cheetah interspecific SCNT followed by embryo aggregation improves *in vitro* development but not pluripotent gene expression. *Reproduction* 150, 1–10.
- Moulavi F, Hosseini SM, Tanhaie-Vash N, Ostadhosseini S, Hosseini SH, Hajinasrollah M, Asghari M, Gourabi H, Shahverdi A and Vosough A (2017) Interspecies somatic cell nuclear transfer in Asiatic cheetah using nuclei derived from post-mortem frozen tissue in absence of cryoprotectant and *in vitro* matured domestic cat oocytes. *Theriogenology* 90, 197–203.
- Oh H, Kim M, Jang G, Kim H, Hong S, Park J, Park K, Park C, Sohn S and Kim DY (2008) Cloning endangered gray wolves (*Canis lupus*) from somatic cells collected *postmortem*. *Theriogenology* 70, 638–47.
- Pan X, Zhang Y, Guo Z and Wang F (2014) Development of interspecies nuclear transfer embryos reconstructed with argali (*Ovis ammon*) somatic cells and sheep ooplasm. *Cell Biol Inter* 38, 211–8.
- Pereira AF and Freitas VJF (2009) Cloning in ruminants: progress and current perspectives. *Rev Bras Reprod Anim* 33, 118–28.
- Pereira AF, Feltrin C, Almeida KC, Carneiro IS, Avelar SRG, Alcântara Neto AS, Sousa FC, Melo CHS, Moura RR, Teixeira DIA, Bertolini LR, Freitas VJF and Bertolini M (2013) Analysis of factors contributing to the efficiency of the *in vitro* production of transgenic goat embryos (*Capra hircus*) by handmade cloning (HMC) *Small Rumin Res* 109, 163–72.
- Pereira AF, Melo LM, Freitas VJF and Salamone DF (2015) Phosphorylated H2AX in parthenogenetically activated, *in vitro* fertilized and cloned bovine embryos. *Zygote* 23, 485–93.
- Pereira AF, Santos MLT, Borges AA, Queiroz Neta LB, Santos MVO and Feitosa AKN (2014) Isolation and characterization of skin-derived donor cells for nuclear transfer. *Acta Vet Brasilica* 8, 311–6.
- Pereira AF, Silva AR, Lima GL and Silva AM (2016) Somatic and gonadal tissue cryopreservation: an alternative tool for the germplasm conservation in wild mammals. In: Melanie Walton (ed.). *Germplasm: Characteristics, Diversity and Preservation*. Nova Science Publishers, New York, pp. 80–117.
- Priya D, Selokar N, Raja A, Saini M, Sahare A, Nala N, Palta P, Chauhan M, Manik R and Singla S (2014) Production of wild buffalo (*Bubalus*

- arnee) embryos by interspecies somatic cell nuclear transfer using domestic buffalo (*Bubalus bubalis*) oocytes. *Reprod Dom Anim* **49**, 343–51.
- Queiroz Neta LB, Lira GPO, Borges AA Santos MVO, Silva MB, Oliveira LRM, Silva AR, Oliveira MF and Pereira AF (2018) Influence of storage time and nutrient medium on recovery of fibroblast-like cells from refrigerated collared peccary (*Pecari tajacu* Linnaeus, 1758) skin. *In Vitro Cell Dev Biol Anim* **54**, 486–95.
- Rao BS, Mahesh YU, Lakshmikantan UR, Suman K, Charan KV and Shivaji S (2010) Developmental competence of oocytes recovered from *postmortem* ovaries of the endangered Indian blackbuck (*Antelope cervicapra*). *J Reprod Dev* **56**, 623–9.
- Rho GJ, Hahnel AC and Betteridge KJ (2001) Comparison of oocyte maturation times and of three methods of sperm preparation for their effects on the production of goat embryos *in vitro*. *Theriogenology* **56**, 503–16.
- Ruiz J, Landeo L, Mendoza J, Correa J, Silva M and Ratto MH (2015) *In vitro* developmental competence of alpaca (*Vicugna pacos*) and llama (*Lama glama*) oocytes after parthenogenetic activation. *Small Rumin Res* **133**, 148–52.
- Saini M, Selokar N, Raja A, Sahare A, Singla S, Chauhan M, Manik R and Palta P (2015) Effect of donor cell type on developmental competence, quality, gene expression, and epigenetic status of interspecies cloned embryos produced using cells from wild buffalo and oocytes from domestic buffalo. *Theriogenology* **84**, 101–8.
- Sansinena M, Hylan D, Hebert K, Denniston R and Godke R (2005) Banteng (*Bos javanicus*) embryos and pregnancies produced by interspecies nuclear transfer. *Theriogenology* **63**, 1081–91.
- Santos MLT, Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR and Pereira AF (2016) *In vitro* culture of somatic cells derived from ear tissue of collared peccary (*Pecari tajacu* Linnaeus, 1758) in medium with different requirements. *Pesq Vet Bras* **36**, 1194–202.
- Saragusty J, Diecke S, Drukker M, Durrant B, Friedrich Ben-Nun I, Galli C, Göritz F, Hayashi K, Hermes R and Holtze S (2016) Rewinding the process of mammalian extinction. *Zool Biol* **35**, 280–92.
- Sharma R, Sharma H, Ahlawat S, Aggarwal R, Vij P and Tania M (2018) First attempt on somatic cell cryopreservation of critically endangered *Camelus bactrianus* of India. *Gene Rep* **10**, 109–15.
- Song J, Hua S, Song K and Zhang Y (2007) Culture, characteristics and chromosome complement of Siberian tiger fibroblasts for nuclear transfer. *In Vitro Cell Dev Biol Anim* **43**, 203–9.
- Sparman ML, Tachibana M and Mitalipov SM (2010) Cloning of non-human primates: the road “less traveled by”. *Int J Dev Biol* **54**, 1671.
- Sukparangsi W, Bootsri R, Sikeao W, Karoon S and Thongphakdee A (2018) Establishment of induced pluripotent stem cells from fishing cat and clouded leopard using integration-free method for wildlife conservation. *Reprod Fertil Dev* **30**, 230.
- Tulake K, Yanagawa Y, Takahashi Y, Katagiri S, Higaki S, Koyama K, Wang X and Li H (2014) Effects of ovarian storage condition on *in vitro* maturation of Hokkaido sika deer (*Cervus nippon yesoensis*) oocytes. *Jpn J Vet Res* **62**, 187–92.
- Wang L, Peng T, Zhu H, Lv Z, Liu T, Shuai Z, Gao H, Cai T, Cao X and Wang H (2007) *In vitro* development of reconstructed ibex (*Capra ibex*) embryos by nuclear transfer using goat (*Capra hircus*) oocytes. *Small Rumin Res* **73**, 135–41.
- Wani NA, Vettical BS and Hong SB (2017) First cloned Bactrian camel (*Camelus bactrianus*) calf produced by interspecies somatic cell nuclear transfer: A step towards preserving the critically endangered wild Bactrian camels. *PLoS One* **12**, e0177800.
- Yamochi T, Kida Y, Oh N, Ohta S, Amano T, Anzai M, Kato H, Kishigami S, Mitani T and Matsumoto K (2013) Development of interspecies cloned embryos reconstructed with rabbit (*Oryctolagus cuniculus*) oocytes and cynomolgus monkey (*Macaca fascicularis*) fibroblast cell nuclei. *Zygote* **21**, 358–66.
- Yelisetti UM, Komjeti S, Katari VC, Sisinthy S and Brahmasani SR (2016) Interspecies nuclear transfer using fibroblasts from leopard, tiger, and lion ear piece collected *postmortem* as donor cells and rabbit oocytes as recipients. *In Vitro Cell Dev Biol Anim* **52**, 632–45.
- Yin Y, Tang L, Zhang P, Kong D, Wang Z, Guan J and Li Z (2013) Optimizing the conditions for *in vitro* maturation and artificial activation of sika deer (*Cervus nippon hortulorum*) oocytes. *Reprod Domest Anim* **48**, 27–32.
- Zhao ZJ, Ouyang YC, Nan CL, Lei ZL, Song XF, Sun QY and Chen DY, (2006) Rabbit oocyte cytoplasm supports development of nuclear transfer embryos derived from the somatic cells of the camel and Tibetan antelope. *J Reprod Dev* **52**, 449–59.
- Zhu HY, Kang JD, Li S, Jin JX, Hong Y, Jin L, Guo Q, Gao QS, Yan CG and Yin XJ (2014) Production of rhesus monkey cloned embryos expressing monomeric red fluorescent protein by interspecies somatic cell nuclear transfer. *Biochem Biophys Res Commun* **6**, 38–43.

ANEXO B – VERSÃO DO ARTIGO PUBLICADO NA PEERJ:

**ISOLATION, CHARACTERIZATION, AND CRYOPRESERVATION OF
COLLARED PECCARY SKIN-DERIVED FIBROBLAST CELL LINES**

QUALIS: Quadriênio 2013-2016: Não classificado pela Medicina Veterinária/ Novo Qualis:
A2

FATOR DE IMPACTO: 2,353

DOI: <https://doi.org/10.7717/peerj.9136>

Isolation, characterization, and cryopreservation of collared peccary skin-derived fibroblast cell lines

Alana Azevedo Borges¹, Gabriela Pereira De Oliveira Lira¹, Lucas Emanuel Nascimento¹, Maria Valéria De Oliveira Santos¹, Moacir Franco De Oliveira², Alexandre Rodrigues Silva³ and Alexsandra Fernandes Pereira¹

¹ Laboratory of Animal Biotechnology, Universidade Federal Rural do Semi-Árido, Mossoró, Rio Grande do Norte, Brazil

² Laboratory of Applied Animal Morphophysiology, Universidade Federal Rural do Semi-Árido, Mossoró, Rio Grande do Norte, Brazil

³ Laboratory of Animal Germplasm Conservation, Universidade Federal Rural do Semi-Árido, Mossoró, Rio Grande do Norte, Brazil

ABSTRACT

Background: Biobanking of cell lines is a promising tool of support for wildlife conservation. In particular, the ability to preserve fibroblast cell lines derived from collared peccaries is of significance as these wild mammals are unique to the Americas and play a large role in maintaining the ecosystem. We identified collared peccary fibroblasts by immunofluorescence and evaluated their morphology, growth and adherence capacity. Further, we monitored the viability and metabolic activity of the fibroblasts to determine the effects of passage number and cryopreservation on establishment of cell lines.

Methods: Skin biopsies were collected from the peripheral ear region from five adult animals in captivity. Initially, cells were isolated from fragments and cultured in the Dulbecco's modified Eagle medium supplemented with 10% fetal bovine serum and 2% antibiotic-antimycotic solution under a controlled atmosphere (38.5 °C, 5% CO₂). We evaluated the maintenance of primary cells for morphology, adherence capacity of explants, explants in subconfluence, cell growth and absence of contamination. Moreover, we identified the fibroblast cells by immunofluorescence. Additionally, to evaluate the influence of the number of passages (first, third and tenth passage) and cryopreservation on establishment of cell lines, fibroblasts were analysed for the viability, metabolic activity, population doubling time (PDT), levels of reactive oxygen species (ROS), and mitochondrial membrane potential ($\Delta\Psi_m$).

Results: All explants (20/20) adhered to the dish in 2.4 days $\pm$ 0.5 with growth around the explants in 4.6 days $\pm$ 0.7, and subconfluence was observed within 7.8 days $\pm$ 1.0. Moreover, by morphology and immunocytochemistry analyses, cells were identified as fibroblasts which presented oval nuclei, a fusiform shape and positive vimentin staining. No contamination was observed after culture without antibiotics and antifungals for 30 days. While there was no difference observed for cell viability after the passages (first vs. third: $P = 0.98$; first vs. tenth: $P = 0.76$; third vs. tenth: $P = 0.85$), metabolic activity was found to be reduced in the tenth passage ($23.2 \pm 12.1\%$) when compared to that in the first and third passage ($100.0 \pm 24.4\%$, $P = 0.006$). Moreover, the cryopreservation did not influence the viability

Submitted 20 November 2019

Accepted 15 April 2020

Published 3 June 2020

Corresponding author

Alexsandra Fernandes Pereira,
alexandra.pereira@ufersa.edu.br

Academic editor

Dorothee Bienzle

Additional Information and
Declarations can be found on
page 13

DOI 10.7717/peerj.9136

© Copyright

2020 Borges et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

($P = 0.11$), metabolic activity ($P = 0.77$), or PDT ($P = 0.11$). Nevertheless, a greater $\Delta\Psi_m$ ($P = 0.0001$) was observed for the cryopreserved cells (2.12 ± 0.14) when compared to that in the non-cryopreserved cells (1.00 ± 0.05). Additionally, the cryopreserved cells showed greater levels of intracellular ROS after thawing (1.69 ± 0.38 vs. 1.00 ± 0.22 , $P = 0.04$).

Conclusions: This study is the first report on isolation, characterization and cryopreservation of fibroblasts from collared peccaries. We showed that adherent cultures were efficient for obtaining fibroblasts, which can be used as donor cells for nuclei for species cloning and other applications.

Subjects Cell Biology, Conservation Biology, Veterinary Medicine

Keywords Biological characterization, Cellular conservation, Cryobanking, Wild mammals

INTRODUCTION

Collared peccaries (*Pecari tajacu* Linnaeus, 1758) are wild mammals found only in the Americas and show a distribution from southern United States to northern Argentina, inhabiting the most diverse environments (Santos et al., 2009). Currently, their population is considered to be stable (Gongora et al., 2011); however, a significant reduction of their population has been seen in some biomes, such as the Caatinga (Desbiez et al., 2012) and the Atlantic forest (Lazure et al., 2010). As excellent seed dispersers (Redford, 1992), they are very important for the maintenance of our ecosystem, whereas, economically, they have been commercialized for their meat and in leather production (Santos et al., 2009). Scientifically, collared peccaries can be used as experimental models for closely related species such as the *Tayassu peccary* and *Catagonus wagneri* that have been listed as “vulnerable” in the IUCN Red List of Threatened Species (Keuroghlian et al., 2013; Altrichter et al., 2015; Gongora et al., 2011).

In this sense, studies related to the conservation of the collared peccary have been intensified, especially aimed at improving the techniques related to the preservation of somatic samples. Using this study, we established a culture condition for explants derived from the skin of adult collared peccaries (Santos et al., 2016) and developed a protocol for cryopreservation (Borges et al., 2017, 2018a, 2018b) and refrigeration of these explants (Queiroz Neta et al., 2018). In order to conduct the cloning experiments on this species by a somatic cell nuclear transfer, as well as to produce induced pluripotent cells, it is necessary to establish properly characterized cell lines.

In general, as observed in other mammals (Guan et al., 2010; Kwong et al., 2014), establishment of an adequate cell line is a prerequisite step for the success of cloning and producing induced pluripotent cells (Borges & Pereira, 2019). For these techniques, fibroblasts and epithelial cells derived from the skin have been widely used (Jyotsana et al., 2016; Siengdee et al., 2018). Initially, epithelial and fibroblast cells were grown simultaneously; nevertheless, fibroblasts can more easily adhere as well as detach by trypsinization as compared to the epithelial cells (Bai et al., 2012; Saadeldin et al., 2019; Siengdee et al., 2018). In these methods, the culture after the second passage has been considered to contain mainly fibroblasts (Mehrabani et al., 2014).

Additionally, for the confirmation of a fibroblast line, it is necessary to verify the possible changes that occur in these cells during culture ([Guan et al., 2010](#); [Song et al., 2007](#)) and cryopreservation ([Magalhães et al., 2017](#)). In general, the number of passages throughout an in vitro study can modify the cellular epigenetic state, affecting the embryonic development after cloning ([Rodriguez-Osorio et al., 2012](#); [Trokovic et al., 2015](#)). [Magalhães et al. \(2017\)](#) observed reduced viability and metabolic activity in the cells derived from the skin of the brown brocket deer in the tenth passage. Thus, the establishment of a cell line ensures a complete knowledge of the parameters that confer quality to the nucleus of the donor cell, named the karyoplast ([Guan et al., 2010](#)). Moreover, identification of damage occurring during cryopreservation is essential for establishment of a cell line. Cryo-variables may affect several cellular processes, including survival, functionality and the cytoskeleton, which may compromise the reprogramming ability of the karyoplasts ([Chatterjee et al., 2017](#)). Therefore, we aimed to isolate, characterize and cryopreserve the fibroblast cells derived from the skin of the ear of collared peccaries for their future application in cloning strategies by a somatic cell nuclear transfer and production of induced pluripotent cells.

MATERIALS AND METHODS

Chemicals and media

The Dulbecco's modified Eagle medium (DMEM), fetal bovine serum (FBS), penicillin, streptomycin and amphotericin solutions were obtained from Gibco-BRL (Carlsbad, CA, USA). Fluorescent probes were purchased from Invitrogen (Carlsbad, CA, USA). Anti-vimentin antibody and goat anti-mouse IgG (Alexa Fluor[®] 488, Warrington, PA, USA) were purchased from Abcam (Cambridge, USA). The other chemicals were obtained from Sigma-Aldrich (St. Louis, MO, USA). Media were filtered using a 0.22-µm system (Corning, New York, USA) and adjusted to pH of 7.2–7.4.

Bioethics and animals

This study was approved by the Ethics Committee of Animal Use of the Federal Rural University of Semi-Arid (CEUA/UFERSA, no. 23091.001072/2015-92) and the Chico Mendes Institute for Biodiversity Conservation (no. 48633-2; ICMBio, Brasília, Brazil). All animals belonged to the Centre of Multiplication of Wild Animals (CEMAS/UFERSA, Mossoró, RN, Brazil, 5°10'S, 37°10'W), registered at the Brazilian Institute of Environment and Renewable Natural Resources (IBAMA) as a scientific breeding site (no. 1478912). The breeder stocks 100 collared peccaries on an average, and for this research four females and one male at ages of 26.8 months ± 2.9 months were used.

Ear tissue explant collection and primary culture

Peripheral skin (1–2 cm²) was recovered from the ear sections used to identify collared peccaries kept in captivity. After the collection, a trichotomy of the tissue followed by a sterilization with 70% alcohol was performed. Samples were transported to the laboratory in DMEM supplemented with 2% antibiotic–antimycotic solution (10,000 units/mL

penicillin, 10,000 µg/mL streptomycin and 25 µg/mL amphotericin B) at 37 °C within 30 min.

In the laboratory, fragments (9.0 mm³) were washed sequentially under laminar flow in the following media: (1) DMEM supplemented with 10% FBS and 10% antibiotic–antimycotic solution; (2) alcohol; and (3) DMEM plus 10% FBS and 2% antibiotic–antimycotic solution. Then, the samples were fragmented (four fragments per animal) and placed in polystyrene culture dishes treated for cell adhesion with the latter medium for cell culture. The skin was cultured at 38.5 °C under a controlled environment with 5% CO₂ and 95% air, according to a method described by [Santos et al. \(2016\)](#).

Evaluation of the somatic cells in primary cultures and subcultures

During primary culture, the medium was changed every 24 h. For evaluation of the somatic cells, the primary culture was analyzed before reaching confluency and until it reached a confluency of 70–80%. Using an inverted microscope (Nikon TS100, Tokyo, Japan), the cells were evaluated for the following parameters: cell morphology, number of adhered samples, number of samples, evident subconfluency, day of sample adherence, day of subconfluent growth of the samples, and total time to reach 70–80% confluence ([Borges et al., 2017](#)).

When the cells reached 70–80% subconfluency, they were subcultured and distributed for other analyses. The 70–80% subconfluence was defined as the stage when 70–80% of the culture dishes consisted of somatic cells ([Santos et al., 2016](#)). Subconfluent cells were washed with PBS then trypsinized with a trypsin/EDTA solution (0.25%/0.2%) for 7 min and centrifuged at 600×g for 10 min. The supernatant was removed, the cell pellet was resuspended in culture medium, and the cell suspension was transferred to another dish for subculturing ([Borges et al., 2018b](#)). The medium was replaced with fresh medium every other day and the cells were monitored daily. With the successful passaging of the cultures, the cells are considered a cell line, following the convention of the Society of In Vitro Biology ([Schaeffer & Terminology Committee Chair Tissue Culture Association, 1990](#)). The cell line was designated as Ptskf.

Thus, in addition to an evaluation of the maintenance of cells in the primary culture, the subcultured cells were initially evaluated for the confirmation of fibroblasts using morphology and immunofluorescence analyses. Moreover, the possibility of contamination was also evaluated. Subsequently, the influence of the number of passages (first, third and tenth passage) and the metabolic activity of the cells were analyzed by a viability assay using trypan blue and the 3-(4, 5-dimethylthiazolyl)-2, 5-diphenyltetrazolium bromide (MTT) assay, respectively. Moreover, the cells were also evaluated for the effects of a slow freezing cryopreservation. Other than the above-mentioned tests, growth dynamics by quantification of the population doubling time (PDT), oxidative stress analysis for quantification of intracellular reactive oxygen species (ROS) levels using the fluorescent probe 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA), and assessment of the mitochondrial membrane potential (ΔΨ_m) using the fluorescent probe MitoTrackerRed[®] were performed.

Morphological characterization of the fibroblasts

Morphological characteristics were observed throughout the in vitro culture under light microscopy for cellular and nuclear shapes and cytoplasmic extensions.

Vimentin immunofluorescence

For a morphological confirmation, the cells were subjected to an immunocytochemistry protocol based on the method described by [Amoli et al. \(2017\)](#). Briefly, the cells were fixed using 4% paraformaldehyde for 10 min at 25 °C, then washed with chilled PBS. Subsequently, cells were incubated with an antigen-retrieval buffer (100 mM Tris, 5% urea, pH 9.5), and then permeabilized for 1 h in 0.4% Triton X-100. Afterwards, the cells were incubated in 0.1% Tween-20 for 1 h to block non-specific binding of the antibodies. Finally, the cells were immuno-stained with mouse anti-vimentin antibody (ab8979, 1:200) for 24 h at 4 °C, and, then incubated with the secondary antibody (goat anti-mouse IgG, Alexa Fluor® 488, Warrington, PA, USA, ab150113, 1:400) for 1 h at 25 °C in the dark. Cells were counter-marked with one µg/mL Hoechst for 1 min and observed under a fluorescence microscope (Olympus BX51TF, Tokyo, Japan).

Confirmation of the absence of bacterial and fungal contamination

Cells of the third passage were cultured for 30 days in DMEM containing 10% FBS in the absence of an antibiotic–antimycotic solution, at 38.5 °C, 5% CO₂ and 95% air. Daily evaluation was performed under light microscopy for the identification of bacterial and fungal contamination.

Influence of the passage number on the quality of fibroblast lines

Initially, the fibroblast cells were analyzed for the effect of the number of passages (first, third and tenth passage) by a viability assay using trypan blue, according to the method described by [Magalhães et al. \(2017\)](#). We evaluated these three cell passages specifically because both fibroblast and epithelial cells were present in the initial (first) passage, only fibroblasts were visualized at the third passage onwards, and the cells of the tenth passage were used for most of the production of embryonic clones ([Shiga et al., 1999](#); [Kubota et al., 2000](#)). The evaluations were performed in triplicate for each animal for each passage.

Briefly, the cells were stained with 0.4% trypan blue in PBS and counted on a hemocytometer. Subsequently, the cells were also analyzed for a metabolic activity using the MTT assay, according to the method described by [Borges et al. \(2018b\)](#). A concentration of 5.0×10^4 cells/mL from the first, third and tenth passages was grown in 12-well polystyrene plates treated for cell adhesion. After 5 days, 1.5 mL of the MTT solution (five mg/mL in DMEM) was added and the polystyrene culture dishes treated for cell adhesion were incubated for 3 h. The MTT solution was then removed and 1.0 mL of dimethyl sulfoxide (DMSO) was added for 5 min under slow stirring to solubilize the MTT. After the total dissolution of formazan crystals, samples were analyzed in a spectrophotometer (Shimadzu® UV-mini-1240, Kyoto, Japan) at an absorbance

wavelength of 595 nm. The evaluations were performed in triplicate for each animal for each passage.

Influence of cryopreservation on the quality of fibroblast lines

To evaluate the effect of cryopreservation on the quality of fibroblast lines, cells of the third passage of the five animals were subjected to slow freezing in the freezing medium (DMEM supplemented with 10% DMSO as a permeating cryoprotectant and 10% FBS and 0.2 M sucrose as non-permeating cryoprotectants). Cells at a concentration of 5.0×10^4 cells/mL were first exposed to DMSO–FBS solution for 15 min at 4 °C, then sucrose solution was added followed by an additional incubation for 15 min at 4 °C. The cryovials containing 1.0 mL of cells in the freezing medium were cooled in a Mr. Frosty freezing container (Thermo Scientific, Waltham, MA, USA) at a cooling rate of 1 °C/min, and later stored in a freezer at –80 °C, reaching –70 °C overnight before being transferred into liquid nitrogen (León-Quinto *et al.*, 2014).

For thawing, the cryovials were exposed for 1 min at 25 °C and immersed in a water bath at 37 °C for 3–4 min. Then, the cell contents were removed from the cryovials and washed to remove the cryoprotectants. Initially, the first wash was performed with DMEM and 10% FBS containing 0.2 M sucrose at 4 °C for 15 min and centrifuged. Subsequently, the second wash was performed using only DMEM and 10% FBS, maintained at 25 °C for 15 min, centrifuged, and the cells were recovered for the evaluations as per a method described previously (Santos *et al.*, 2016).

After thawing, the non-cryopreserved and the cryopreserved cells were evaluated for growth dynamics by quantification of PDT. The evaluations were performed in triplicate for each animal. Briefly, the growth kinetics was studied for nine days using 3.0×10^4 cells/mL, and cells were counted daily to determine the number of growing cells. Data on the cell growth and density were monitored and recorded, mean values of which were used to plot a growth curve and calculate PDT (Roth, 2006) using the following formula:

$$PDT = T \ln_2 / \ln (X_e / X_b)$$
 where PDT is the time of the culture (in hours), T is the incubation time, X_b is the number of cells at the beginning of the time incubation, X_e is the number of cells at the end of the incubation time, and $\ln$ is the Napierian logarithm.

Moreover, for evaluation of an oxidative stress by quantification of the intracellular ROS levels, cells were stained with the fluorescent probe H₂DCFDA, according to a method described by Santos *et al.* (2019). Thawed cells were washed with PBS and placed into polystyrene culture dishes treated for cell adhesion containing 500 µL of 5 µM H₂DCFDA. The cells obtained after a 70% confluency were incubated at 38.5 °C in 5% CO₂ for 30 min. Stained cells were washed with PBS, placed on glass slides, photographed under a fluorescence microscope (Olympus BX51TF, Tokyo, Japan), and fluorescence signal intensity (pixels) was quantified. Ten images (two/animal) obtained were evaluated using the ImageJ software (version 1.49v, Java 1.8.0_201, Wayne Rasband, U.S. National Institutes of Health, Bethesda, MD, USA; website: <http://rsb.info.nih.gov/ij/download.html>). The background signal intensity was subtracted from the values obtained for the treated samples. Measured mean value of the micrograph for the non-cryopreserved cells

was taken as a calibrator. Relative expression levels (arbitrary fluorescence units) were generated by dividing the measured value of each micrograph for the cryopreserved cells by the mean of the calibrator.

Finally, for the assessment of $\Delta\Psi_m$, cells were stained using 500 nM of the fluorescent probe MitoTracker Red[®] (CMXRos), according to a method described by Santos *et al.* (2019). The procedure, incubation, and evaluation of the ten images (two/animal) were performed as described for the quantification of ROS.

Statistical analysis

All data have been expressed as the mean $\pm$ standard error (one animal/one repetition) and were analyzed using the StatView 5.0 software (Graph-Pad Software Incorporation, La Jolla, CA, USA). Normality of all results was verified by the Shapiro–Wilk test and homoscedasticity was verified by the Levene’s test. ROS levels, $\Delta\Psi_m$, viability, and metabolic activity were altered with arcsine and analysed by variance analysis (ANOVA) followed by the Tukey’s test. PDT was compared with ANOVA followed by the unpaired *t*-test. Statistical significance was set at $P < 0.05$.

RESULTS

Evaluation of the somatic cells in the primary cultures and subcultures

The total culture time was 95 days with an evaluation of cells until the tenth passage. The adhesion of the fragments (Fig. 1A), detachment of cells (Figs. 1B and 1C), and proliferative capacity were observed in all the explants until reaching a confluence (and later, a subconfluence) around the adhered fragments (Figs. 1D–1F; Table 1). All explants had adhesion ability and reached subconfluence. Number of days for each explant to reach a 100% tissue adherence ($2.4 \text{ days} \pm 0.5 \text{ days}$), to grow around the explants ($4.6 \text{ days} \pm 0.7 \text{ day}$), and to reach subconfluence ($7.8 \text{ days} \pm 1.0 \text{ day}$), were different.

Morphological characterization of the fibroblasts

In cultures, monolayers of cells with a fibroblast-like morphology were observed (Fig. 1E). The cells had an oval nuclei and extensions with a fusiform shape, showing rapid growth that replaced the epithelial cells.

Vimentin immunofluorescence

Morphology of the fibroblast-like cells in the initial culture was observed by light microscopy, which was further confirmed for the cell type identification as vimentin-labeled fibroblasts under fluorescence microscopy (Figs. 2A–2F). Cells exhibited a high expression of vimentin that marked the cytoplasm completely, and the spindle-like shape and ovoid nucleus was highlighted by the Hoechst labeling. Therefore, the identification of a fibroblast cell was evident.

Confirmation of the absence of bacterial and fungal contamination

No sign of contamination (turbidity, colony, or hyphal growth) was observed for 30 days in the culture without antibiotics and antifungals. The culture medium did not show any change in the appearance when observed under a light microscope. We did not observe




Figure 1 Outgrowth of fibroblast cells from ear skin samples of collared peccaries. Exhibit a skin explant cultured in (A) day 1, (B) day 3 and (C) day 5 of primary culture and exhibit a fibroblast population cultured in (D) day 15 and (E) day 19 of subculture and (F) exhibit cells after the trypsinization process. Scale bar = 100 μ m. [Full-size !\[\]\(09aa87a6a2999ef70143e2d9125279f4_img.jpg\) DOI: 10.7717/peerj.9136/fig-1](https://doi.org/10.7717/peerj.9136/fig-1)

Table 1 Establishment of primary culture and subcultures of somatic cells derived from collared peccary ear skin.

Animal	No. samples			No. attached samples		
	Initial	Attached (%)	Day all attached explants	Grow to subconfluence (%)	Day all cell grow explants	Subconfluence total time (days)
F1	4	100	2	100	4	6
F2	4	100	2	100	4	5
F3	4	100	1	100	3	9
F4	4	100	3	100	5	10
M1	4	100	4	100	7	9
Mean $\pm$ S.E.	20	100	2.4 $\pm$ 0.5	100	4.6 $\pm$ 0.7	7.8 $\pm$ 1.0

turbidity or any specific odor. In addition, there was no change in the biological characteristics of growth and proliferation indicating a complete absence of contamination.

Influence of the passage number on the quality of fibroblast lines

No significant difference was observed in the cell viability (74.5–84.4%) when evaluated by trypan blue staining after the passages (first vs. third: $P=0.98$; first vs. tenth: $P=0.76$; third




Figure 2 Immunocytochemical detection of vimentin protein for identification of collared peccary fibroblasts. (A–D) Cells stained with vimentin antibody. (B–E) Nucleus of cells stained by Hoechst. (C–F) Merged vimentin (green) and Hoechst (blue). A–C ($\times 5$), D–F ($\times 10$). Scale bar = 10 μm .

Full-size [DOI: 10.7717/peerj.9136/fig-2](https://doi.org/10.7717/peerj.9136/fig-2)




Figure 3 The growth dynamics of cryopreserved and non-cryopreserved cells derived from collared peccary skin. (A) Growth curves of cryopreserved and non-cryopreserved cells. (B) Values of population doubling time (PDT) after culture for nine days.

Full-size [DOI: 10.7717/peerj.9136/fig-3](https://doi.org/10.7717/peerj.9136/fig-3)

vs. tenth: $P = 0.85$). However, the metabolic activity was reduced in the tenth passage ($23.2 \pm 12.1\%$) as compared to that of the first and third passages ($100.0 \pm 24.4\%$, $P = 0.006$).

Influence of cryopreservation on the quality of fibroblast lines

Cryopreservation did not affect the viability when evaluated by trypan blue staining ($87.4 \pm 0.3\%$ vs. $74.0 \pm 5.9\%$, $P = 0.11$). Moreover, after two passages of the thawed cells, the viability was $86.4 \pm 3.2\%$. In addition, no difference ($P = 0.77$) was observed for the metabolic activity between the cryopreserved ($85.2 \pm 10.0\%$) and the non-cryopreserved cells ($100.0 \pm 36.4\%$).

Moreover, the cryopreserved and the non-cryopreserved cells were compared for growth dynamics (Fig. 3). The growth curve of both groups showed a typical “S-shaped”




Figure 4 Evaluation of intracellular reactive oxygen species (ROS) and mitochondrial membrane potential ($\Delta\Psi m$). Cell stained using fluorescent probe (A) 2',7'-dichlorodihydrofluorescein diacetate (H2DCFDA) and (B) MitoTrackerRed® (CMXRos) ($\times 10$). Quantification of (C) ROS and (D) $\Delta\Psi m$ levels. Scale bar = 10 μm . (*) Indicate statistical difference ($P < 0.05$).

Full-size DOI: 10.7717/peerj.9136/fig-4

pattern from the 9-day culture of cells. The latency time was 2 days, followed by an exponential phase until the fourth day, the stationary phase until the 7th day, and the plateau phase from day eight (Fig. 3A). No difference was observed for the PDT values of the cryopreserved and the non-cryopreserved cells ($P = 0.11$, Fig. 3B).

Nevertheless, the cryopreserved cells showed greater levels of intracellular ROS (Fig. 4A) in arbitrary fluorescence units when compared to that of the non-cryopreserved cells (1.69 ± 0.38 vs. 1.00 ± 0.22 , $P = 0.04$) (Fig. 4C). In addition, an alteration in the $\Delta\Psi m$ (Fig. 4B) in arbitrary fluorescence units ($P = 0.0001$) was observed for the cryopreserved cells (2.37 ± 0.07) when compared to that of the non-cryopreserved cells (1.00 ± 0.05) (Fig. 4D).

DISCUSSION

In this study, we isolated, characterized, and cryopreserved the fibroblast cells derived from the skin of collared peccaries. Moreover, we established the fibroblast cell lines of these animals with an aim to use these cells in cloning experiments by a somatic cell nuclear transfer in the future. The cell line can be considered as the first constituent of the peccary invitrome and a resource for future studies in many disciplines (Barrioch, 2018; Bols et al., 2017). Thus, the ear tissues of collared peccaries can be isolated and grown into fibroblasts in an adherent culture for establishment of cell lines and development of a cryobank. The development of these somatic cell banks has been increasing in the interest of conserving genetic samples of wild mammals to preserve valuable species, and as

sources for biological research (*León-Quinto et al., 2009; Mehrabani et al., 2014; Saadeldin et al., 2019; Siengdee et al., 2018*).

All explants adhered to the flask surface within 2–4 days, with cellular growth around the explant within 3 days, and demonstrating confluency within 5–10 days after a culture initiation. These characteristics of explants during in vitro culture were similar to the explants derived from other domestic and wild mammals. In studies using tissues from horses, the migration of fibroblast and epithelial-like cells from explants have been observed after 5–7 days of an in vitro culture (*Amoli et al., 2017*). In the case of goat-derived tissues, the explants reportedly adhered to the flasks within 5–7 days and the cells became confluent within 3–5 days post adhesion (*Bai et al., 2012*). In the Iranian Sistani cattle-derived tissues, the explants adhered to the culture flasks within 7–14 days and were observed to allow the growth of fibroblast-like cells from the margins of explants (*Gorji et al., 2017*).

For the Luxi cattle-derived tissues, fibroblast-like or epithelial-like cells could be seen migrating from the tissues within 5–12 days post adhesion (*Liu et al., 2008*). In the tissues derived from wild camels, fibroblast-like or epithelial-like cells could be seen migrating from the sides of explants within 8–10 days post adhesion (*Sharma et al., 2018*). In tissues derived from the domestic porcine, a species phylogenetically close to the collared peccaries, all the explants adhered within 3–8 days (*Silvestre, Sánchez & Gómez, 2004*). The similarity among these data can be related to the culture medium because in a majority of these studies, DMEM containing FBS, antibiotic and antimycotic solution was used (*Magalhães et al., 2017; Saadeldin et al., 2019; Siengdee et al., 2018*). Since primary culture needs to mimic the in vivo environment of the cells (*Guo et al., 2018*), we observed previously (*Santos et al., 2016*) that the medium for growth of somatic cells derived from collared peccaries was DMEM with 10% FBS and 2% antibiotic–antimycotic solution.

We showed that ear explant cultures obtained from the tissues of the collared peccaries were efficient for cell recovery allowing a culture for 95 days and up to 10 passages. Moreover, we confirmed that the ear skin was a common source where fibroblasts cells could be separated and eventually used as karyoplasts for cloning purposes (*Luo et al., 2014*). These cells were identified by vimentin, an intermediate filament that indicates the mesenchymal origin of endothelial and fibroblast cells (*Yajing et al., 2018*). Initially, during in vitro culture, epithelial and fibroblastic cells grew simultaneously. However, fibroblasts can be trypsinized more rapidly and adhere more easily as compared to the epithelial cells (*Bai et al., 2012; Saadeldin et al., 2019*). Therefore, in this work, cells from the third passage were confirmed as fibroblasts by morphology and immunofluorescence analyses.

The clear medium observed during the 30-day assay demonstrated the ability to allow the growth of the culture without any biological contamination. The propagation of fungi causes turbidity by accumulation of their metabolites. In addition, the colonies can be seen under a light microscope, or sometimes with the naked eye (*Li et al., 2009*). Bacterial contamination can also be identified by the naked eye as turbidity. One simple way to avoid this contamination is to filter the culture medium (*Bai et al., 2012*). Sources of

contamination may include, but are not limited to, the equipment, air, culture medium, serum and explant. Therefore, microbial contaminations are quite frequent in cell culture ([Bai et al., 2012](#)). Thus, the use of antibiotic and antimycotic combined with careful handling, is essential to ensure the absence of contamination.

After two passages, it was possible to separate fibroblast cells from other primary cells because different cell types exhibit different cellular behavior upon trypsinization. The fibroblast cells detach quicker than epithelial cells ([Gorji et al., 2017](#)). However, fibroblasts detach in response to trypsin more rapidly than epithelial cells and adhere more quickly ([Bai et al., 2012](#)). In wild camels, the initial lag phase of 48 h representing the adaptation of fibroblasts and recovery from a protease damage is followed by the exponential phase ([Sharma et al., 2018](#)). In collared peccaries, the replication of cells begins to slow down after 7 days because of contact inhibition, which, in wild camels, has been observed after 6 days ([Sharma et al., 2018](#)). No difference in the cell viability was observed among the first, third and the tenth passage, corroborating with the studies that used cells from these passages for production of competent cloned embryos ([Shiga et al., 1999](#); [Kubota et al., 2000](#)).

In contrast, through the metabolic activity test evaluated by the formation of formazan crystals, a significant reduction in the metabolic activity at the tenth passage was observed, indicating a reduced cellular functionality. Similar behavior was observed in cells from the brown brocket deer in which the metabolic activity measured by the MTT assay showed significantly lower values in the tenth passage than the values in the fourth passage ([Magalhães et al., 2017](#)). Therefore, the number of passages can reduce the metabolic activity rate and cell proliferation, thereby conserving cells of the early passages ([Li et al., 2009](#)). After several passages, genetic characteristics of the cells can be modified by culture conditions; hence, a minimum number of passages have been recommended to conserve the cellular characteristics ([Mehrabani et al., 2014](#)). Owing to this reason, the cells were cryopreserved in the third passage for the conservation of the somatic germplasm of collared peccaries.

The cell survival rate after thawing is the most commonly used criteria to evaluate the success of a cryopreservation ([Chatterjee et al., 2017](#)). The cellular viability and the functional metabolic activity of the cells were maintained after thawing the fibroblasts isolated from the collared peccaries. This factor demonstrates that optimal in vitro culture conditions significantly influence the recovery from cellular damages caused by the freezing process ([Gorji et al., 2017](#)). As for the growth curve, cryopreserved cells presented a very similar profile to that of the non-cryopreserved cells, showing their normal proliferation capacity regardless of the cryopreservation process. The establishment of somatic cell banks using cryopreservation technology is an easy and effective approach towards storing the genetic information of diverse species ([Li et al., 2009](#)). However, the cells should be handled with the utmost care during cryopreservation to maintain a high-quality cell bank in the long term ([Mehrabani et al., 2014](#)).

Moreover, epigenetic alterations, such as DNA fragmentation, free radical accumulation, ionic imbalances, apoptosis, biochemical alterations, DNA methylation and

histone modification can be a result of the cryopreservation (*Chatterjee et al., 2017*). These after-effects of cryopreservation may have caused mitochondrial structural abnormalities, thereby promoting an increased ROS production and H_2O_2 content, increased lipid peroxidation and increased expression of autophagic proteins harbored by the cells (*Mata et al., 2012*). A failure in the mitochondrial membrane potential is a hallmark of apoptosis, leading to the collapse of the organelle and release of cytochrome-C into the cytoplasm and ultimately activation of the apoptotic cascade (*Magalhães et al., 2012*). Moreover, a high $\Delta\Psi_m$ mitochondrial respiratory chain becomes a significant ROS producer (*Korshunov, Skulachev & Starkov, 1997*). Therefore, a higher $\Delta\Psi_m$ in cryopreserved cells can be linked mainly to an increase in the oxidative stress.

Finally, parameters like cryovariables, including cooling and thawing rates, type and concentration of the cryoprotectant, cell type and shape and nucleation temperature may affect the success of cryopreservation (*Chatterjee et al., 2017*). This suggests that the optimization of related cryopreservation methods for the collared peccary fibroblasts to minimize an altered $\Delta\Psi_m$ and increased levels of intracellular ROS production is essential.

CONCLUSIONS

To our knowledge, this study is the very first report on a successful isolation, characterization and cryopreservation of fibroblast lines derived from adult collared peccaries (Ptskf). We showed that the adherent culture was efficient for obtaining fibroblasts, which can be used as donor cells for nuclei for cloning of this species. Moreover, it was possible to maintain the viability of the cells until the tenth passage. In addition, cryopreservation did not affect the viability, metabolic activity and proliferative activity of the fibroblasts after slow freezing. However, cryopreservation altered the ROS levels and $\Delta\Psi_m$, indicating necessary optimization of the cryopreservation protocol. Lastly, the establishment of fibroblast cell lines derived from collared peccaries may be a source of experimental models for many biological studies such as nuclear reprogramming and animal cloning.

ACKNOWLEDGEMENTS

The authors thank the Centre for Wild Animals Multiplication (CEMAS/UFERSA) for providing the animals, and the Laboratory Biochemistry and Molecular Biology (BIOMOL/UERN) for technical assistance.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This study was supported by the Brazilian Council of Scientific Development (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior— Brasil (CAPES, Financial Code 001). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Brazilian Council of Scientific Development (CNPq).

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES): 001.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Alana Azevedo Borges conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Gabriela Pereira de Oliveira Lira performed the experiments, prepared figures and/or tables, and approved the final draft.
- Lucas Emanuel Nascimento performed the experiments, prepared figures and/or tables, and approved the final draft.
- Maria Valéria de Oliveira Santos performed the experiments, prepared figures and/or tables, and approved the final draft.
- Moacir Franco de Oliveira conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Alexandre Rodrigues Silva conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Alexsandra Fernandes Pereira conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

This study was approved by the Ethics Committee of Animal Use of the Federal Rural University of Semi-Arid (CEUA/UFERSA, no. 23091.001072/2015-92) and the Chico Mendes Institute for Biodiversity Conservation (ICMBio, no. 48633-2).

Data Availability

The following information was supplied regarding data availability:

The raw data are available in a [Supplemental File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.9136#supplemental-information>.

REFERENCES

- Altrichter M, Taber A, Noss A, Maffei L, Campo J. 2015. *Catagonus wagneri*. The IUCN Red List of Threatened Species 2015:e.T4015A72587993
DOI 10.2305/IUCN.UK.2015-2.RLTS.T4015A72587993.en.

- Amoli AD, Mohebbi N, Farzaneh P, Fazeli SAS, Nikfarjam L, Movasagh SA, Moradmam Z, Ganjibakhsh M, Nasimian A, Izadpanah M. 2017. Establishment and characterization of Caspian horse fibroblast cell bank in Iran. *In Vitro Cellular & Developmental Biology—Animal* 53:337–343 DOI 10.1007/s11626-016-0120-3.
- Bai C, Wang D, Su X, Zhang M, Guan W, Ma Y. 2012. Establishment and biological research of the Jining grey goat fibroblast line. *Turkish Journal of Veterinary and Animal Sciences* 36:659–667 DOI 10.3906/vet-1106-15.
- Barioch A. 2018. The Cellosaurus: a cell knowledge resource. *Journal of Biomolecular Techniques* 29(2):25–38 DOI 10.7171/jbt.18-2902-002.
- Bols NC, Pham PH, Dayeh VR, Lee LEJ. 2017. Invitromatics, invitrome, and invitroomics: introduction of three new terms for in vitro biology and illustration of their use with the cell lines from rainbow trout. *In Vitro Cellular & Developmental Biology—Animal* 53:383–405 DOI 10.1007/s11626-017-0142-5.
- Borges AA, Lima GL, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, Pereira AF. 2017. Conservation of somatic tissue derived from collared peccaries (*Pecari tajacu* Linnaeus, 1758) using direct or solid-surface vitrification techniques. *Cytotechnology* 69(4):643–654 DOI 10.1007/s10616-017-0074-7.
- Borges AA, Lira GPO, Nascimento LE, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, Pereira AF. 2018a. Influence of cryopreservation solution on the in vitro culture of skin tissues derived from collared peccary (*Pecari tajacu* Linnaeus, 1758). *Biopreservation and Biobanking* 16(2):77–81 DOI 10.1089/bio.2017.0090.
- Borges AA, Pereira AF. 2019. Potential role of intraspecific and interspecific cloning in the conservation of wild mammals. *Zygote* 27(3):111–117 DOI 10.1017/S0967199419000170.
- Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, Pereira AF. 2018b. Combination of ethylene glycol with sucrose increases survival rate after vitrification of somatic tissue of collared peccaries (*Pecari tajacu* Linnaeus, 1758). *Pesquisa Veterinária Brasileira* 38(2):350–356 DOI 10.1590/1678-5150-pvb-5193.
- Chatterjee A, Saha D, Niemann H, Gryshkov O, Glasmacher B, Hofmann N. 2017. Effects of cryopreservation on the epigenetic profile of cells. *Cryobiology* 74:1–7 DOI 10.1016/j.cryobiol.2016.12.002.
- Desbiez ALJ, Keuroghlian A, Mello Beisiegel B, Medici EP, Gatti A, Pontes ARM, Campos CB, Tófoli CF, Júnior EAM, Azevedo FC. 2012. Avaliação do risco de extinção do cateto *Pecari tajacu* Linnaeus, 1758, no Brasil. *Biodiversidade Brasileira* 3:74–83.
- Gongora J, Reyna-Hurtado R, Beck H, Taber A, Altrichter M, Keuroghlian A. 2011. *Pecari tajacu*. *The IUCN Red List of Threatened Species* 2011:e.T41777A10562361 DOI 10.2305/IUCN.UK.2011-2.RLTS.T41777A10562361.en.
- Gorji ZE, Khaledi KJ, Amoli AD, Ganjibakhsh M, Nasimian A, Gohari NS, Izadpanah M, Vakhshiteh F, Farghadan M, Moghanjoghi SM. 2017. Establishment and characteristics of Iranian Sistani cattle fibroblast bank: a way to genetic conservation. *Conservation Genetics Resources* 9(2):305–312 DOI 10.1007/s12686-016-0640-x.
- Guan W, Liu C, Li C, Liu D, Zhang W, Ma Y. 2010. Establishment and cryopreservation of a fibroblast cell line derived from Bengal tiger (*Panthera tigris tigris*). *Cryoletters* 31:130–138.
- Guo H, Xing Y, Zhang Y, He L, Deng F, Ma X, Li Y. 2018. Establishment of an immortalized mouse dermal papilla cell strain with optimized culture strategy. *PeerJ* 6:e4306 DOI 10.7717/peerj.4306.

- Jyotsana B, Sahare AA, Raja AK, Singh KP, Nala N, Singla S, Chauhan M, Manik R, Palta P. 2016. Use of peripheral blood for production of buffalo (*Bubalus bubalis*) embryos by handmade cloning. *Theriogenology* 86(5):1318–1324 DOI 10.1016/j.theriogenology.2016.04.073.
- Keuroghlian A, Desbiez R, Reyna-Hurtado M, Altrichter H, Beck A, Taber JMV. 2013. *Tayassu pecari*. The IUCN Red List of Threatened Species 2013:e.T41778A44051115 DOI 10.2305/IUCN.UK.2013-1.RLTS.T41778A44051115.en.
- Korshunov SS, Skulachev VP, Starkov AA. 1997. High protonic potential actuates a mechanism of production of reactive oxygen species in mitochondria. *FEBS Letters* 416(1):15–28 DOI 10.1016/S0014-5793(97)01159-9.
- Kubota C, Yamakuchi H, Todoroki J, Mizoshita K, Tabara N, Barber M, Yang X. 2000. Six cloned calves produced from adult fibroblast cells after long-term culture. *Proceedings of the National Academy of Sciences of the United States of America* 97(3):990–995 DOI 10.1073/pnas.97.3.990.
- Kwong P, Nam H, Wan Khadijah W, Kamarul T, Abdullah R. 2014. Comparison of in vitro developmental competence of cloned caprine embryos using donor karyoplasts from adult bone marrow mesenchymal stem cells vs. ear fibroblast cells. *Reproduction in Domestic Animals* 49(2):249–253 DOI 10.1111/rda.12262.
- Lazure L, Bachand M, Anseau C, Almeida-Cortez J. 2010. Fate of native and introduced seeds consumed by captive white-lipped and collared peccaries (*Tayassu pecari*, Link 1795 and *Pecari tajacu*, Linnaeus 1758) in the Atlantic rainforest, Brazil. *Brazilian Journal of Biology* 70(1):47–53 DOI 10.1590/S1519-69842010000100008.
- León-Quinto T, Simón MA, Cadenas R, Jones J, Martinez Hernandez FJ, Moreno JM, Soria B. 2009. Developing biological resource banks as a supporting tool for wildlife reproduction and conservation: the Iberian lynx bank as a model for other endangered species. *Animal Reproduction Science* 112(3–4):347–361 DOI 10.1016/j.anireprosci.2008.05.070.
- León-Quinto T, Simón MA, Cadenas R, Martínez Á, Serna A. 2014. Different cryopreservation requirements in foetal vs. adult skin cells from an endangered mammal, the Iberian lynx (*Lynx pardinus*). *Cryobiology* 68(2):227–233 DOI 10.1016/j.cryobiol.2014.02.001.
- Li LF, Guan WJ, Li H, Zhou XZ, Bai XJ, Ma YH. 2009. Establishment and characterization of a fibroblast cell line derived from Texel sheep. *Biochemistry and Cell Biology* 87:485–492 DOI 10.1139/O09-005..
- Liu C, Guo Y, Guan W, Ma Y, Zhang H, Tang X. 2008. Establishment and biological characteristics of Luxi cattle fibroblast bank. *Tissue and Cell* 40(6):417–424 DOI 10.1016/j.tice.2008.04.005.
- Luo J, Liang M, Yang X, Xu H, Shi D, Lu S. 2014. Establishment and biological characteristics comparison of Chinese swamp buffalo (*Bubalus bubalis*) fibroblast cell lines. *In Vitro Cellular & Developmental Biology—Animal* 50:7–15 DOI 10.1007/s11626-013-9677-2.
- Magalhães LC, Bhat MH, Freitas JL, Melo LM, Teixeira DI, Pinto LC, Câmara LM, Duarte JM, Freitas VJ. 2017. The effects of cryopreservation on different passages of fibroblast cell culture in brown brocket deer (*Mazama gouazoubira*). *Biopreservation and Biobanking* 15(5):463–468 DOI 10.1089/bio.2017.0060.
- Magalhães R, Nugraha B, Pervaiz S, Yu H, Kuleshova LL. 2012. Influence of cell culture configuration on the post-cryopreservation viability of primary rat hepatocytes. *Biomaterials* 33(3):829–836 DOI 10.1016/j.biomaterials.2011.10.015.
- Mata M, Garrido-Maraver J, Cotán D, Cordero MD, Oropesa-Ávila M, Izquierdo LG, De Miguel M, Lorite JB, Infante ER, Ybot P. 2012. Recovery of MERRF fibroblasts and cybrids

pathophysiology by Coenzyme Q 10. *Neurotherapeutics* **9**(2):446–463
DOI 10.1007/s13311-012-0103-3.

Mehrabani D, Mahboobi R, Dianatpour M, Zare S, Tamadon A, Hosseini SE. 2014. Establishment, culture, and characterization of Guinea pig fetal fibroblast cell. *Veterinary Medicine International* **2014**(2):1–5 DOI 10.1155/2014/510328.

Queiroz Neta LB, Oliveira Lira GPO, Borges AA, Santos MVO, Silva MB, Oliveira LRM, Silva AR, Oliveira MF, Pereira AF. 2018. Influence of storage time and nutrient medium on recovery of fibroblast-like cells from refrigerated collared peccary (*Pecari tajacu* Linnaeus, 1758) skin. *In Vitro Cellular & Developmental Biology—Animal* **54**:486–495
DOI 10.1007/s11626-018-0270-6.

Redford KH. 1992. The empty forest. *BioScience* **42**(6):412–422 DOI 10.2307/1311860.

Rodriguez-Ororio N, Urrego R, Cibelli J, Eilertsen K, Memili E. 2012. Reprogramming mammalian somatic cells. *Theriogenology* **78**(9):1869–1886
DOI 10.1016/j.theriogenology.2012.05.030.

Roth V. 2006. Doubling time calculator. Available at <http://www.doubling-time.com/compute.php> (accessed 1 September 2019).

Saadeldin IM, Swelum AAA, Noreldin AE, Tukur HA, Abdelazim AM, Abomughaid MM, Alowaimer AN. 2019. Isolation and culture of skin-derived differentiated and stem-like cells obtained from the Arabian Camel (*Camelus dromedarius*). *Animals* **9**(6):378–395
DOI 10.3390/ani9060378.

Santos MLT, Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, Pereira AF. 2016. In vitro culture of somatic cells derived from ear tissue of collared peccary (*Pecari tajacu* Linnaeus, 1758) in medium with different requirements. *Pesquisa Veterinária Brasileira* **36**(12):1194–1202 DOI 10.1590/s0100-736x2016001200010.

Santos DO, Mendes A, Da Nogueira C, Nogueira Filho SLG. 2009. Criação comercial de caítilus (*Pecari tajacu*): uma alternativa para o agronegócio. *Revista Brasileira de Saúde e Produção Animal* **10**:1–10.

Santos MVO, Nascimento LE, Praxedes EA, Borges AA, Silva AR, Bertini LM, Pereira AF. 2019. *Syzygium aromaticum* essential oil supplementation during in vitro bovine oocyte maturation improves parthenogenetic embryonic development. *Theriogenology* **128**:74–80
DOI 10.1016/j.theriogenology.2019.01.031.

Schaffer WI, Terminology Committee Chair Tissue Culture Association. 1990. Terminology associated with cell, tissue and organ culture, molecular biology and molecular genetics. *In Vitro Cellular & Developmental Biology* **26**(1):97–101 DOI 10.1007/BF02624162.

Sharma R, Sharma H, Ahlawat S, Aggarwal R, Vij P, Tandia M. 2018. First attempt on somatic cell cryopreservation of critically endangered *Camelus bactrianus* of India. *Gene Reports* **10**:109–115 DOI 10.1016/j.genrep.2017.11.007.

Shiga K, Fujita T, Hirose K, Sasae Y, Nagai T. 1999. Production of calves by transfer of nuclei from cultured somatic cells obtained from Japanese black bulls. *Theriogenology* **52**(3):527–535
DOI 10.1016/S0093-691X(99)00149-1.

Siengdee P, Klinhom S, Thitaram C, Nganvongpanit K. 2018. Isolation and culture of primary adult skin fibroblasts from the Asian elephant (*Elephas maximus*). *PeerJ* **6**:e4302
DOI 10.7717/peerj.4302.

Silvestre MA, Sánchez JP, Gómez EA. 2004. Vitrification of goat, sheep, and cattle skin samples from whole ear extirpated after death and maintained at different storage times and temperatures. *Cryobiology* **49**(3):221–229 DOI 10.1016/j.cryobiol.2004.08.001.

- Song J, Hua S, Song K, Zhang Y. 2007.** Criação comercial de caititus (*Pecari tajacu*): uma alternativa para o agronegócio. *In Vitro Cellular & Developmental Biology—Animal* **43**:203–209 DOI [10.1007/s11626-007-9043-3](https://doi.org/10.1007/s11626-007-9043-3).
- Trokovic R, Weltner J, Noisa P, Raivio T, Otonkoski T. 2015.** Combined negative effect of donor age and time in culture on the reprogramming efficiency into induced pluripotent stem cells. *Stem Cell Research* **15**(1):254–262 DOI [10.1016/j.scr.2015.06.001](https://doi.org/10.1016/j.scr.2015.06.001).
- Yajing S, Rajput IR, Ying H, Fei Y, Sanganyado E, Ping L, Jingzhen W, Wenhua L. 2018.** Establishment and characterization of pygmy killer whale (*Feresa attenuata*) dermal fibroblast cell line. *PLOS ONE* **13**(3):e0195128 DOI [10.1371/journal.pone.0195128](https://doi.org/10.1371/journal.pone.0195128).

ANEXO C – VERSÃO DO ARTIGO PUBLICADO NA CRYO-LETTERS

**CRYOPRESERVATION OF COLLARED PECCARY (*Pecari tajacu* LINNAEUS,
1758) SOMATIC CELLS IS IMPROVED BY SUCROSE AND HIGH
CONCENTRATIONS OF FETAL BOVINE SERUM**

QUALIS: Quadriênio 2013-2016: B1/ Novo Qualis: A2

FATOR DE IMPACTO: 0,694

CRYOPRESERVATION OF COLLARED PECCARY (*Pecari tajacu* LINNAEUS, 1758) SOMATIC CELLS IS IMPROVED BY SUCROSE AND HIGH CONCENTRATIONS OF FETAL BOVINE SERUM

Gabriela Pereira de Oliveira Lira¹, Alana Azevedo Borges¹, Matheus Barbosa do Nascimento¹, Leonardo Vitorino Costa de Aquino¹, Moacir Franco de Oliveira², Alexandre Rodrigues Silva³ and Alessandra Fernandes Pereira^{1*}

¹Laboratory of Animal Biotechnology, Federal Rural University of Semi-Arid (UFERSA), Mossoro, RN, Brazil.

²Laboratory of Applied Animal Morphophysiology, UFERSA, Mossoro, RN, Brazil.

³Laboratory of Animal Germplasm Conservation, UFERSA, Mossoro, RN, Brazil

*Corresponding author e-mail: alexandra.pereira@ufersa.edu.br

Abstract

BACKGROUND: The formation of somatic cell banks is affected by, amongst other factors, the cryoprotectant solution used. The selection of an effective solution, therefore, is a primary parameter. **OBJECTIVE:** We optimized the cryoprotectant used for collared peccary somatic cell cryopreservation. **MATERIALS AND METHODS:** We categorized cells into different groups based on their cryopreservation and evaluated the morphology, viability, proliferative activity, metabolism, and oxidative stress. One group was cryopreserved in 10% DMSO with 10% fetal bovine serum (DMSO-10FBS), and another with 50% FBS (DMSO-50FBS). The cryopreservation of both groups included the presence of 0.2 M sucrose (DMSO-SUC-10FBS and DMSO-SUC-50FBS). Non-cryopreserved cells and cells cryopreserved with 10% DMSO (DMSO) supplemented with 0.2 M sucrose (DMSO-SUC) were used as controls. **RESULTS:** There was no difference observed in morphology or viability among the groups. Proliferative activity was reduced in DMSO-10FBS when compared to controls. Although cryopreservation reduced metabolism, no difference was observed among solutions. A lower level of reactive oxygen species was observed in cells of DMSO-SUC-50FBS when compared to other cryoprotectants. Only cells of DMSO-SUC-50FBS had mitochondrial potential similar to non-cryopreserved cells. **CONCLUSION:** 10% DMSO supplemented with 50% FBS and 0.2 M SUC was observed to be the most efficient cryoprotectant for preserving collared peccary somatic cells.

Keywords: Peccaries, extracellular cryoprotectants, slow freezing, cryobanking.

INTRODUCTION

Collared peccaries (*Pecari tajacu* Linnaeus, 1758), also known as “wild pigs”, are wild mammals belonging to the Tayassuidae family of the order Artiodactyla (21). These animals play an important role in maintaining the global ecosystem where they act as seed dispersers, promote flora renewal, and contribute to the

food chain (34). Although globally classified as “Least Concern”, this species is already extinct in eastern and southern Argentina and is declining in some biomes (9), thus necessitating conservation strategies for the maintenance of its population.

One of the conservation tools applied to wild mammals has been the formation of somatic resource banks (6). These banks allow

long-term storage of tissues and somatic cells to be employed in different proposals, such as multiplying individuals by somatic cell nuclear transfer (SCNT) (43) and generating induced pluripotent cells (41). The efficiency of these procedures depends on the quality of cells after slow freezing, which is dependent on the choice of the cryoprotectant, a crucial step for the success of these cryobanks (28). In general, an effective cryoprotectant is the appropriate combination of intracellular cryoprotectants such as dimethyl sulfoxide (DMSO) (25), ethylene glycol (EG) (16), and extracellular cryoprotectants such as sucrose (SUC) (23) and/or fetal bovine serum (FBS) (15).

Specifically, our group has successfully established somatic tissue banks of collared peccaries based on our knowledge of the tissues (2), techniques (3), and solutions required (4, 5). Subsequently, we initiated somatic cell banks of this species, demonstrating that 10% DMSO in the presence of 0.2 M SUC was more efficient in somatic cell slow freezing when compared to EG with 0.2 M SUC, showing a viability of 69.8% and 58.5%, respectively (17). Thus, one way to improve the rates of cell recovery would be to optimize the extracellular agents employed. Studies have shown that FBS, when associated with 10% DMSO and 0.2 M SUC, promoted a beneficial effect in some species (*Lynx pardinus* and *Elephas maximus*) (15, 36). This agent can reduce oxidative stress, caused by excessive production of reactive oxygen species (ROS) during slow freezing (11).

Nevertheless, variations in FBS concentration can be observed in some species with its use at 10% (*Felis silvestris libica*) (8) and 50% (*Bubalus bubalis*) (18), necessitating the evaluation of the most appropriate concentration, as well as its relationship with SUC. Therefore, we aimed to optimize the cryoprotectant by varying the different extracellular combinations of SUC and FBS. We assessed the impact of the cryoprotectants on the morphology, viability, proliferative activity, metabolism, and oxidative stress of cryopreserved collared peccary cells.

MATERIALS AND METHODS

Chemicals and media

All chemicals used were purchased from Sigma-Aldrich (St. Louis, MO, USA). Dulbecco's modified Eagle's medium (DMEM), FBS, trypsin-EDTA, Antibiotic-Antimycotic

(ATB-ATM) solution and fluorescent probes were obtained from Gibco-BRL (Carlsbad, CA, USA). The 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) assay was purchased from Grainer (Miami, FL, USA).

Compliance with ethical standards and animals

The experimental protocols and animal handling procedures were performed with the approval of the Animal Ethics Committee of the Federal Rural University of Semi-Arid (CEUA/UFERSA, no. 23091.001072/2015-92), in compliance with the Chico Mendes Institute for Biodiversity Conservation (ICMBio, no. 48633-2). A total of four adult collared peccaries, 41.5 ± 1.8 months old, provided by the Centre of Multiplication of Wild Animals (CEMAS/UFERSA, Mossoró, RN, Brazil; $5^{\circ}10'S$, $37^{\circ}10'W$) were used.

Skin biopsy, primary culture, and subcultures

Sections (1-2 cm²) used for collared peccary identification were taken from peripheral ear tissue using pliers. After collection, tissue samples were washed in 70% ethanol (38) and incubated in DMEM supplemented with 10% FBS and 2% ATB-ATM solution at 37 °C for 30 min.

In the laboratory, tissues were trichotomized, washed in 70% ethanol followed by DMEM supplemented with 10% FBS and 2% ATB-ATM solution. Subsequently, tissue samples were fragmented in 9.0 mm³ sections and cultured in DMEM supplemented with 10% FBS and 2% ATB-ATM solution at 38.5 °C in an incubator with 5% CO₂ and 95% air (31). During the primary culture, the medium was changed every 24 h. Cells were subcultured after reaching 70-80% confluence until the third passage, followed by cryopreservation. The subconfluence of 70-80% was defined when 70-80% of the Petri dishes presented somatic cells (31).

Study design and cell cryopreservation

The morphology, viability, proliferative activity, metabolism, and oxidative stress were evaluated in non-cryopreserved and cryopreserved cells. Cells were cryopreserved in either 10% DMSO with 10% FBS (DMSO-10FBS group), or 50% FBS (DMSO-50FBS group). In both the groups, 0.2 M SUC was added and these were further designated as the DMSO-SUC-10FBS and DMSO-SUC-50FBS groups, respectively. Moreover, non-

cryopreserved cells (non-cryopreserved group), those cryopreserved in 10% DMSO (DMSO group) and those cryopreserved with 10% DMSO supplemented with 0.2 M SUC (DMSO-SUC group), were used as controls.

For cryopreservation, cells were subjected to slow cooling in freezing medium (DMEM supplemented with 10% DMSO and extracellular cryoprotectants according to experimental groups). Briefly, cells at a concentration of 1.0×10^4 cells/mL were first exposed to DMSO solution for 15 min at 4 °C in DMEM, followed by the addition of SUC and FBS solutions according to experimental groups and incubated for an additional 15 min at 4 °C. The cryovials containing 1.0 mL of cells in freezing solution were cooled in a Mr. Frosty freezing container (Thermo Scientific Nalgene, Rochester, NY, USA) at a cooling rate of 1 °C/min, and kept in a freezer -80 °C until reaching -70 °C, before being transferred to liquid nitrogen (15).

After 2 weeks, the cryovials were removed from liquid nitrogen and kept at room temperature of 25 °C for up to 1 min and then warmed in a water bath at 37 °C for 3-4 min. For the removal of cryoprotectants, the cells were washed twice with DMEM. Cell suspension was centrifuged for 10 min at 400×g and washed again with DMEM and centrifuged according to León-Quinto et al. (15). In the groups containing SUC, the cells were washed with DMEM with 0.2 M SUC, prior to centrifugation as previously described.

Evaluation of morphological characteristics and cell viability

The cells' morphology was evaluated daily using an inverted microscope (Nikon TS100, Tokyo, Japan). The following morphological characteristics were assessed: size, aspect, shape and adhesion (29). The viability analysis was performed by cells stained with trypan blue. For each animal and each group, the viability test was performed in duplicate. Briefly, the cells were centrifuged and suspended in 1.0 mL of the cell culture medium (DMEM supplemented with 10% FBS and 2% ATB-ATM solution); an

aliquot of cells was stained with 0.4% trypan blue (in phosphate buffered saline, PBS) in the ratio 1:1 and counted in a Neubauer chamber (39).

Analysis of proliferative activity and metabolism

Proliferative activity was quantified according to the population doubling time (PDT), which was calculated by seeding cells at a density of 1.0×10^4 cells/well in a 24-well plate. Cells were trypsinized in duplicate and counted at 24 h intervals for up to 216 h of culture. The mean cell counts were recorded every time and the cell growth curve was delineated. Finally, the PDT was calculated using the following formula (30):

$$\text{PDT} = T \ln 2 / \ln (X_e / X_b),$$

where PDT is the time of the culture (in hours), T is the incubation time, X_b is the number of cells at the beginning of the time incubation, X_e is the number of cells at the end of the incubation time, and $\ln$ is Napierian logarithm.

For evaluation of metabolism, cells were seeded into 12-well dishes at a density of 5.0×10^4 cells/mL and cultured at 38.5 °C in a humid atmosphere with 5% CO₂ (4). After five days, MTT solution (5 mg/mL in DMEM) was added to each well for 3 h under the same conditions. After incubation, the MTT solution was removed, and DMSO was added to each dish to solubilize the formazan crystals. Finally, the absorbance (595 nm) of the samples was read using a spectrophotometer (Shimadzu® UV-mini-1240, Kyoto, Japan). The mean values of non-cryopreserved cells were considered 100% as compared to the other groups.

Assessment of oxidative stress

Cellular oxidative stress was evaluated by quantification of intracellular reactive oxygen species (ROS) levels and measurement of mitochondrial membrane potential ($\Delta\Psi_m$). Briefly, intracellular levels of ROS were quantified using 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA) as a fluorescent probe, according to Santos et al. (32).




Figure 1. Subcultures of fibroblast-like cells from ear skin samples of collared peccaries. **a)** Non-cryopreserved cells and cell cryopreserved in **b)** DMSO, **c)** DMSO-SUC, **d)** DMSO-10FBS, **e)** DMSO-50FBS, **f)** DMSO-SUC-10FBS, **g)** DMSO-SUC-50FBS. Scale bar = 100 μm .

For this, the cells, independent of the experimental group, were seeded in glass slides and grown *in vitro* until reaching 70% confluence. Subsequently, cells were washed in PBS and incubated with 1.0 mL 5 μM H_2DCFDA at 38.5 $^{\circ}\text{C}$ with 5% CO_2 for 30 min protected from light. After this period, the stained cells were washed twice in PBS and imaged using a fluorescence microscope (Olympus BX51TF, Tokyo, Japan). The intensity of the signal fluorescence (pixels) was measured using ImageJ software (version 1.49v, Java 1.8.0_201, Wayne Rasband, U.S. National Institutes of Health, Bethesda, MD, USA; website: <http://rsb.info.nih.gov/ij/download.html>). The

background signal intensity was subtracted from the values obtained for the treated samples. The measured mean value of the micrograph for non-cryopreserved cells was used as a calibrator. Relative expression levels (arbitrary fluorescence units) were generated by dividing the measured value of each micrograph for the cryopreserved cells by the mean of the calibrator.

The $\Delta\Psi\text{m}$ was assessed using the fluorescent probe MitoTracker Red[®] (CMXRos) at 500 nm according to Santos et al. (32). The procedure, incubation and evaluation of the images, were performed as described for the quantification of ROS.




Figure 2. Viability of collared peccary non-cryopreserved and cryopreserved cells using different extracellular cryoprotectants. Non-cryopreserved group, 10% DMSO (DMSO group), 10% DMSO and 0.2 M SUC (DMSO-SUC group), 10% DMSO and 10% FBS (DMSO-10FBS group), 10% DMSO and 50% FBS (DMSO-50FBS group), 10% DMSO, 0.2 M SUC and 10% FBS (DMSO-SUC-10FBS group), 10% DMSO, 0.2 M SUC and 50% FBS (DMSO-SUC-10FBS group). $P > 0.05$.

Statistical analysis

Data were expressed as mean $\pm$ standard error (one animal/one repetition) and analyzed using the GraphPad software (Graph-Pad Software Incorporated, La Jolla, CA, USA). All results were verified for normality by the Shapiro-Wilk test and homoscedasticity by Levene's test. Since data did not show a normal distribution, they were arcsine transformed and analyzed by ANOVA followed by Tukey test. Significance was set at $P < 0.05$.

RESULTS

Evaluation of morphological characteristics and cell viability

All cells derived from collared peccary tissue had consistent morphological characteristics in all experimental groups (Fig. 1a-g). Slow cooling did not alter the capacity of the cells to grow and reach 70-80% confluence,

presenting cells with similar morphology to fibroblasts; with fusiform aspect, cytoplasmic prolongations, abundant cytoplasm and central nucleus, and with an overall large and oval morphology.

Moreover, no difference was observed among groups for cell viability evaluated by trypan blue (Fig. 2, $P > 0.05$). Thus, while non-cryopreserved cells had a viability of $87.0 \pm 5.2\%$, cryopreserved cells of different groups showed viabilities in the range of $62.5 \pm 5.2\%$ to $91.7 \pm 0.0\%$.

Analysis of proliferative activity and metabolism

Although the growth curve did not show a typical "S" shape in all cryopreserved groups, it was possible to observe from the shape of the curve the occurrence of the initial adaptation phase of the cells, followed by exponential growth until reaching the stable phase of decline (Fig. 3a). In addition, non-cryopreserved cells




Figure 3. The growth dynamics and metabolism of collared peccary non-cryopreserved and cryopreserved cells using different extracellular cryoprotectants. **a)** Growth curves of cryopreserved and non-cryopreserved cells. **b)** Values of population doubling time (PDT) after culture for nine days. Non-cryopreserved group, 10% DMSO (DMSO group), 10% DMSO and 0.2 M SUC (DMSO-SUC group), 10% DMSO and 10% FBS (DMSO-10FBS group), 10% DMSO and 50% FBS (DMSO-50FBS group), 10% DMSO, 0.2 M SUC and 10% FBS (DMSO-SUC-10FBS group), 10% DMSO, 0.2 M SUC and 50% FBS (DMSO-SUC-10FBS group). Different letters show significant differences among the groups ($P < 0.05$). Bars indicate standard error.

showed a reduction in their growth; however, this reduction stabilized from day 8. Thereafter the decline in the curve was no longer observed, which showed that there was no cell death only the decrease of proliferative activity of cells from this group. Additionally, the proliferative activity evaluated by population doubling time was reduced in cells derived from DMSO-10FBS group ($10.5 \text{ h} \pm 0.3$) when compared to non-cryopreserved cells (19.4 ± 2.6 , Fig. 3b)

Although cryopreservation reduced cell metabolism, no difference was observed among cryoprotectant solutions ($P > 0.05$). Thus, while cryopreserved cells had a metabolic activity of $100.0 \pm 0.0\%$, cryopreserved cells among the different groups varied among $63.6 \pm 22.6\%$ and $97.7 \pm 25.1\%$.

Assessment of oxidative stress

A lower level of ROS was observed in cells derived from DMSO-SUC-50FBS when compared to other cryopreserved groups (Fig 4a and 4c).

Moreover, only cells from the DMSO-SUC-50FBS group had mitochondrial membrane potential similar to the non-cryopreserved cells (Fig 4b and 4d).

DISCUSSION

The successful formation of a bank of biological resources is correlated to the efficiency of the cryopreservation protocol used. As such, the initial combination of intracellular and extracellular cryoprotectants is key in ensuring cell quality after thawing. In this study, we showed that the combination of 0.2 M SUC with 50% FBS added to the cryoprotectant solution containing 10% DMSO, ensured a better quality of the recovered collared peccary somatic cells, particularly by reducing the oxidative stress caused by slow freezing and by maintaining normal cellular characteristics. Therefore, our results demonstrated three relevant points of cryopreservation of somatic cells: (a) the cells obtained from collared peccary tissue showed characteristics similar to fibroblast cells used as the main donor nucleus in SCNT; (b) some alterations resulting from the cryopreservation may not be observed from the analysis of the viability and metabolic activity of the thawed cells; therefore, more specific assessments such as the analysis of the oxidative stress may be necessary; and (c) the favorable effect of the addition of 50% FBS on the cryoprotectant solution containing 10% DMSO and 0.2 M SUC. Thus, these findings ensure that




Figure 4. Assessment of oxidative stress in collared peccary non-cryopreserved and cryopreserved cells using different extracellular cryoprotectants. Non-cryopreserved group, 10% DMSO (DMSO group), 10% DMSO and 0.2 M SUC (DMSO-SUC group), 10% DMSO and 10% FBS (DMSO-10FBS group), 10% DMSO and 50% FBS (DMSO-50FBS group), 10% DMSO, 0.2 M SUC and 10% FBS (DMSO-SUC-10FBS group), 10% DMSO, 0.2 M SUC and 50% FBS (DMSO-SUC-50FBS group). **a)** Quantification of intracellular reactive oxygen species (ROS) levels **b)** Measurement of mitochondrial membrane potential ($\Delta\Psi_m$). **c)** Cells labeled with H_2DCFDA for quantification of ROS levels. **d)** Cells labeled with MitoTracker Red[®] for measurement of $\Delta\Psi_m$. Different letters within each analysis show significant differences among the groups ($P < 0.05$). Scale bar = 200 μm .

higher quality cells are employed in future conservation strategies.

In general, cryopreservation protocols can induce lesions such as lipid peroxidation of membranes and damage to DNA in fibroblasts, resulting from the formation of ROS (33). Moreover, the natural antioxidant defense mechanism of the cells is not enough to eliminate excess ROS (11). Therefore, studies have shown that supplementation with antioxidants and/or factors that modulate the cell death process caused by excess ROS may be a potential solution to reduce oxidative stress induced by cryopreservation (14). Additionally, a decrease in $\Delta\Psi_m$ has been observed in cells with excess ROS formation (13). Thus, as suggested by Poot et al. (27), we used a fluorescent probe that has an affinity for molecules present in cells. Such probes are responsible for evaluating the relationship among mitochondria and the triggering of apoptotic mechanisms. With this, it was possible to demonstrate that among the cryoprotectant solutions tested, the cells obtained from the DMSO-SUC-50FBS treatment showed high $\Delta\Psi_m$ after thawing.

Thus, one compound that has such antioxidant potential is FBS, which has been frequently used in cryoprotectant solutions (12). In general, FBS is routinely added to stabilize cell membranes and adjust intra- and extracellular osmotic pressure. However, the role of this cryoprotectant as an antioxidant has become increasingly evident, which may be related to its constituents, mostly amino acids (20). This could be due to the synergistic effect of their constituents, which may have attached to cell membranes, promoting stabilization and avoiding excessive concentration of solutes that may be related to ROS formation (12). In addition, their constituents may be responsible for the increased activity of antioxidant intracellular enzymes (26). Thus, it is likely that the FBS in the DMSO-SUC-FBS solution acted as a source of antioxidants. However, the mechanisms underlying the antioxidant role of FBS under the formation of ROS suffered by cells during cryopreservation are still unclear. Moreover, when the cells were cryopreserved with 10% FBS (DMSO-10FBS and DMSO-SUC-10FBS groups), the fluorescence intensity increased, indicating an increase in ROS production. This suggests that the reduction and/or absence of antioxidative agents in the DMSO, DMSO-10FBS and DMSO-SUC-10FBS

groups may be associated with the formation of ROS (24, 42, 44).

This result may be due to FBS counteracting the cytotoxicity induced by DMSO, as demonstrated by Miki et al. (22), when analyzing the $\Delta\Psi_m$ of stem cells subjected to a cryoprotectant solution with 10% DMSO in association with 90% FBS, which resulted in a $\Delta\Psi_m$ of 1.00. Therefore, treatments with 10% FBS caused the cells to be more exposed to cytotoxicity caused by DMSO during the freeze-thaw cycle. The DMSO penetrates the cell membrane and forms hydrogen bonds with the water molecules present in the cytosol. Thus, the freezing point of the water decreases, and there is a lower probability of the formation of ice crystals inside the cell. Furthermore, the integrity of the membrane is conserved and for this reason, DMSO is an indispensable intracellular cryoprotectant. However, even though its positive effect is observed, its toxicity remains high, suggesting its association with extracellular cryoprotectants such as SUC and FBS. This association can accelerate the cellular dehydration process and balance cytotoxicity. Therefore, the ideal concentrations for achieving this cytotoxic balance must be established so that the concentrations of extracellular cryoprotectants are sufficient to minimize the negative effect of the high toxicity of DMSO. Additionally, the presence of SUC can promote osmotic maintenance of the medium and perform synergistically with the intracellular cryoprotectant, resulting in improved effects when used in combination (37).

The ROS levels of the DMSO-SUC-10FBS group were higher in comparison to the DMSO group, probably due to the use of more cryoprotective agents in their composition. It should be borne in mind that such cryoprotective agents (DMSO-SUC-10FBS), despite promoting cell protection, present a certain toxicity that may be responsible for greater oxidative stress and subsequent increased production of ROS. However, this does not mean that the use of fewer cryoprotective agents is recommended, but rather that a combination promoting balance between the concentration used, toxicity, protection of cells at cryogenic temperatures and lower production of ROS, should be utilized.

In addition, the morphological and confluence aspects observed in the collared cells were similar to those observed by Queiroz Neta et al. (29) and Borges et al. (4) describing collared peccary cells recovered from

refrigerated and vitrified tissues, respectively. Thus, proper maintenance of the morphological characteristics of cultured cells may be associated with a decrease in ice crystal formation due to the cryopreservation process, which when exposed to cells, may have deleterious effects on cell characteristics (15).

The viability data showed that cryopreserved cells and non-cryopreserved cells were stable regardless of treatments. This can be attributed to the quality of the 10% DMSO slow freezing technique that is commonly associated with better results for somatic cells derived from wild mammals (23, 35).

Further, the higher viability rates and the in vitro culture performance of cells shown in this study, corroborate the findings of Subramanian et al. (40) and Barnes et al. (1). These authors affirmed that cultures derived from cryopreserved and non-cryopreserved cells have a similar growth performance, since the cell population after culture can recover with rates similar to non-cryopreserved cells.

Finally, all groups showed no change in PDT values except DMSO-10FBS. In general, cells cultured in specific treatments including FBS in their composition required less time to replicate (19). Our PDT rates and metabolism activity evaluated by MTT may be associated with the use of an adequate system of cryopreservation that promoted an optimum cooling rate (10) associated with compounds present in FBS such as transport proteins, adhesion proteins, enzymes, hormones, growth factors, cytokines, fatty acids, lipids, vitamins, carbohydrates, and nitrogen of non-protein origin (7).

In conclusion, 10% DMSO supplemented with 0.2 M SUC and 50% FBS was the most efficient cryoprotectant solution for conserving collared peccary somatic cells. This work presents a breakthrough for the improvement and establishment of an ideal freezing protocol for collared peccary somatic cells that can be used in future biotechnical techniques such as cloning, since each stage involved should be optimized for the species studied.

Acknowledgements: The authors thank the Centre for Wild Animals Multiplication (CEMAS/UFERSA) for providing the animals. This study was financed in part by the Higher Education Personnel Improvement Coordination - Brazil (CAPES, Financial Code 001) and National Counsel of Technological and

Scientific Development (CNPq). MF Oliveira, AR Silva and AF Pereira were CNPq investigators.

REFERENCES

1. Barnes LM, Bentley CM & Dickson AJ (2003) *Biotechnol Prog* **19**, 233–237.
2. Borges AA, Bezerra FVF, Costa FN, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR & Pereira AF (2017) *Arq Bras Med Vet Zootec* **69**, 948–954.
3. Borges AA, Lima GL, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR & Pereira AF (2017) *Cytotechnology* **69**, 643–654.
4. Borges AA, Lira GPO, Nascimento LE, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR & Pereira AF (2018) *Biopreserv Biobank* **16**, 77–81.
5. Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR & Pereira AF (2018) *Pesq Vet Bras* **38**, 350–356.
6. Borges AA & Pereira AF (2019) *Zygote* **27**, 111–117.
7. Brunner D, Frank J, Appl H, Schöffl H, Pfaller W & Gstraunthaler G (2010) *Altex* **27**, 53–62.
8. Gómez MC, Jenkins JA, Giraldo A, Harris RF, King A, Dresser AL & Pope CE (2003) *Biol Reprod* **69**, 1032–1041.
9. Gongora J, Biondo C, Cooper JD, Taber A, Keuroghlian A, Altrichter M, Nascimento FF, Chong AY, Miyaki CY, Bodmer R, Mayor P & González S (2011) *Bonn Zool Bull* **60**, 95–101.
10. Gurruchaga H, Ciriza J, del Burgo LS, Rodriguez-Madoz JR, Santos E, Prosper F, Hernández RM, Orive G & Pedraz JL (2015) *Int J Pharm* **485**, 15–24.
11. Gurruchaga H, del Burgo LS, Hernández RM, Orive G, Selden C, Fuller B, Ciriza J & Pedraz JL (2018) *J Control Release* **281**, 119–138.
12. Kimura Y, Okitsu T, Xibao L, Teramae H, Okonogi A, Toyoda K, Uemoto S & Fukushima M (2013) *Islets* **5**, 45–52.
13. Lançoni R, Celeghini ECC, Alves MBR, Lemes KM, Gonella-Díaz AM, Oliveira LZ & Arruda AR (2018) *J Equine Vet Sci* **69**, 78–83.
14. Len JS, Koh WSD & Tan WST (2019) *Biosci Rep* **39**, BSR20191601.
15. León-Quinto T, Simón MA, Cadenas R, Martínez A & Serna A (2014) *Cryobiology* **68**, 227–233.

16. Li Y, Lu RH, Luo GF, Pang WJ & Yang GS (2006) *Cryobiology* **53**, 240–247.
17. Lira GPO, Borges AA, Nascimento LE, Santos MVO, Queiroz Neta LB, Oliveira MF & Pereira AF (2017) in *Anais do Encontro de Biotecnologia do Nordeste* (RENORBIO).
18. Luo C, Lu C, Wang X, Wang Z, Li X, Gong F, Jiang J, Liu Q & Shi D (2013) *Theriogenology* **80**, 878–886.
19. Mahesh YU, Rao BS, Katari VC, Komjeti S, Christo D, Lakshmikantan U, Pawar RM & Shivaji S (2012) *Reprod Domest Anim* **47**, 799–805.
20. Marco-Jiménez F, Garzón DL, Peñaranda DS, Pérez L, Viudes-de-Castro MP, Vicente JS, Jover M & Asturiano JF (2006) *Cryobiology* **53**, 51–57.
21. Mayer JJ & Wetzel RM (1987) *Mamm. Species* **293**, 1–7.
22. Miki T, Wong W, Zhou E, Gonzalez A, Garcia I & Grubbs BH (2016) *Stem Cell Res Ther* **7**, 8.
23. Moulavi F, Hosseini SM, Tanhaie-Vash N, Ostadhosseini S, Hosseini SH, Hajinasrollah M, Asghari MH, Gourabi H, Shahverdi A, Vosough AH & Nasr-Esfahani MH (2017) *Theriogenology* **90**, 197–203.
24. Nirmala JG, Akila S, RT Narendhirakannan & Chatterjee S (2017) *Adv Powder Technol* **28**, 1170–1184.
25. Oliveira LRM, Silva MB, Praxedes EA, Ribeiro LR, Silva HVR & Pereira AF (2020) *An Acad Bras Ciênc* In press.
26. Piao MJ, Hyun YJ, Cho SK, Kang HK, Yoo ES, Koh YS, Lee NH, Ko MH & Hyun JW (2012) *Mar Drugs* **10**, 2826–2845.
27. Poot M, Gibson LL & Singer VL (1997) *Cytometry A* **27**, 358–364.
28. Praxedes EA, Borges AA, Santos MVO & Pereira AF (2018) *Zoo Biol* **37**, 258–263.
29. Queiroz Neta LB, Lira GPO, Borges AA, Santos MVO, Silva MB, Oliveira LRM, Silva AR, Oliveira MF & Pereira AF (2018) *In Vitro Cell Dev Biol Anim* **54**, 486–495.
30. Roth V (2006) <http://www.doubling-time.com/compute.php>, (2006), Accessed date: 12 February 2019.
31. Santos MLT, Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR & Pereira AF (2016) *Pesq Vet Bras* **36**, 194–1202.
32. Santos MVO, Nascimento LE, Praxedes EA, Borges AA, Silva AR, Bertini LM & Pereira AF (2019) *Theriogenology* **128**, 74–80.
33. Santos TRM, Silva ACG, Carvalho FS, Sanz G, Vaz BG, Lião LM, Menegatti R & Valadares MC (2018) *Chem-Biol Interact* **293**, 20–27.
34. Schlichting PE (2018) *J. Wildl. Manage* **82**, 1556–1557.
35. Sharma R, Sharma H, Ahlawat S, Aggarwal RAK, Vij PK & Tania MS (2018) *Gene Rep* **10**, 109–115.
36. Siengdee P, Klinhom S, Thitaram C & Nganvongpanit C (2018) *PeerJ* **6**, e4302.
37. Silva AR, Campos LB, Maia KM & Borges AA (2017) *Rev Bras Reprod Anim* **41**, 110–115.
38. Singh M & Degala V (2017) *Int J Biol* **9**, 9–13.
39. Strober W (2015) *Curr Protoc Immunol* **111**, A3-B.
40. Subramanian J, Aulakh RPS, Grewal PS, Sanford M, Pynn AFJ & Yuk IH (2018) *Biotechnol Prog* **34**, 463–477.
41. Verma R, Holland MK, Temple-Smith P & Verma PJ (2012) *Theriogenology* **77**, 220–228.
42. von Bomhard A, Elsässer A, Ritschl LM, Schwarz S & Rotter N (2016) *PloS One* **11**, e0149660.
43. Wani NA, Vettical BS & Hong SB (2017) *PloS One* **12**, e0177800.
44. Zhang L, Xue X, Yan J, Yan LY, Jin HH, Zhu XH, He ZZ, Liu J, Li R & Qiao J (2016) *Sci Rep* **6**, 26326.

**ANEXO D – VERSÃO DO ARTIGO PUBLICADO NA PESQUISA VETERINÁRIA
BRASILEIRA**

***IN VITRO* MATURATION OF COLLARED PECCARY (*Pecari tajacu* LINNAEUS,
1758) OOCYTES AFTER DIFFERENT INCUBATION TIMES**

QUALIS: Qualis 2013-2016: A2/ Novo Qualis: A4

FATOR DE IMPACTO: 0,1474

DOI: <https://doi.org/10.1590/1678-5150-pvb-5471>

***In vitro* maturation of collared peccary (*Pecari tajacu* Linnaeus, 1758) oocytes after different incubation times¹**

Alana A. Borges², Maria V.O. Santos², Luiza B. Queiroz Neta², Moacir F. Oliveira³,
Alexandre R. Silva⁴ and Alexsandra F. Pereira^{2*}

ABSTRACT.- Borges A.A., Santos M.V.O., Queiroz Neta L.B., Oliveira M.F., Silva A.R. & Pereira A.F. 2018. [*In vitro* maturation of collared peccary (*Pecari tajacu* Linnaeus, 1758) oocytes after different incubation times.] *Maturação in vitro de oócitos de cateto (Pecari tajacu Linnaeus, 1758) após diferentes períodos de incubação.* *Pesquisa Veterinária Brasileira* 38(9):. Universidade Federal Rural do Semi-Árido, Av. Francisco Mota 572, Mossoró, RN 59625-900, Brazil. E-mail: alexsandra.pereira@ufersa.edu.br

Oocyte *in vitro* maturation (IVM) is the first step of the *in vitro* reproductive technologies that enables mature oocytes to be generated *ex vivo* and after used for embryo production. In this sense, the establishment of culture environment, as oocyte incubation time, is essential for the success of the IVM. Therefore, the study was carried out to investigate the relationship between the meiotic potential and the IVM times of collared peccary oocytes, wild mammals of great commercial and ecological interest. Thus, ovaries were collected of females derived from captivity and transported to the laboratory within 1 hour of slaughtering. The oocytes derived from follicles (3-6mm in diameter) were recovered by aspirated and sliced. Good quality oocytes (evenly granulated cytoplasm with a least one layer of surrounding *cumulus* cells) were selected and subjected to culture in TCM 199 supplemented with 10µg/mL FSH, 10% FBS and 100µM cysteamine at 38.5°C, 5% CO₂ and maximum humidity for 24 or 48 hours. After the incubation period, the nuclear status, the presence of first polar body and the expansion of *cumulus* cells of oocytes were assessed. The data obtained were analyzed by Fisher exact test (P<0.05). A total of four sessions (2-3 females per session) were performed, resulting in eighteen aspirated and sliced ovaries with normal morphological characteristics. An oocyte recovery rate of about 83.1% (59/71) was obtained with 3.3 oocytes/ovary and 2.3 viable oocytes/ovary. After different incubation times, differences (P<0.05) were observed in 24 and 48 hours for expansion of the *cumulus* cells (38.1% vs. 100%), presence of first polar body (52.4% vs. 90.5%) and nuclear status in second metaphase (19.0% vs. 76.2%), respectively. In conclusion, 48 hours is suitable time for the *in vitro* maturation of oocytes derived from collared peccaries when compared to the time of 24 hours, according to the meiotic potential observed. Additional studies should be conducted to improve the quality of the oocyte culture environment, as medium composition, aiming to obtain viable mature oocytes for other *in vitro* biotechnologies.

INDEX TERMS: Collared peccary, *Pecari tajacu*, oocytes, incubation, reproduction, meiotic competence, nuclear maturation, wild mammals.

¹ Received on July 31, 2017.

Accepted for publication on October 18, 2017.

² Laboratório de Biotecnologia Animal, Universidade Federal Rural do Semi-Árido (UFERSA), Av. Francisco Mota 572, Mossoró, RN 59625-900, Brazil. *Corresponding author: alexsandra.pereira@ufersa.edu.br

³ Laboratório de Morfofisiologia Animal Aplicada, UFERSA, Av. Francisco Mota 572, Mossoró, RN 59625-900.

⁴ Laboratório de Conservação de Germoplasma Animal, UFERSA, Av. Francisco Mota 572, Mossoró, RN 59625-900.

RESUMO.- [*Maturação in vitro de oócitos de cateto (Pecari tajacu Linnaeus, 1758) após diferentes períodos de incubação.*] A maturação *in vitro* (MIV) oocitária é a primeira etapa das tecnologias reprodutivas *in vitro* que permite que oócitos maturados sejam gerados *ex vivo* e depois usados para a produção de embriões. Nesse sentido, o estabelecimento do ambiente de cultivo, como o período de incubação de oócitos, é essencial para o sucesso da MIV.

Portanto, o estudo foi realizado para investigar a relação entre o potencial meiótico e os períodos de MIV de oócitos derivados de catetos, mamíferos selvagens de grande interesse comercial e ecológico. Para tanto, os ovários foram coletados de fêmeas derivadas de cativeiro e transportados ao laboratório dentro de 1 h após o abate. Os oócitos derivados de folículos (3-6mm de diâmetro) foram recuperados por aspiração e fatiados. Oócitos de boa qualidade (citoplasma uniformemente granuloso com pelo menos uma camada circundante de células *cumulus*) foram selecionados e submetidos ao cultivo em TCM 199 suplementado com 10µg/mL de FSH, 10% de SFB e 100µM de cisteamina a 38,5°C, 5% de CO₂ e umidade máxima por 24 e 48 h. Após o período de incubação, o estado nuclear, a presença do primeiro corpúsculo polar e a expansão das células do *cumulus* dos oócitos foi avaliada. Os dados obtidos foram analisados pelo teste exato de Fisher ($P < 0,05$). Um total de quatro sessões (2-3 fêmeas por sessão) foi realizado, resultando em dezoito ovários aspirados e fatiados com características morfológicas normais. Uma taxa de recuperação oocitária de aproximadamente 83,1% (59/71) foi obtida com 3,3 oócitos/ovário e 2,3 oócitos viáveis/ovário. Após diferentes períodos de incubação, diferenças ($P < 0,05$) foram observadas entre 24 e 48 h para a expansão das células *cumulus* (38,1% vs. 100%), presença de primeiro corpúsculo polar (52,4% vs. 90,5%) e estado nuclear na segunda metáfase (19,0% vs. 76,2%), respectivamente. Em conclusão, 48 h é o período adequado para a maturação *in vitro* de oócitos derivados de catetos quando comparado ao tempo de 24 h, de acordo com o potencial meiótico observado. Estudos adicionais devem ser conduzidos para melhorar a qualidade do ambiente de cultivo oocitário, como a composição de meio, objetivando obter oócitos maturados viáveis para outras biotecnologias *in vitro*.

TERMOS DE INDEXAÇÃO: Maturação, oócitos, cateto, *Pecari tajacu*, incubação, reprodução, competência meiótica, maturação nuclear, mamíferos silvestres.

INTRODUCTION

Oocyte *in vitro* maturation (IVM) is the first step of the *in vitro* reproductive technologies that enables mature oocytes to be generated *ex vivo* (Rahman et al. 2008) and further destined to systems of *in vitro* embryo production (IVEP), as *in vitro* fertilization (IVF), intracytoplasmic sperm injection (ICSI) and somatic cell nuclear transfer (SCNT). All these biotechniques are interesting tools for applying the reproductive management (Kumar et al. 2016) and biodiversity conservation (Moulavi et al. 2017).

The capability of embryonic development depends on successful IVM, since it stimulates the oocyte to achieve the meiotic competence required (Lee et al. 2014) through the nuclear and cytoplasmic events that promote the synthesis of RNA and proteins (Sirard et al. 2006). Therefore, the IVM quality is a determinant factor for the success of the IVEP (Zhang et al. 2015, Park et al. 2017). In this sense, the establishment of culture environment, as oocyte incubation time, should be clarified, once all IVM events require a specific time that could vary according to the peculiarities of each species (Higaki et al. 2017).

The optimization of IVM conditions for collared peccaries is interesting under various concerns. Although this species is

not threatened (IUCN, 2015), their phylogenetic proximity to endangered species as the *Catagonus wagneri* (Keuroghlian et al. 2014), as well as the commercial importance on their meat and leather (Nogueira-Filho et al. 2004), have been leading to the development of studies focused on their reproductive physiology (Maia et al. 2014, Miranda-Moura et al. 2016). Based on these studies, it was possible to know that specifically for females, they have symmetrical and oval ovaries with a relatively smooth surface, which can present irregular appearance due to corpus luteum or follicles (Garcia et al. 2009). Moreover, the ovaries are enveloped by an ovarian bursa (Guimarães et al. 2012), as also observed in swine (Męczyński 1974). Additionally, collared peccaries' estrous cycle lasts 21 days, with a follicular phase of 6 days and 15 days for the luteal phase (Maia et al. 2014). On the other hand, these individuals have a lower ovulation rate (2.3 follicles) when compared to swine (Sowls 1997) and usually produce 1.7-1.9 offspring per female after 141-151 days of gestation (Mayor et al. 2005).

Therefore, in view of the need for studies related to the reproductive biotechniques in collared peccaries, the initial point has been to observe reproductive relationship with close domestic phylogenetic species, as swine and others artiodactyls. Thus, similarities were observed for folliculogenesis (Guimarães et al. 2012), hormonal profiles (Ahuja-Aguirre et al. 2017) and follicular histological characteristics (Lima et al. 2013) between collared peccaries and swine and/or artiodactyla species, as buffalos.

In this context, IVM time used for buffalo oocytes is of 24 h (Kadom et al. 2014), while maturation time of 48 h is used for porcine oocytes (Pereyra-Bonnet et al. 2008). Moreover, it had observed that a suboptimal culture environment for porcine oocytes may promote nuclear maturation without the structures reaching cytoplasmic maturation (Abeydeera 2002), confirming that the IVM time to be precise and defined (Sirard et al. 2006). Therefore, this is first study was carried out to investigate the relationship between the meiotic potential and the IVM times of collared peccary oocytes. This information will be the first about the IVM of oocytes derived from collared peccary, aiming to direct pathways to obtain an embryo *in vitro* of this species. This step can serve as a basis for both enlightening studies of embryogenesis and physiological aspects of the animal. In addition, it may be the starting point for advanced biotechnology that can increase zootechnical potential and ways to ensure effective tools that can be used for conservation of phylogenetically near species.

MATERIALS AND METHODS

Animal ethics and care. The experiment was conducted in accordance with the Committee for Ethics in Animal Use of Federal Rural University of Semi-Arid (CEUA/UFERSA, no. 23091.001072/2015-92) and the Chico Mendes Institute for Biodiversity Conservation (ICMBio, no. 48633-2). The animals used in this research belong to the Centre of Multiplication of Wild Animals (CEMAS/UFERSA, Mossoró, RN, Brazil, 5°10'S, 37°10'W), registered at the Brazilian Institute of Environment and Renewable Natural Resources (IBAMA) as a scientific breeding site (no. 1478912). The breeder contains 200 collared peccaries on average. Annually, a programmed slaughter is conducted for population control and the biological material is destined for several experiments.

Reagents and media. All reagents and media used were obtained from Sigma-Aldrich (St. Louis, MO, USA), but fetal bovine serum

(FBS) was obtained from Gibco-BRL (Carlsbad, CA, USA). When necessary, the pH was adjusted to 7.2-7.4 for all media. Media were filtered using 0.22µm system (Corning, NY, USA).

Oocyte collection, selection and *in vitro* maturation.

Ovaries were obtained from nine adult female collared peccaries (age 4-10 years). Eighteen ovaries were removed from the ovarian bursa and transported to the laboratory in 0.9% (w/v) NaCl solution at 37°C for up to 1 h. Antral follicles (3-6 mm) were aspirated with 21 gauge needle attached to a 5.0 mL syringe. After the aspiration, for a greater oocyte recovery, ovaries were sliced with a scalpel blade in 100x20 mm plates. All follicular fluid was pooled in conical tubes for ten minutes. The sediment was diluted in the oocyte collection medium composed of TCM 199 culture medium (M2520) supplemented with 2.2g/L bicarbonate (S5761), 0.2 mM sodium pyruvate (P45622), 1% antibiotic-antimycotic solution (A5955) and 10% FBS (12657-029). Thus, *cumulus*-oocyte complexes (COCs) were selected according to the number of *cumulus* cells layers and homogeneity of the cytoplasm, being classified in four grades. Grade 1 (3-5 *cumulus* cells layers and homogeneous cytoplasm); Grade 2 (1-2 *cumulus* cells layers and homogeneous cytoplasm); Grade 3 (partially denuded cytoplasm and slightly heterogeneous); Grade 4 (denuded and heterogeneous cytoplasm), according to Kumar et al. (2016). Only grade 1 and 2 COCs were considered viable to be matured.

After three washes in oocyte collection medium, the COCs were incubated in 100 µL drops of maturation medium composed of TCM 199 with 2.2g/L bicarbonate, 0.2mM sodium pyruvate, 1% antibiotic-antimycotic solution and supplemented 10µg/mL follicle stimulating hormone (Folltropin®-V, Bioniche, Canada), 100µM cysteamine (M9768) and 10% FBS. All drops were covered with mineral oil and cultured at 38.5°C in humidified atmosphere of 5% CO₂ and divided in two groups, 24 and 48 h.

Assessment of *in vitro* maturation. After the IVM, COCs were evaluated by the analysis of *cumulus* cells expansion using a stereomicroscope at 20-40x magnification. Thus, oocytes with *cumulus* cells were considered matured (Appeltant et al. 2015). Subsequently, *cumulus* cells were removed by pipetting the oocytes

in oocyte collection medium and denuded structures were verified for the presence of the first polar body under stereomicroscope at 20-40X magnification. Thus, oocytes with first polar body were considered matured (Pereyra-Bonnet et al. 2008).

The nuclear status of oocytes was determined by marking it with Hoechst 33342 (B2261). For this, denuded oocytes were fixed in paraformaldehyde buffered in 4% phosphate buffered solution (PBS) for 30 min and then washed in PBS with 0.4% bovine serum albumin (BSA). Afterward, Hoechst 33342 (10µg/mL) labeling was performed for 15 min and exposed under UV light in fluorescence microscope. Finally, oocytes presenting nucleus in metaphase II (metaphase plate, MII) were considered matured and those in other nuclear phases (prophase I, metaphase I, anaphase I and telophase I) were considered immature (Appeltant et al. 2015).

Data analysis. Data were obtained from four sessions, with 2-3 females per session, totaling in nine experimental animals. For statistical analysis, the Graphpad Instat 3.06 software (GraphPad Software Inc., La Jolla, CA, USA) was used. Mean numbers in percentage ± standard error of the matured oocytes in two different times (24 vs. 48 h) were compared using Fisher exact test (P<0.05) for expansion of *cumulus* cells, presence of the first polar body and nuclear status.

RESULTS

Under gross evaluation, the bursa surrounded all peccaries' ovaries, which were symmetric, presenting an oval shape. Moreover, they had a uniform appearance on most of its surface (Fig.1A); nevertheless, some elevations were observed derived from large follicles or the presence of corpus luteum (Fig.1B).

Afterward all sessions, a total of 18 ovaries resulted in 59 immature oocytes from follicles aspirated and sliced follicles with an estimated recovery rate of 83.1% (approximately, 59/71; oocytes recovered/follicles aspirated) and 3.3 oocytes per ovary. Of these, 42 structures (71.2%, 42 structures viable/59 recovered structures) were classified as viable according to the number of *cumulus* cells layers and homogeneity of the cytoplasm, resulting in 2.3 viable oocytes per ovary, (Fig.2A).

After the IVM, the time of 48 h was more adequate than 24 h for the IVM in all evaluated parameters, as *cumulus* cells




Fig.1. Macroscopic aspects of collared peccary ovaries. (A) Uniform appearance with small follicles. (B) Presence of large follicles and corpus luteum. The arrowhead indicates a small follicle and arrow shows a large follicle.




Fig.2. *In vitro* maturation (IVM) of oocytes of collared peccary. (A) Immature oocytes before IVM. Obj.10x. (B) Cumulus cell expansion after 24 h IVM, Obj.10x. (C) Cumulus cell expansion after 48 h IVM. Obj.10x. (D) Oocyte with extrusion of first polar body. Obj.40x. (E) Oocyte in MII stage. Obj.40x.

Table 1. *In vitro* maturation of collared peccary oocytes after different incubation times

Evaluation of <i>in vitro</i> maturation	Incubation times (in hours)	
	24	48
Expansion of cumulus cells	38.1 ± 1.4 (8/21) ^b	100.0 ± 0.0 (21/21) ^a
Presence of the first polar body	52.4 ± 2.2 (11/21) ^b	90.5 ± 2.0 (19/21) ^a
Nuclear status in second metaphase	19.0 ± 1.4 (4/21) ^b	76.2 ± 1.3 (16/21) ^a
Mean in % ± standard error (n). ^{a,b} In same row differ (P<0.05).		

expansion ($P=0.0002$), presence of the first polar body ($P=0.0148$) and nuclear status in second metaphase ($P=0.0003$) (Table 1). Moreover, a 100% cumulus cells expansion was observed for oocytes cultured for 48 h (Fig.2C) when compared to 24 h (Fig.2B). Additionally, oocytes with first polar body (Fig.2D) and MII presence (Fig.2E) were evidenced in more of 50% and 19% of the oocytes, respectively.

DISCUSSION

The results indicated that the most appropriate incubation period for collared peccaries' oocytes achieve meiotic competence was 48 h. This time is similar to that observed for porcine oocytes (Pereyra-Bonnet et al. 2008). In this sense, we can infer that similar follicular characteristics described between collared peccaries and swine (Lima et al. 2013) justify this relationship, since the *in vitro* conditions for maturation should mimic the *in vivo* conditions that occur in the females.

Moreover, other peculiar feature of collared peccary oocytes similar to the porcine oocytes is the presence of cytoplasmic lipid droplets (Lima et al. 2013). This aspect is crucial for the establishment of the appropriate requirements for IVM, evidencing that further IVM protocols should be adapted from porcine to peccaries' oocytes. Nevertheless, a factor that still needs to be further investigated by the similarity that collared peccary oocytes had in relation to porcine oocytes is that the

IVM in swine oocytes have their altered medium to adapt to the variable requirements of IVM (Gruppen 2014) and this step may improve the quality of matured oocytes derived from collared peccaries. Thus, the tracks of such similarities between peccaries and porcine oocytes indicate a starting point, but not the direct application of protocols without further experimentation, considering that differences occur between these species for follicular morphology (Lima et al. 2013).

On the other hand, the follicular development in collared peccaries was also similar to that found for other artiodactyl species regarding the stages of folliculogenesis, i.e. recruitment, selection and dominance (Guimarães et al. 2012) when compared to buffalos (Manik et al. 2002) and camelids (Basiouni 2007). Nevertheless, despite this similarity, the 24 h IVM time that is frequently used for buffaloes (Kadoom et al. 2014) was not suitable for oocyte maturation in the peccaries. The IVM time to guarantee the formation of the metaphase plate is essential to ensure that the oocytes possess the ability to develop quality embryos (Gruppen 2014). In this sense, the 24 h of IVM for collared peccary oocytes was not long enough for the organization of the metaphase plate (19.0%), while the majority of oocytes were able to reach metaphase II in 48 h of IVM (76.2%). As developmental competence is also reached in cytoplasmic maturation, which occurs after meiotic maturation and refers to the ability of the oocyte to

be fertilized and to block polyspermy, this type of maturation can be visualized by migration of the cortical granules to the oocyte periphery and by the expansion of *cumulus* cells (Milakovic et al. 2015). The analysis of *cumulus* cells expansion at 24 and 48 h showed different degrees of expansion (38.1% vs. 100.0%, respectively); these may be related to cytoplasmic maturation, being an indicator of successful maturation (Abeydeera 2002). Therefore, 24 h of IVM is not sufficient for IVM of collared peccary oocytes (Fig. 2B).

Several important factors for the IVM success are dependent of the incubation time, which can influence the final quality of the embryonic development (Miyoshi et al. 2002). The signaling pathway of 3',5' cyclic adenosine monophosphate (cAMP) is highly time-dependent and restricted to time mechanisms, which is what triggers germinal vesicle breakdown, so nuclear maturation depends on cAMP so the oocyte can reach metaphase II (Sirard et al. 2006). Likewise, another time dependent factor is the secretion of steroid hormones by COCs derived small and large follicles during IVM. Topfer et al. (2016) observed that oocytes derived from small follicles have a lesser ability to support estradiol synthesis in the first half of IVM (28 h) and to suppress progesterone synthesis at the end of IVM (48 h) compared with oocytes from large follicles. Thus, the steroid hormone secretion by COCs is time-dependent and therefore, the meiotic maturation time should be to be precise and defined (Sirard et al. 2006, Topfer et al. 2016).

Moreover, the *cumulus* cells play a bidirectional role for oocyte development. One of the factors that contribute to this role is their mitochondria and mitochondrial DNA, besides contributing to the passage of nutrients and growth factors to the oocyte (Pawlak et al. 2016). Additionally, it is known that the *cumulus* cells play a regulatory role for cAMP, showing great relevance for oocyte maturation (Gruppen 2014). Therefore, the clear expansion of *cumulus* cells with 48 h (Fig. 2C) in this study ensured the optimal performance of the cells during IVM. Further adjustments of cAMP levels during maturation may improve fertilization capacity and oocyte development (Appeltant et al. 2015).

Finally, other factors, as follicles size and technique employed for oocyte collection, may influence the IVM conditions (Sun et al. 2001). In this study, 3–6 mm follicles aspirated and matured for 48 h allowed the obtaining of oocytes to reach meiotic competence. Moreover, the recovery rate obtained can be due to the use of two methods of oocytes collection, first the follicular aspiration and then the slicing technique. The slicing technique allows the increase oocyte recovery, especially when there is limited availability of ovaries (Ferraz et al. 2016). Moreover, viability rate of 71.2% can be attributed to the good development of immature oocytes in the follicular environment (Gruppen 2014) and the efficiency of the methods for oocyte recovery.

In conclusion, 48 h IVM for collared peccary is the most suitable incubation period for oocytes reaching the meiotic competence when compared to 24 h time. This was the first study with *in vitro* maturation, in an attempt to improve a protocol of IVEP for this species. Although this time has been defined in the present work, further details for the improvement of IVM, as medium composition, should be considered not only to achieve meiotic competence but also to guarantee appropriate embryonic development.

Acknowledgement. This study was supported by National Counsel of Technological and Scientific Development (CNPq) and Coordination for the Improvement of Higher Education Personnel (CAPES). The authors thank the CEMAS/UFERSA for providing the animals. A.R. Silva and M.F. Oliveira were recipients of CNPq grants.

REFERENCES

- Abeydeera L.R. 2002. *In vitro* production of embryos in swine. *Theriogenology* 57(1):256-273. <[http://dx.doi.org/10.1016/S0093-691X\(01\)00670-7](http://dx.doi.org/10.1016/S0093-691X(01)00670-7)> <PMid:11775974>
- Ahuja-Aguirre C., López-deBuen L., Rojas-Maya S. & Hernández-Cruz B.C. 2017. Progesterone and estradiol profiles in different reproductive stages of captive collared peccary (*Pecari tajacu*) females assessed by fecal metabolites. *Anim. Reprod. Sci.* 180:121-126. <<http://dx.doi.org/10.1016/j.anireprosci.2017.03.008>> <PMid:28336266>
- Appeltant R., Beek J., Vandenberghe L., Maes D. & Van Soom A. 2015. Increasing the cAMP concentration during *in vitro* maturation of pig oocytes improves *cumulus* maturation and subsequent fertilization *in vitro*. *Theriogenology* 83(3):344-352. <<http://dx.doi.org/10.1016/j.theriogenology.2014.09.023>> <PMid:25442019>
- Basiouni G.F. 2007. Follicular wave pattern, folliculogenesis and assisted reproductive techniques in the non-pregnant female dromedary camel (*Camelus dromedarius*): a review. *J. Biol. Sci.* 7(6):1038-1045. <<http://dx.doi.org/10.3923/jbs.2007.1038.1045>>
- Ferraz M.S., Moraes Júnior F.J., Feitosa M.L.T., Almeida H.M., Bezerra D.O., Pessoa G.T., Albuquerque D.M.N. & Carvalho M.A.M. 2016. Ovarian slicing technique to obtain oocytes from agoutis (*Dasyprocta prymnolopha*). *Pesq. Vet. Bras.* 36(3):204-208. <<http://dx.doi.org/10.1590/S0100-736X2016000300009>>
- Garcia A.R., Kahwage P.R. & Ohashi O.M. 2009. Aspectos reprodutivos de caíto (*Tayassu tajacu*). *Revta Bras. Reprod. Anim.* 33:71-81.
- Gruppen C.G. 2014. The evolution of porcine embryo *in vitro* production. *Theriogenology* 81(1):24-37. <<http://dx.doi.org/10.1016/j.theriogenology.2013.09.022>> <PMid:24274407>
- Guimarães D.A., Garcia S.C.G., Pantoja M.A.F., Silva S.S., Albuquerque N.I. & Le Pendu Y. 2012. Ovarian folliculogenesis in collared peccary: *Pecari tajacu* (Artiodactyla: Tayassuidae). *Revta Biol. Trop.* 60(1):437-455. <PMid:22458237>
- Higaki S., Kishi M., Koyama K., Nagano M., Katagiri S., Takada T. & Takahashi Y. 2017. Early germinal vesicle breakdown is a predictor of high preimplantation developmental competent oocytes in mice. *Zygote* 25(1):41-48. <<http://dx.doi.org/10.1017/S0967199416000290>> <PMid:27873567>
- IUCN 2015. *Catagonus Wagneri*. IUCN Red List of Threatened Species. Version 2015.4. International Union for Conservation of Nature. <<http://dx.doi.org/10.2305/IUCN.UK.2015-2.RLTS.T4015A72587993.en>>
- Kadom A.K., Abdel-Khalek A.E., Shamiyah S.M., El-Sharawy M.E. & El-Razek I.A. 2014. *In vitro* maturation, fertilization and development of prepubertal and mature buffalo oocytes. *Egyptian J. Anim. Prod.* 51:65-69.
- Keuroghlian A., Andrade Santos M.D.C. & Eaton D.P. 2014. The effects of deforestation on white-lipped peccary (*Tayassu pecari*) home range in the southern Pantanal. *Mammalia* 79(4):1-7.
- Kumar S., Devi B., Purkayastha A., Bharti P.K., Doley S. & Kadirvel G. 2016. Porcine ovarian biometry, oocyte retrieval and quality of oocytes under different seasons in north east hill region. *Indian J. Anim. Sci.* 86:300-303.
- Lee J., Lee Y., Park B., Elahi F., Jeon Y., Hyun S.H. & Lee E. 2014. Developmental competence of IVM pig oocytes after SCNT in relation to the shrinkage pattern induced by hyperosmotic treatment. *Theriogenology* 81(7):974-981. <<http://dx.doi.org/10.1016/j.theriogenology.2014.01.022>> <PMid:24576710>
- Lima G.L., Santos E.A.A., Luz V.B., Rodrigues A.P.R. & Silva A.R. 2013. Morphological characterization of the ovarian preantral follicle population

- of collared peccaries (*Tayassu tajacu* Linnaeus, 1758). Anat. Histol. Embryol. 42(4):304-311. <http://dx.doi.org/10.1111/ahe.12021> <PMid:23278244>
- Maia K.M., Peixoto G.C., Campos L.B., Bezerra J.A.B., Ricarte A.R., Moreira N., Oliveira M.F. & Silva A.R. 2014. Estrus cycle monitoring of captive collared peccaries (*Pecari tajacu*) in semiarid conditions. Pesq. Vet. Bras. 34(11):1115-1120. <http://dx.doi.org/10.1590/S0100-736X2014001100014>
- Manik R.S., Palta P., Singla S.K. & Sharma V. 2002. Folliculogenesis in buffalo (*Bubalus bubalis*): a review. Reprod. Fertil. Dev. 14(5-6):315-325. <http://dx.doi.org/10.1071/RD01126> <PMid:12467356>
- Mayor P., López-Gatius F. & López-Béjar M. 2005. Integrating ultrasonography within the reproductive management of the collared peccary (*Tayassu tajacu*). Theriogenology 63(7):1832-1843. <http://dx.doi.org/10.1016/j.theriogenology.2004.08.007> <PMid:15823342>
- Męczyński S. 1974. Morphohistological structure of female genital organs in sousliks. Acta Theriologica 19(1-13):91-106. <http://dx.doi.org/10.4098/ATArch.74-7> <PMid:4478856>
- Milakovic I., Jeseta M., Hanulakova S., Knitlova D., Hanzalova K., Hulinska P., Machal L., Kempisty B., Antosik P. & Machatkova M. 2015. Energy status characteristics of porcine oocytes during *in vitro* maturation is influenced by their meiotic competence. Reprod. Domest. Anim. 50(5):812-819. <http://dx.doi.org/10.1111/rda.12592> <PMid:26280917>
- Miranda-Moura M.T.M., Oliveira G.B., Peixoto G.C.X., Pessoa J.M., Papa P.C., Maia M.S., Moura C.E.B. & Oliveira M.F. 2016. Morphology and vascularization of the corpus luteum of peccaries (*Pecari tajacu* Linnaeus, 1758) throughout the estrous cycle. Arq. Bras. Med. Vet. Zootec. 68(1):87-96. <http://dx.doi.org/10.1590/1678-4162-8486>
- Miyoshi K., Rzucidlo S.J., Pratt S.L. & Stice S.L. 2002. Utility of rapidly matured oocytes as recipients for production of cloned embryos from somatic cells in the pig. Biol. Reprod. 67(2):540-545. <http://dx.doi.org/10.1095/biolreprod67.2.540> <PMid:12135893>
- Moulavi F., Hosseini S.M., Tanhaie-Vash N., Ostadhosseini S., Hosseini S.H., Hajinasrollah M., Asghari M.H., Gourabi H., Shahverdi A., Vosough A.D. & Nasr-Esfahani M.H. 2017. Interspecies somatic cell nuclear transfer in Asiatic cheetah using nuclei derived from *post-mortem* frozen tissue in absence of cryo-protectant and *in vitro* matured domestic cat oocytes. Theriogenology 90:197-203. <http://dx.doi.org/10.1016/j.theriogenology.2016.11.023> <PMid:28166968>
- Nogueira-Filho S., Nogueira S., Mendes A. & Jori F. 2004. A large-scale commercial farming of collared peccary (*Tayassu tajacu*) in north-eastern Brazil. Game Wildl. Sci. 21:413-420.
- Park S.J., So K.H. & Hyun S.H. 2017. Effect of zeaxanthin on porcine embryonic development during *in vitro* maturation. J. Biomed. Res. 31(2):154-161. <PMid:28808197>
- Pawlak P., Chabowska A., Malyszka N. & Lechniak D. 2016. Mitochondria and mitochondrial DNA in porcine oocytes and *cumulus* cells - A search for developmental competence marker. Mitochondrion 27:48-55. <http://dx.doi.org/10.1016/j.mito.2015.12.008> <PMid:26705762>
- Pereyra-Bonnet F., Fernández-Martín R., Olivera R., Jarazo J., Vichera G., Gibbons A. & Salamone D. 2008. A unique method to produce transgenic embryos in ovine, porcine, feline, bovine and equine species. Reprod. Fertil. Dev. 20(7):741-749. <http://dx.doi.org/10.1071/RD07172> <PMid:18842176>
- Rahman A.N.M.A., Abdullah R.B. & Wan-Khadijah W.E. 2008. *In vitro* maturation of oocytes with special reference to goat: a review. Biotechnology 7:599-611.
- Sirard M.A., Richard F., Blondin P. & Robert C. 2006. Contribution of the oocyte to embryo quality. Theriogenology 65(1):126-136. <http://dx.doi.org/10.1016/j.theriogenology.2005.09.020> <PMid:16256189>
- Sowls L.K. 1997. Javelinas and Other Peccaries: the biology, management and use. Texas A&M University Press, College Station, Texas, p.86-104.
- Sun Q.Y., Lai L., Bonk A., Prather R.S. & Schatten H. 2001. Cytoplasmic changes in relation to nuclear maturation and early embryo developmental potential of porcine oocytes: effects of gonadotropins, *cumulus* cells, follicular size, and protein synthesis inhibition. Mol. Reprod. Dev. 59(2):192-198. <http://dx.doi.org/10.1002/mrd.1022> <PMid:11389554>
- Topfer D., Ebeling S., Weitzel J.M. & Spannbrücker A.C. 2016. Effect of follicle size on *in vitro* maturation of pre-pubertal porcine *cumulus* oocyte complexes. Reprod. Domest. Anim. 51(3):370-377. <http://dx.doi.org/10.1111/rda.12688> <PMid:27040379>
- Zhang J.Y., Jiang Y., Lin T., Kang J.W., Lee J.E. & Jin D.I. 2015. Lysophosphatidic acid improves porcine oocyte maturation and embryo development *in vitro*. Mol. Reprod. Dev. 82(1):66-77. <http://dx.doi.org/10.1002/mrd.22447> <PMid:25564987>


1 **ANEXO E – VERSÃO DO ARTIGO PUBLICADO NA THERIOGENOLOGY**

2
3 **PRODUCTION OF COLLARED PECCARY (*Pecari tajacu* LINNAEUS, 1758)**
4 **PARTHENOGENIC EMBRYOS FOLLOWING DIFFERENT OOCYTE CHEMICAL**
5 **ACTIVATION AND *IN VITRO* MATURATION CONDITIONS**

6
7
8 **QUALIS:** Quadriênio 2013-2016: A2/ Novo Qualis: A1

9 **FATOR DE IMPACTO:** 2,299

10 **DOI:** <https://doi.org/10.1016/j.theriogenology.2019.10.016>



Production of collared peccary (*Pecari tajacu* Linnaeus, 1758) parthenogenic embryos following different oocyte chemical activation and *in vitro* maturation conditions

Alana Azevedo Borges^a, Maria Valéria de Oliveira Santos^a, Lucas Emanuel Nascimento^a, Gabriela Pereira de Oliveira Lira^a, Érika Almeida Praxedes^a, Moacir Franco de Oliveira^b, Alexandre Rodrigues Silva^c, Alessandra Fernandes Pereira^{a,*}

^a Laboratory of Animal Biotechnology, Federal Rural University of Semi-Arid, Mossoro, RN, Brazil

^b Laboratory of Applied Animal Morphophysiology, Federal Rural University of Semi-Arid, Mossoro, RN, Brazil

^c Laboratory of Animal Germplasm Conservation, Federal Rural University of Semi-Arid, Mossoro, RN, Brazil

ARTICLE INFO

Article history:

Received 2 July 2019

Received in revised form

13 September 2019

Accepted 13 October 2019

Available online 14 October 2019

Keywords:

Wildlife

Somatic cell nuclear transfer

Artificial activation

Embryo development

Epidermal growth factor

ABSTRACT

To optimize the protocols for assisted reproductive techniques (ARTs) in collared peccary (*Pecari tajacu* Linnaeus, 1758), we evaluated various conditions for oocyte *in vitro* maturation (IVM) and chemical activation. Initially, we assessed the IVM rates, cumulus-oocyte complex (COC) quality, and oocyte morphometry in the absence or presence of epidermal growth factor (EGF). There was no difference between the COCs matured in absence or presence of EGF for the expansion of cumulus cells ($97.6\% \pm 1.2$ vs. $100\% \pm 0.0$), presence of first polar body ($65.9\% \pm 1.2$ vs. $70.5\% \pm 1.8$), nuclear status in second metaphase ($62.5\% \pm 11.6$ vs. $68.4\% \pm 4.9$), cytoplasmic maturation ($100.0\% \pm 0.7$ vs. $75.0\% \pm 0.7$), reactive oxygen species levels (0.5 ± 0.2 vs. 0.3 ± 0.1), and mitochondrial membrane potential (1.1 ± 0.2 vs. 1.1 ± 0.1). However, the zona pellucida thickness of matured COCs was reduced in the presence of EGF. Thus, the EGF group was used for further experiments. The oocytes were artificially activated with ionomycin and four secondary activator combinations [6-dimethylaminopurine (6D), 6D and cytochalasin B (6D + CB), cycloheximide (CHX), and CHX and CB (CHX + CB)]. The effect of immature COCs based on cumulus cell layers and cytoplasm homogeneity (GI and GII or GIII COCs) on embryonic development and quality was evaluated. There was no difference in the cleavage rates among the groups of secondary activators. The cleavage rates of embryos derived from GI/GII and GIII COCs were greater than 72.2% and 25.0%, respectively. Moreover, treatment with CHX showed a reduction in the cleavage rate of embryos derived from GIII COCs when compared to the cleavage rate of embryos derived from GI/GII COCs ($P < 0.05$). Nevertheless, higher rates of blastocyst/total GI and GII COCs were observed in the 6D group ($27.6\% \pm 0.3$) compared to CHX group ($6.9\% \pm 0.3$). Additionally, only 6D treatment resulted in the production of embryos derived from GIII COCs ($25.0\% \pm 0.2$). The percentage of the ICM/total cell ratio was also greater in blastocysts derived from 6D ($42.5\% \pm 19.0$), 6D + CB ($37.9\% \pm 21.9$), and CHX + CB ($43.8\% \pm 19.6$) groups when compared to CHX ($3.6\% \pm 0.1$) group. Thus, the combination of ionomycin and 6D could produce collared peccary embryos by activation of both GI/GII COCs and GIII COCs. These optimized IVM conditions using EGF and chemical activation using ionomycin and 6D in collared peccaries form the first steps for establishing ARTs to conserve this species.

© 2019 Elsevier Inc. All rights reserved.

1. Introduction

In the last decade, a rapid decline in the population of various mammalian species was reported in the South American countries [1]. The collared peccary (*Pecari tajacu* Linnaeus, 1758), a wild pig, is found only in the Americas and is classified as the Least Concern

* Corresponding author. Laboratory of Animal Biotechnology Federal Rural University of Semi-Arid, Av. Francisco Mota, 572, Mossoró, RN, 59625-900, Brazil.

E-mail address: alexandra.pereira@ufersa.edu.br (A.F. Pereira).

species [2]. However, the species is already extinct in eastern and southern Argentina and is declining in some biomes, such as Catinga and the Atlantic forest [1]. Generally, this species plays an important role in the ecosystem as a seed dispersing agent. Thus, the maintenance of these animals in captivity is essential for their sustainable use and conservation [3].

The collared peccary populations can be conserved through assisted reproductive techniques (ARTs), such as cloning by somatic cell nuclear transfer (SCNT) and *in vitro* fertilization (IVF). Cloning by SCNT enables the conservation of species as it can be used for increasing the species population [4], basic research [5], and obtaining induced pluripotent cells [6]. Therefore, establishment of all the stages involved in SCNT can aid in the optimal use of this technology for species conservation [5].

There are various steps involved in cloning via SCNT. *In vitro* maturation (IVM) and artificial oocyte activation are the fundamental steps for the development of SCNT technique to obtain cytoplasmic donor cells (or cytoplasm) and evaluate their developmental competence. Recently, we had demonstrated that the collared peccary cumulus-oocyte complexes (COCs) require maturation times of greater than 24 h for complete meiotic development, which resulted in an IVM rate of 76.2% [7]. However, developmental competence was not evaluated in this study. It is important to understand the development of *in vitro*-matured oocytes to blastocysts.

Moreover, the optimization of IVM medium composition can potentially result in higher IVM rates. In some mammals (swine [8]; sheep [9]; canine [10]), supplementation of IVM medium with 10 ng/mL of epidermal growth factor (EGF) increases the meiotic development and subsequent embryonic development. In pigs, EGF along with gonadotropins enhances oocyte nuclear and cytoplasmic maturation [11]. Although this domestic species is phylogenetically close to collared peccary [12], there are differences in the reproductive features between these species, such as ovulation rate [13,14] and litter size [14]. Although swine and collared peccary are phylogenetically close, they exhibit substantial phylogenetic distance. Thus, direct extrapolations between the collared peccary and swine were not possible. Therefore, we aimed to specifically evaluate the effects of EGF on the collared peccary oocyte development. Additionally, different species may exhibit differential responses, as observed in the collared peccary semen cryopreservation. Treatment with 20% low-density lipoproteins (LDL) improved post-thaw sperm motility and plasma membrane integrity of collared peccaries [15], whereas treatment with LDL at concentrations above 10% did not result in a beneficial effect on porcine semen cryopreservation [16].

To the best of our knowledge, we are not aware of any published reports on the establishment of oocyte artificial activation protocols in collared peccaries. Artificial activation of oocytes is also a critical step of SCNT [5] and the activation procedures vary among species (bovine and equine [17]; goat [18]; swine [19]). This indicates that the response to these artificial activation protocols also may vary depending on the species. Previously, Campos Junior et al. [20] artificially activated the collared peccary oocytes using ionomycin and 6-dimethylaminopurine (6D), which resulted in 10% four-cell embryo stage. Increasing the number of oocytes and optimizing the artificial activation protocols may potentially promote higher rates of embryonic development.

An efficient artificial activation protocol promotes *in vitro* embryonic development competencies by decreasing the level of the maturation promoter factor (MPF) through Ca^{2+} oscillation [18]. Thus, different chemical activation methods may be employed using ionomycin as the primary activator, which allows Ca^{2+} transport and induces calcium influx by activation of endogenous entry pathways [21]. Moreover, primary activation can be used in

combination with secondary activators, such as protein phosphorylation inhibitors (6D) that inhibit the protein kinase activity and promote mitosis, and protein synthesis inhibitors (cycloheximide or CHX) that inhibit the production of cyclin B, which is a regulatory component of MPF [22]. Additionally, cytochalasin B (CB) has been used to prevent the fragmentation of embryos [23] and increase the rate of diploid embryos as well as to effectively inhibit the secondary polar body extrusion [24].

Generally, COCs used for IVM and chemical activation are selected based on cumulus cell layers and cytoplasm homogeneity (Grade I to IV). Only GI and GII COCs are used for IVM, which are considered more appropriate according to morphological classification described above. However, due to the reduced availability of oocytes in wild mammals, evaluation of development in GIII COCs is important to increase the quantities of oocytes in the subsequent stages of SCNT. In some mammals, similar molecular patterns were observed between GI/GII and GIII COCs [25,26]. Thus, this study aimed to evaluate various conditions for IVM and chemical activation of collared peccary oocytes.

2. Materials and methods

All experiments were conducted in accordance with the Animal Ethics Committee of the Federal Rural University of Semi-Arid (Opinion N° 23091.001072/2015–92) and Chico Mendes Institute for Biodiversity Conservation (ICMBio, N° 48633–2). Unless otherwise stated, all chemicals were obtained from Sigma Chemical Co. (St. Louis, MO, USA).

2.1. Cumulus-oocyte complex (COC) collection

Twenty collared peccaries ($n=20$) aged 25.1 ± 2.5 months housed at the Centre of Multiplication of Wild Animals (CEMAS/UFERSA, Mossoró, RN, Brazil; $5^{\circ}10'S$, $37^{\circ}10'W$) and registered at the Brazilian Institute of Environment and Renewable Natural Resources (IBAMA, N° 1478912) were used in this study. The animals were stimulated with 600 IU PG600® (400 IU eCG and 200 IU hCG, MSD Saúde Animal, São Paulo, SP, Brazil) before COC collection, following the methods of Peixoto et al. [27] with modifications. At day 4 post-hormone administration, the animals were euthanized and the ovaries were excised. The ovaries were transported to the laboratory within 1 h and maintained in saline solution (NaCl, 0.9%) supplemented with 0.05 mg/mL penicillin at $35-37^{\circ}\text{C}$.

At the laboratory, all visible follicles were classified as small (<0.3 cm), medium (0.3–0.5 cm), or large (>0.5 cm) [28]. The follicles with 0.2–0.6 cm diameter were aspirated for COC collection using a 21 G needle attached to a 5.0 mL syringe containing COC collection medium (CCM; TCM199 (Gibco-BRL, Carlsbad, CA, USA) supplemented with 0.2 mM sodium pyruvate, 10% fetal bovine serum (FBS, serum (FBS, Gibco-BRL), and 1% antibiotic-antimycotic solution).

2.2. COC classification and IVM

The COCs were observed under a stereomicroscope and classified into four grades following the classification system proposed by Kumar et al. [29], which was based on the number of cumulus cell layers and homogeneity of the cytoplasm: Grade I (3–5 cumulus cells layers and homogeneous cytoplasm); Grade II (1–2 cumulus cells layers and homogeneous cytoplasm); Grade III (partially denuded and slightly heterogeneous cytoplasm); Grade IV (denuded and heterogeneous cytoplasm). Only Grade I/II and Grade III COCs were used for IVM in separate drops, according to the experimental design.

The COCs were divided into drops (15–30 COCs per 100 μL) and

the drops were covered with mineral oil. The COCs were incubated for 44 h at 38.5 °C and 5% CO₂ in a humidified atmosphere. The IVM medium comprised CCM supplemented with 20 µg/mL FSH/LH (Pluset®, Hertape Calier, Juatuba, MG, Brazil), 100 µM cysteamine, and an absence or presence of 10 ng/mL EGF.

2.3. Evaluation of cumulus cells

After IVM, cumulus cell expansion was evaluated using a stereomicroscope. The COCs exhibiting cumulus cell expansion were considered mature. The cumulus cell expansion was scored as follows: score 4, total expansion; score 3, partial expansion; score 2, slight expansion; score 1, minimum expansion; score 0, no expansion [30].

The cumulus cells were removed by successive pipetting. The viability of cumulus cells was evaluated by staining the cells with trypan blue (0.2%) for 2 min. The viable cells were unstained, whereas the non-viable cells were stained blue. All cells were counted in the 4 outer quadrants of the Neubauer chamber [31].

2.4. Assessment of nuclear maturation

The denuded oocytes were observed under a stereomicroscope to assess the presence of first polar body (1 PB). The oocytes exhibiting 1 PB were considered mature. To assess the nuclear stage, denuded oocytes were fixed in 4% paraformaldehyde prepared in phosphate buffer saline (PBS) for 30 min. Next, the oocytes were washed with PBS containing 0.4% bovine serum albumin (BSA) fraction V. The cells were then stained with Hoechst 33342 (10 µg/mL) for 15 min and observed under a fluorescent microscope. The oocytes with nucleus at metaphase II (metaphase plate, MII) stage and 1 PB were considered mature, while those in other nuclear phases were considered immature.

2.5. Quantification of intracellular ROS levels by a dichlorofluorescein assay

For evaluation of oocyte quality after IVM, reactive oxygen species (ROS) levels in the oocytes were determined using the fluorescent probe, 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA; Invitrogen, Carlsbad, CA, USA), according to Santos et al. [31]. Briefly, denuded oocytes were washed twice with PBS and incubated in 500 µL of 5 µM H₂DCFDA at 38.5 °C and 5% CO₂ for 30 min. The stained oocytes were washed twice with PBS and imaged using a fluorescence microscope (Olympus BX51TF, Tokyo, Japan). The fluorescence signal intensity (pixels) was quantified using the ImageJ software (version 1.49v, Java 1.8.0_201, Wayne Rasband, U.S. National Institutes of Health, Bethesda, MD, USA; website: <http://rsb.info.nih.gov/ij/download.html>). The background signal intensity was subtracted from the fluorescent intensity values of the treated oocytes. Immature oocytes were assessed as the calibrator. The relative expression levels were calculated by dividing the measured value of each treatment micrograph with the mean value of the calibrator.

2.6. Assessment of mitochondrial membrane potential ($\Delta\Psi_m$) and cytoplasmic maturation

The $\Delta\Psi_m$ was measured by incubating the oocytes with MitoTracker Red® (CMXRos, Invitrogen) at 500 nM for 30 min [31]. The staining procedures and evaluation of images were similar to those used for the quantification of ROS levels. Additionally, the cytoplasmic maturation of the labeled oocytes was evaluated based on the mitochondrial distribution as follows: peripheral (immature oocytes), dispersed (mature oocytes), and transitional (between

peripheral and dispersed distributions; partially mature oocytes).

2.7. Morphometric evaluation of oocytes after maturation

The denuded oocytes were observed under an inverted microscope for assessing the oocyte morphometric parameters. The images were captured using an inverted microscope (Nikon TS100, Tokyo, Japan) equipped with a camera and image processing software. The morphometric parameters of the captured images were analyzed in the ImageJ software using the scale bar as an arbitrary scale for pixel analysis. The measured morphometric parameters included outer oocyte diameter (ZPO), zona pellucida thickness (ZPT), inner oocyte diameter (ZPI), and ooplasm diameter (OD), as described by Saadeldin et al. [32]. Thus, perivitelline space (PVS) diameter, inner oocyte area (IA), ooplasm area (OA), and PVS area were mathematically calculated based on previous parameters using the following formulae: $IA = 3.14 \times (ZPI/2)^2$; $OA = 3.14 \times (OD/2)^2$; $PVS \text{ diameter} = ZPI - OD$ and $PVS \text{ area} = IA - OA$. Diameters were measured in µm and areas in µm².

2.8. Artificial activation and in vitro development (IVD) of embryos

After IVM, the cumulus cells were removed by pipetting the oocytes in the presence of hyaluronidase for up to 2 min. The oocytes with 1 PB were activated using 5 µM ionomycin (I24222; Gibco-BRL) prepared in CCM for 4 min at 37 °C. Next, the oocytes were washed and incubated with drops of secondary activators at 38.5 °C and 5% CO₂ for 3 h, according to the experimental design. Subsequently, the oocytes were washed and incubated in 50 µL drops of IVD medium covered with mineral oil. Synthetic oviductal fluid (SOF) supplemented with 0.2 mM sodium pyruvate, 0.2 mM L-glutamine, 0.34 mM sodium citrate, 2.8 mM myo-inositol, 2% essential amino acid solution, 1% non-essential amino acid solution, 1% antibiotic-antimycotic solution, 5.0 mg/mL BSA, and 2.5% FBS was employed as the IVD medium. The day of activation was considered as D0 and the cells were cultured until D7. On D3, 50% of the culture medium was replaced with fresh medium.

2.9. Evaluation of blastocysts and differential staining of inner cell mass and trophectoderm cells

After seven days of IVD, the blastocyst cells were fixed in ethanol and labeled with Hoechst 33342 (10 µg/mL) and propidium iodide (PI) (10 µg/mL) for 15 min at 38.5 °C. Next, the cells were washed twice with PBS and placed on glass slides in glycerol droplets. A coverslip was placed over the droplet and the images were captured under UV light at 330–385 nm using a fluorescent microscope. The embryos were visualized individually and the nuclei were counted in the captured fluorescent images. The nuclei of the inner cellular mass (ICM) appeared blue as they were labeled only with Hoechst 33342, while those of the trophectoderm (TE) appeared red or pink as they were labeled with both PI and Hoechst 33342. The numbers of ICM, TE, and total cells were quantified using the ImageJ software [33].

2.10. Experimental design

To evaluate various conditions for IVM and chemical activation of collared peccary oocytes, two experiments were performed. Initially, we assessed the IVM rates, COC quality, and oocyte morphometric parameters after IVM in the absence or presence of 10 ng/mL of EGF (experiment 1). Furthermore, we activated the oocytes with ionomycin and four secondary activator combinations (6D, 6D and CB, CHX, and CHX and CB). The effect of immature COC grades (GI and GII or GIII COCs) on the embryonic development and

quality was evaluated (experiment 2). For the second experiment, only the oocytes grown under optimal IVM conditions were used.

2.10.1. Experiment 1: effects of EGF supplementation on IVM of collared peccary COCs

The COCs were washed with the medium and four replicates (two females per replicate) were used to assess the effect of EGF on IVM. GI and GII COCs derived from eight females were matured in the absence (EGF negative group) or presence of 10 ng/mL of EGF (EGF positive group). Meiotic competence was measured by visualization of the 1 PB and nuclear stage (MII). Moreover, oocytes were evaluated for cytoplasmic maturation based on mitochondrial distribution using MitoTracker Red®. Additionally, the cumulus cells were assessed and the ROS levels and $\Delta\Psi_m$ were quantified. The matured oocytes were compared by morphometric evaluation according to the ZPO, ZPT, OD, PVS area, and PVS diameter.

2.10.2. Experiment 2: effects of secondary activators and COC quality on the development of collared peccary parthenogenetic embryos

After 44 h of IVM, matured GI/GII and GIII COCs derived from twelve females distributed in three replicates were artificially activated and cultured for seven days. The COCs were artificially activated with ionomycin and four secondary activator combinations: 1.9 mM 6D (6D group), 1.9 mM 6D and 7.5 μ g/mL CB (6D + CB group), 10 μ g/mL CHX (CHX group), and 10 μ g/mL CHX and 7.5 μ g/mL CB (CHX + CB group). On day 3 of IVD, the total cleavage rate and number of embryos containing 2 cells, 3–7 cells, or 8 or more cells were quantified. On day 7 of IVD, the total blastocyst formation rate and the developmental phase [initial blastocyst (iB), blastocyst (B), expanded blastocyst (eB), and blastocyst hatching/hatched (hB)] were quantified. To evaluate the quality, the total numbers of cells were counted by differentiating the ICM and TE.

2.11. Statistical analysis

All data are expressed as the mean $\pm$ standard error and were analyzed using StatView 5.0 (SAS Institute Inc., Cary, NC, USA). The normality of the data was verified using the Shapiro-Wilk test, while the homoscedasticity was verified with Levene's test. The morphometric oocyte data were examined by variance analysis (ANOVA) and Tukey's test. The levels of ROS and $\Delta\Psi_m$, as well as the number of blastomers were transformed with arcsine and analyzed by ANOVA followed by Tukey's test. All other data were compared with Kruskal-Wallis test and Dunn (multiple

comparisons) test. The difference was considered statistically significant when the P value was less than 0.05.

3. Results

In total, 630 visible follicles (31.5 follicles per female on average) were obtained upon stimulation of 40 ovaries (Fig. 1A) derived from 20 collared peccaries with PG600®. The number of medium follicles (385 ± 14.1) was higher than that of small follicles (113 ± 1.9 , $P = 0.03$) and large follicles (132 ± 5.4 , $P = 0.04$). The number of small and large follicles was similar ($P > 0.05$).

Moreover, 333 ± 10.7 immature oocytes (Fig. 1B) were obtained from all ovaries with a recovery rate of $52.9\% \pm 4.2$ ($333/630$) and 16.2 ± 2.8 oocytes per female. Of these, 253 ± 8.5 (76.0%) oocytes were classified as GI and GII (12.1 ± 2.2 COCs per female), 37.0 ± 3.3 (11.1%) as GIII (3.1 ± 0.8 COCs per female), and 43.0 ± 1.8 (12.9%) as GIV (2.8 ± 1.0 COCs per female) based on the cytoplasm homogeneity and the number of cumulus cell layers.

3.1. Experiment 1: effects of EGF supplementation on IVM of collared peccary COCs

The effect of EGF on the IVM of collared peccary COCs is shown in Tables 1 and 2. Initially, there was no difference in the IVM rates of COCs exhibiting cumulus cell expansion (Fig. 1C) between EGF positive and negative groups. All COCs exhibited cumulus cell expansion scores of 3 and 4 and the cumulus cell viability was greater than 76.0% for both EGF positive and negative groups. Moreover, there was no difference in the IVM rates of COCs selected on the basis of 1 PB presence (Fig. 1D), MII assessments (Fig. 1E), and cytoplasmic maturation evaluated based on mitochondrial distribution. Additionally, the intracellular ROS levels (Fig. 2A, C, and D) and $\Delta\Psi_m$ in the matured COCs were similar between the EGF positive and EGF negative groups.

There was no difference for most of the evaluated oocyte morphometric assessment parameters. However, the thickness of the zona pellucida in the matured oocytes reduced in the presence of EGF (Table 2). Thus, the IVM medium supplemented with EGF was used for experiment 2.

3.2. Experiment 2: effects of secondary activators and COC quality on the development of collared peccary parthenogenetic embryos

Assessment of embryonic development is described in Tables 3 and 4. The cleavage rate (Fig. 3A–C) after 3 days of IVD was similar




Fig. 1. Representative images of (A) ovaries of collared peccary obtained after hormonal induction with PG600®; (B) Cumulus-oocyte complex (COC) recovered with homogeneous cytoplasm and more one layer of cumulus cells; (C) COCs matured after 44 h IVM; (D) Denuded oocyte with presence of 1 PB; (E) Visualization of the metaphasic plate (MII).

Table 1
Effects of EGF supplementation on *in vitro* maturation of collared peccary cumulus-oocyte complexes (COCs).

Group	Evaluation of <i>cumulus</i> cells				Nuclear maturation		Cytoplasmic maturation mitochondrial distribution		
	Expansion and grade of expansion, %			Viability, %	1 PB,%	MII,%	Peripheral, %	Transition, %	Dispersed, %
	Total	Grade 3	Grade 4						
EGF positive	100.0 ± 0.0 (44/44)	63.6 ± 3.0 (28/44)	29.5 ± 0.3 (13/44)	76.0 ± 3.1 (784/1031)	70.5 ± 1.8 (31/44)	68.4 ± 4.9 (13/19)	0.0 ± 0.0 (0/9)	25.0 ± 0.0 (2/8)	75.0 ± 0.7 (6/8)
EGF negative	97.6 ± 1.2 (40/41)	70.7 ± 2.5 (29/41)	29.3 ± 2.1 (12/41)	79.4 ± 1.4 (765/963)	65.9 ± 1.2 (27/41)	62.5 ± 11.6 (10/16)	0.0 ± 0.0 (0/9)	0.0 ± 0.0 (0/9)	100.0 ± 0.0 (9/9)

No differences were observed among groups ($P > 0.05$). 1 PB, first polar body, MII, metaphase II.

Table 2
Morphometric assessment (in μm and μm^2) in of matured COCs derived from collared peccaries (mean ± standard error).

Groups	Outer oocyte diameter (ZPO)	Zona pellucida thickness (ZPT)	Ooplasm diameter (OD)	Perivitelline space (PVS) area	Perivitelline space (PVS) diameter
EGF positive	373.9 ± 4.4 (37)	33.3 ± 1.2 (37) ^a	284.3 ± 4.2 (37)	7312.6 ± 823.1 (35)	16.2 ± 2.1 (35)
EGF negative	382.2 ± 5.3 (34)	38.6 ± 1.5 (34) ^b	285.6 ± 3.0 (34)	6284.0 ± 390.9 (27)	13.8 ± 0.8 (27)

^{a,b}: Values with different superscript letters within columns are significantly different ($P < 0.05$).

in the evaluated groups. The cleavage rates were greater than 72.2% and 25.0% for the embryos derived from GI/GII and GIII COCs, respectively. Moreover, there was no reduction in the cleavage rates of embryos derived from GI/GII and GIII COCs among the 6D and CHX + CB groups (Table 3).

Nevertheless, the 6D group exhibited higher rates of blastocyst/total cell in the embryos derived from GI/GII COCs than CHX group (Table 3). Moreover, only 6D treatment resulted in the production of embryos derived from GIII COCs (25.0% ± 0.2). Further, the percentages of ICM/total cell ratio in blastocysts derived from 6D, 6D + CB, and CHX + CB group COCs were higher than those in blastocysts derived from CHX group COCs. Interestingly, the total number of blastocyst cells (Fig. 3G and H) derived from 6D + CB and CHX + CB oocyte groups was similar (Table 4).

In relation to embryo development kinetics (Fig. 3D–F), or the proportion of embryos at different stages of development, there was no significant difference among the groups in the percentage of embryos classified with 2, 3–7, or ≥8 cells (Fig. 4A and B). On D7, the percentages of iB, B, eB, or hB embryos were similar among the treatment groups (Fig. 4C, $P > 0.05$).

4. Discussion

Establishment of SCNT is an important strategy to develop this technology as a conservation tool for collared peccary. In this study, we optimized the protocols for improving matured oocyte quality and artificial oocyte activation, which are important for the success of SCNT. Recently [34–37], our group has developed strategies for the establishment of donor nuclei (or karyoplast), which form a database of somatic resources for the species. We have established the conditions for obtaining cytoplasts and evaluated their competence after artificial oocyte activation using different protocols.

Initially, we evaluated the effects of EGF on the IVM of collared peccary oocytes. We, for the first time, demonstrated that the zona pellucida thickness in oocytes matured with EGF was lower than that in oocytes matured without EGF. The thickness of the zona pellucida is an indicator of the success of hatching and implantation of embryos [38]. The elasticity and thinning of zona pellucida are essential for the hatching process. Thick zona pellucida is associated with low-quality embryos. Increased thickness of zona




Fig. 2. Oocytes of collared peccaries after maturation evaluated as intracellular concentration of reactive oxygen species (ROS) levels and mitochondrial membrane potential ($\Delta\Psi\text{m}$). (A) Measurement of fluorescence of ROS levels in matured oocytes in the presence of EGF (EGF positive group) or in the absence (EGF negative group) ($P > 0.05$). (B) Quantification of mitochondrial membrane potential in oocytes matured in the presence of EGF (EGF positive group) or in the absence (EGF negative group) ($P > 0.05$). (C) Oocytes labeled with fluorescent probe H_2DCFDA for ROS quantification with EGF (EGF positive group) and (D) without EGF (EGF negative group). (E) Oocytes labeled with fluorescent probe MitoTracker Red® for $\Delta\Psi\text{m}$ with EGF (EGF positive group) and (F) without EGF (EGF negative group).

Table 3

Embryonic development of collared peccary GI/GII and GIII cumulus-oocyte complexes (COCs) activated with ionomycin in combination with different secondary activators.

Groups	No. COCs		Cleavage, %		Blastocyst/Oocyte, %		Blastocyst/Cleaved, %	
	GI/GII	GIII	GI/GII COCs	GIII COCs	GI/GII COCs	GIII COCs	GI/GII COCs	GIII COCs
6D	34	4	82.9 ± 2.4(29)	75.0 ± 0.4(3)	23.5 ± 0.3(8) ^a	25.0 ± 0.2 (1) ^a	27.6 ± 0.3(8) ^a	33.3 ± 0.3 ^a (1)
6D + CB	36	9	72.2 ± 1.4(26)	77.8 ± 0.7(7)	16.7 ± 0.5(6) ^{a,b}	0.0 ± 0.0(0) ^b	23.1 ± 0.5(6) ^{a,b}	0.0 ± 0.0 (0) ^b
CHX	37	4	78.4 ± 2.2(29) ^A	25.0 ± 0.2(1) ^B	5.4 ± 0.2(2) ^b	0.0 ± 0.0(0) ^b	6.9 ± 0.3(2) ^b	0.0 ± 0.0 (0) ^b
CHX + CB	34	7	76.5 ± 2.2(26)	57.1 ± 0.4(4)	17.6 ± 0.3(6) ^{a,b}	0.0 ± 0.0(0) ^b	23.1 ± 0.2(6) ^{a,b}	0.0 ± 0.0 (0) ^b

^{a,b}: Values with different superscript letters within columns are significantly different ($P < 0.05$). ^{A,B}: Values with different superscript letters within rows are significantly different ($P < 0.05$). 6D group: 6D; 6D + CHX group: 6D and cytochalasin B; CHX group: cycloheximide; CHX + CB group: cycloheximide and cytochalasin B.

Table 4

Cell count of cumulus-oocyte complexes (COCs) as viable and activated with ionomycin in combination with different secondary activators.

Secondary activators	No. of embryos	ICM, %	TE, %	Total	%ICM/Total
6D	5	78.8 ± 35.2	93.2 ± 41.7	172 ± 76.9 ^{a,b}	42.5 ± 19.0 ^a
6D + CB	3	33.7 ± 19.4	65.0 ± 37.5	98.7 ± 57.0 ^a	37.9 ± 21.9 ^a
CHX	2	7.0 ± 0.1	188.0 ± 0.1	195 ± 0.1 ^b	3.6 ± 0.1 ^b
CHX + CB	5	40.2 ± 18.0	57.6 ± 25.8	97.8 ± 43.7 ^a	43.8 ± 19.6 ^a

^{a,b}: Values with different superscript letters within columns are significantly different ($P < 0.05$). ICM: internal cellular mass; TE: trophectoderm; %ICM/total: internal cellular mass/total cell ratio.

pellucida may be caused by an inappropriate *in vitro* environment [38]. Zhou et al. [39] demonstrated that thinning solutions enhance the nuclear maturation of oocytes that previously had thick zona pellucida. The study hypothesized that thinner zona pellucida could facilitate first polar body extrusion. Khanmohammadi et al. [38] observed that a suitable culture medium promotes optimal environment that allows zona pellucida thinning and subsequently facilitates embryonic implantation. Furthermore, morphometric evaluation of matured oocytes has been used to estimate the oocyte developmental competence [32].

The expansion of cumulus cells was observed in almost all

oocytes derived from both EGF positive and negative groups (>97.8%). A similar response was also observed in swine oocytes [8]. Further, both groups exhibited high rates of cumulus cell expansion after maturation. There was no difference in the viability of cumulus cells, with values higher than 76% for all groups. This response may have occurred due to the presence of gonadotrophins and/or serum




Fig. 3. Kinetics of embryonic development in collared peccaries. After 3 days of IVD different embryos with (A) 2 cells; (B) 4 cells; (C) more than 8 cells. Evaluation of embryonic development on day 7 of IVD (D) blastocyst; (E) blastocyst hatching; (F) blastocyst hatched already without zona pellucida. Counting the total number of blastocyst (G) cells inner mass labeled with Hoechst 33342. (H) trophectoderm cells stained with propidium iodide. (I) merge total cells.




Fig. 4. Embryonic development of classified activated with ionomycin in combination with different secondary activators. Embryonic development in 3 days with 2 cells, 3–7 cells and 8 cells after activation with ionomycin in combination with secondary activators in (A) GI/GII ($P > 0.05$) and (B) GIII COCs ($P > 0.05$). (C) Percentage of embryos in D7 classified in different stages of blastocyst after activation of GI/GII COCs ($P > 0.05$).

in the IVM medium, which can interfere with the stimulation induced by EGF [40]. Additionally, FSH may act through EGF receptors to exert a synergistic effect on cytoplasmic maturation [11] and may interfere with the effects of EGF on *cumulus* cell expansion.

The quality of matured oocytes is important for the evaluation of meiotic competence. Thus, we evaluated the ROS levels and $\Delta\Psi_m$ in matured oocytes and observed no difference between the groups for both parameters. Generally, the decrease in ROS levels is accompanied with an increase in glutathione (GSH) levels, a natural antioxidant that reduces ROS in oocytes [41], where ROS generation depends on $\Delta\Psi_m$ [31]. Fan et al. [41] observed that treating the goat oocytes with EGF and cysteamine resulted in lower ROS levels in matured oocytes. In our study, both groups were cultured in a medium containing cysteamine, and EGF probably did not enhance its activity.

Similarly, EGF did not increase the IVM rates of oocytes at MII phase and have 1 PB. Although Kishida et al. [42] observed a higher rate of nuclear maturation in porcine oocytes when EGF was present in the medium (10 ng/mL), another study demonstrated that EGF does not influence the nuclear maturation [43]. This may be due to the interaction of EGF with different supplements of the IVM medium, which are highly variable in each study. Further, EGF was involved primarily in the maintenance of embryonic quality where EGF-activated pathways promote greater competence of the oocyte to support embryonic development. Therefore, we suggest the use of EGF in the IVM medium of collared peccary oocytes for future experiments.

In this study, we observed that 6D was the most optimal secondary activator for collared peccary oocytes. However, the combination of 6D and CB did not have a beneficial effect on embryonic development. Further, CHX did not exhibit good potential for oocyte activation, which was improved only in combination with CB. Several studies have demonstrated the ability of 6D, CHX, and CB to inhibit meiotic resumption and second polar body extrusion in porcine oocytes [44,45]. Moreover, the higher efficiency of 6D compared to CHX was also observed in porcine oocytes [46]. Although 6D and CHX exerted positive effect on oocyte activation and morula-blastocyst formation rates, 6D was more effective than CHX on both matured and immature oocytes [46]. This may be due to the mechanism of action of CHX, which may result in the inhibition of proteins involved in embryonic development [47], whereas 6D specifically inhibits protein kinases [48]. Additionally, Khadijah et al. [49] and Zhang et al. [50] demonstrated that the combination of ionomycin and 6D was better than the combination of ionomycin and CHX for embryonic development in cattle (7.1% vs. 2.2%) and rabbit oocytes (8.6% vs. 1.2%).

Moreover, the embryonic development rates in the presence of ionomycin and 6D for collared peccary oocytes were higher than those observed for swine oocytes [51] evaluated based on the cleavage percentage (82.9% vs. 47.2%), blastocyst percentage (23.5% vs. 11.3%), and ICM/total cell ratio (42.5% vs. 24.8%). This indicates that despite similarities in the responses of protocols used between species, some differences are observed, suggesting the importance of establishing a species-specific protocol. Additionally, Campos Junior et al. [20] activated the collared peccary oocyte with the combination of ionomycin and 6D and obtained up to two-cell (40%) and 4-cell (10%) embryos. Thus, this is the first study that achieved the blastocyst stage of embryonic development in collared peccaries through artificial oocyte activation.

The low performance of CHX was observed in both artificial oocyte activation of collared peccaries and blastocyst quality. Although our results demonstrated a higher total number of cells in the 6D and CHX groups, CHX group exhibited a higher number of trophectoderm cells and a very low number of cells of the internal mass, which would later impair the embryonic development of the

CHX group that had a low blastocyst rate compared to the other groups. The CHX + CB group exhibited higher ICM/total cell rates compared to the CHX group. This may be because CHX has no effect on blocking the extrusion of the second polar body. Thus, parthenotes treated with CHX were largely haploid and exhibited low blastocyst development, which does not improve blastocyst formation further [52]. The combination of 6D and CB was more effective in inhibiting the extrusion of the second polar body, which promotes the development of diploid embryos with higher developmental competence compared to haploid embryos [53].

The combination of ionomycin and 6D exhibited optimal performance as this was the only combination that produced embryos from GIII oocytes. As CB enhances the effect of CHX on GI and GII oocytes, its mechanism cannot be as efficient as 6D because the combination of CHX and CB could not promote the development of GIII oocytes to blastocysts. However, matured GIII oocytes were activated and developed into embryonic blastocyst only in the 6D group. Therefore, embryonic development depends on the efficiency of the activation protocol [53,54]. Thus, use of oocytes with heterogeneous cytoplasm and less than one layer of *cumulus* cells may be suitable when there is low availability of oocytes.

In conclusion, EGF can be used to supplement the maturation medium to obtain a greater quality of matured oocytes and to improve embryonic development. Moreover, the combination of ionomycin and 6D could produce collared peccary embryos from the activation of both GI/GII and GIII COCs. These optimized IVM conditions and artificial activation in collared peccaries represent the first steps for cloning this species through ARTs.

Acknowledgments

The authors thank the CEMAS/UFERSA for providing biological material for the experiments carried out in this research and MSD Animal Health Brazil for the donation of hormones. This study was supported by Brazilian Council of Scientific Development – CNPq and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES, Financial Code 001). AR Silva, MF Oliveira and AF Pereira are recipients of CNPq grants.

References

- [1] Desbiez ALJ, Keuroghlian A, Beisiegel BM, Medici EP, Gatti A, Pontes ARM, et al. Avaliação do risco de extinção do cateto *Pecari tajacu* Linnaeus, 1758, no Brasil. *Biodiversidade Brasileira* 2012;74–83.
- [2] Gongora J, Reyna-Hurtado R, Beck H, Taber A, Altrichter M, Keuroghlian A. *Pecari tajacu*. In: IUCN, editor. The IUCN red list of threatened species. IUCN; 2011. eT41777A10562361. <https://doi.org/10.2305/IUCN.UK.2011.2.RLTS.T41777A10562361.en>; 2011. [Accessed 6 May 2019].
- [3] Albuquerque N, Dias H, Guimarães DDA, Pendu Y, Garcia A, Kahwage P, et al. Criação de caimitos em cativeiro: sistema intensivo de produção na Amazônia Oriental. n.d.. In: Embrapa Amazônia Oriental, editor. Embrapa Amazônia Oriental (INFOTEC-A). Embrapa Amazônia Oriental; 2016. p. 1–108.
- [4] Wani NA, Vettical BS, Hong SB. First cloned Bactrian camel (*Camelus bactrianus*) calf produced by interspecies somatic cell nuclear transfer: a step towards preserving the critically endangered wild Bactrian camels. *PLoS One* 2017;12:e0177800.
- [5] Borges AA, Pereira AF. Potential role of intraspecific and interspecific cloning in the conservation of wild mammals. *Zygote* 2019;27:111–7.
- [6] Honda A, Chojookhuu N, Izu H, Kawano Y, Inokuchi M, Honsho K, et al. Flexible adaptation of male germ cells from female iPSCs of endangered *Tokudaia osimensis*. *Sci Adv* 2017;3:e1602179.
- [7] Borges AA, Santos MVO, Queiroz Neta LB, Oliveira MF, Silva AR, Pereira AF. *In vitro* maturation of collared peccary (*Pecari tajacu* Linnaeus, 1758) oocytes after different incubation times. *Pesqui Vet Bras* 2018;38:1863–8.
- [8] Abeydeera LR, Wang W, Cantley TC, Rieke A, Prather RS, Day BN. Presence of epidermal growth factor during *in vitro* maturation of pig oocytes and embryo culture can modulate blastocyst development after *in vitro* fertilization. *Mol Reprod Dev* 1998;51:395–401.
- [9] Guler A, Poulin N, Mermillod P, Terqui M, Cognie Y. Effect of growth factors, EGF and IGF-I, and estradiol on *in vitro* maturation of sheep oocytes. *Theriogenology* 2000;54:209–18.
- [10] Song H, Kang E, Maeng G, Ock S, Lee S, Yoo J, et al. Influence of epidermal

- growth factor supplementation during *in vitro* maturation on nuclear status and gene expression of canine oocytes. *Res Vet Sci* 2011;91:439–45.
- [11] Uhm S, Gupta M, Yang J, Chung H-J, Min T, Lee H. Epidermal growth factor can be used in lieu of follicle-stimulating hormone for nuclear maturation of porcine oocytes *in vitro*. *Theriogenology* 2010;73:1024–36.
 - [12] Orliac MJ, Pierre-Olivier A, Ducrocq S. Phylogenetic relationships of the suidae (mammalia, cetartiodactyla): new insights on the relationships within suidea. *Zool Scr* 2010;39:315–30.
 - [13] Hellgren EC, Syntzke DR, Oldenberg PW, Guthery FS. Demography of a collared peccary population in South Texas. *J Wildl Manag* 1995;59:153–63.
 - [14] Gottdenker N, Bodmer RE. Reproduction and productivity of white-lipped and collared peccaries in the Peruvian Amazon. *J Zool* 1998;245:423–30.
 - [15] Souza ALP, Lima GL, Peixoto GCX, Castelo TS, Oliveira MGC, Paula VV, et al. Sperm characteristics following freezing in extenders supplemented with whole egg yolk and different concentrations of low-density lipoproteins in the collared peccary (*Pecari tajacu*). *Reprod Biol* 2015;15:223e8.
 - [16] Hu J, Li OW, Li G, Chen XY, Yang H, Zhang SS, et al. The cryoprotective effect on frozen-thawed boar semen of egg yolk low density lipoproteins. *Asian Aust J Anim Sci* 2006;19:486–94.
 - [17] Fernandes CB, Devito LG, Martins LR, Blanco IDP, Lima Neto JF, Tsuribe PM, et al. Artificial activation of bovine and equine oocytes with cycloheximide, roscovitine, strontium, or 6-dimethylaminopurine in low or high calcium concentrations. *Zygote* 2014;22:387–94.
 - [18] Pathak J, Kharche S, Goel A. Effects of different activation protocols on cleavage rate and blastocyst production of caprine oocytes. *Iran J Vet Res* 2017;18:243.
 - [19] Macedo MP, Glanzner WG, Rissi VB, Gutierrez K, Currin L, Baldassarre H, et al. A fast and reliable protocol for activation of porcine oocytes. *Theriogenology* 2019;123:22–9.
 - [20] Campos Junior PHA, Costa GMJ, Avelar GF, Lacerda SMSN, Costa NN, Ohashi OM, et al. Derivation of sperm from xenografted testis cells and tissues of the peccary (*Tayassu tajacu*). *Reproduction* 2014;147:291–9.
 - [21] Cervera R, Silvestre M, Marti N, García-Mengual E, Moreno R, Stojkovic M. Effects of different oocyte activation procedures on development and gene expression of porcine pre-implantation embryos. *Reprod Domest Anim* 2010;45:12–20.
 - [22] Ock S, Kwack D, Mohana Kumar B, Han J, Kim S, Rho G. Effects of activation methods on DNA synthesis and development of parthenogenetic porcine embryos. *Reprod Domest Anim* 2011;46:1082–9.
 - [23] Collas P, Robl JM. Factors affecting the efficiency of nuclear transplantation in the rabbit embryo. *Biol Reprod* 1990;43:877–84.
 - [24] Qiu X, Li N, Xiao X, Li Y. Aggregation of a parthenogenetic diploid embryo and a male embryo improves the blastocyst development and parthenogenetic embryonic stem cell derivation. *Turk J Biol* 2017;41:629–39.
 - [25] Assidi M, Dufort I, Ali A, Hamel M, Algriany O, Dielemann S, et al. Identification of potential markers of oocyte competence expressed in bovine *cumulus* cells matured with follicle-stimulating hormone and/or phorbol myristate acetate *in vitro*. *Biol Reprod* 2008;79:209–22.
 - [26] Almeida KC, Pereira AF, Alcantara Neto AS, Avelar SRG, Bertolini LR, Bertolini M, et al. Real-time qRT-PCR analysis of EGF receptor in *cumulus*-oocyte complexes recovered by laparoscopy in hormonally treated goats. *Zygote* 2011;19:127–36.
 - [27] Peixoto GCX, Lima GL, Maia KM, Souza ALP, Castelo TS, Paiva ALC, Paula VV, et al. Single injection of eCG/hCG leads to successful estrous synchronization in the collared peccary (*Pecari tajacu* Linnaeus, 1758). *Anim Reprod Sci* 2019;298:106112.
 - [28] Viana ACS, Freitas JLS, Campos LB, Silva AR, Oliveira MF, Freitas VJF, et al. Analysis of ovarian response to superstimulation treatment in collared peccary (*Pecari tajacu* Linnaeus, 1758). *Anim Reprod* 2017;4:804.
 - [29] Kumar S, Devi B, Purkayastha A, Bharti PK, Doley S, Kadirvel G. Porcine ovarian biometry, oocyte retrieval and quality of oocytes under different seasons in north east hill region. *Indian J Anim Sci* 2016;86:300–3.
 - [30] Kubo N, Cayo-Colca IS, Miyano T. Effect of estradiol-17 β during *in vitro* growth culture on the growth, maturation, *cumulus* expansion and development of porcine oocytes from early antral follicles. *Anim Sci J* 2015;86:251–9.
 - [31] Santos MVO, Nascimento LE, Praxedes EA, Borges AA, Silva AR, Bertini LM, et al. *Syzygium aromaticum* essential oil supplementation during *in vitro* bovine oocyte maturation improves parthenogenetic embryonic development. *Theriogenology* 2019;128:74–80.
 - [32] Saadeldin IM, Swelum AAA, Yaqoob SH, Alowaimier AN. Morphometric assessment of *in vitro* matured dromedary camel oocytes determines the developmental competence after parthenogenetic activation. *Theriogenology* 2017;95:141–8.
 - [33] Paul AK, Liang Y, Sritrattana K, Nagai T, Parnpai R. Vitrification of bovine matured oocytes and blastocysts in a paper container. *Anim Sci J* 2018;89:307–15.
 - [34] Borges AA, Lima GL, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, et al. Conservation of somatic tissue derived from collared peccaries (*Pecari tajacu* Linnaeus, 1758) using direct or solid-surface vitrification techniques. *Cyto-technology* 2017;69:643–54.
 - [35] Borges AA, Lira GPO, Nascimento LE, Queiroz Neta LB, Santos MVO, Oliveira MF, et al. Influence of cryopreservation solution on the *in vitro* culture of skin tissues derived from collared peccary (*Pecari tajacu* Linnaeus, 1758). *Biopreserv Biobanking* 2018;16:77–81.
 - [36] Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, Pereira AF. Combination of ethylene glycol with sucrose increases survival rate after vitrification of somatic tissue of collared peccaries (*Pecari tajacu* Linnaeus, 1758). *Pesqui Vet Bras* 2018;38:77–81.
 - [37] Santos MLT, Borges AA, Queiroz Neta LB, Santos MVO, Oliveira MF, Silva AR, et al. *In vitro* culture of somatic cells derived from ear tissue of collared peccary (*Pecari tajacu* Linnaeus, 1758) in medium with different requirements. *Pesqui Vet Bras* 2016;36:1194–202.
 - [38] Khanmohammadi N, Movahedin M, Safari M, Sameni HR, Yousefi B, Jafari B, et al. Effect of L-carnitine on *in vitro* developmental rate, the zona pellucida and hatching of blastocysts and their cell numbers in mouse embryos. *Int J Reprod Biomed* 2016;14:649.
 - [39] Zhou HX, Ma YZ, Liu YL, Chen Y, Zhou CJ. Assessment of mouse germinal vesicle stage oocyte quality by evaluating the *cumulus* layer, zona pellucida, and perivitelline space. *PLoS One* 2014;9:e105812.
 - [40] Lorenzo P, Illera M, Illera J, Illera M. Enhancement of *cumulus* expansion and nuclear maturation during bovine oocyte maturation *in vitro* by the addition of epidermal growth factor and insulin-like growth factor I. *Reproduction* 1994;101:697–701.
 - [41] Fan Z, Yang M, Regowski M, Polejaeva IA. Effects of three different media on *in vitro* maturation and development, intracellular glutathione and reactive oxygen species levels, and maternal gene expression of abattoir-derived goat oocytes. *Small Rumin Res* 2017;147:106–14.
 - [42] Kishida R, Lee E, Fukui Y. *In vitro* maturation of porcine oocytes using a defined medium and developmental capacity after intracytoplasmic sperm injection. *Theriogenology* 2004;62:1663–76.
 - [43] Abeydeera L, Wang W, Cantley T, Rieke A, Murphy C, Prather R, et al. Development and viability of pig oocytes matured in a protein-free medium containing epidermal growth factor. *Theriogenology* 2000;54:787–97.
 - [44] Beux G, Richard FJ, Sirard MA. Effect of cycloheximide, 6-DMAP, roscovitine and butyrolactone I on resumption of meiosis in porcine oocytes. *Theriogenology* 2003;60:1049–58.
 - [45] Yi Y, Park C. Parthenogenetic development of porcine oocytes treated by ethanol, cycloheximide, cytochalasin B and 6-dimethylaminopurine. *Anim Reprod Sci* 2005;86:297–304.
 - [46] Demir K, Can A, Ertürk E, Özdemirci S, Karacam H, Sandal AI, et al. Effect of different activation techniques on immature and *in vitro* matured cat oocytes. *Kafkas Univ Vet Fak Derg* 2014;20:565–70.
 - [47] Che L, Lalonde A, Bordignon V. Chemical activation of parthenogenetic and nuclear transfer porcine oocytes using ionomycin and strontium chloride. *Theriogenology* 2007;67:1297–304.
 - [48] Alexander B, Coppola G, Berardino D, Rho G, St John E, Betts D, et al. The effect of 6-dimethylaminopurine (6-DMAP) and cycloheximide (CHX) on the development and chromosomal complement of sheep parthenogenetic and nuclear transfer embryos. *Mol Reprod Dev* 2006;73:20–30.
 - [49] Khadijah S, Rahman M, Abdullah R, Wan K. Comparison of bovine embryonic development of parthenogenetically activated oocytes by different combinations of chemical treatments. *Mal J Anim Sci* 2014;17:85–94.
 - [50] Zhang S, Xiang S, Yang J, Shi J, Guan X, Jiang J, et al. Optimization of parthenogenetic activation of rabbit oocytes and development of rabbit embryo by somatic cell nuclear transfer. *Reprod Domest Anim* 2019;54:258–69.
 - [51] García-Mengual E, Alfonso J, Salvador I, Duque C, Silvestre M. Oocyte activation procedures and influence of serum on porcine oocyte maturation and subsequent parthenogenetic and nuclear transfer embryo development. *Zygote* 2008;16:279–84.
 - [52] Lee J, Tian XC, Yang X. Optimization of parthenogenetic activation protocol in porcine. *Mol Reprod Dev* 2004;68:51–7.
 - [53] Kim Y, Lee S, Ock S, Balasubramanian S, Choe S, Rho G. Development of cloned pig embryos by nuclear transfer following different activation treatments. *Mol Reprod Dev* 2005;70:308–13.
 - [54] Felmer R, Arias M. Activation treatment of recipient oocytes affects the subsequent development and ploidy of bovine parthenogenetic and somatic cell nuclear transfer (SCNT) embryos. *Mol Reprod Dev* 2015;82:441–9.

**ANEXO F – COMPROVANTE DO ARTIGO SUBMETIDO NA ANNALS OF
ANIMAL SCIENCE**

**EFFECTS OF INCUBATION TIME AND METHOD OF CELL CYCLE
SYNCHRONIZATION ON COLLARED PECCARY SKIN-DERIVED FIBROBLAST
CELL LINES**

QUALIS: Quadriênio 2013-2016: Não classificado pela Medicina Veterinária/ Novo Qualis:
A3

FATOR DE IMPACTO: 1,586



Alexsandra Fernandes Pereira <alexandra.pereira@ufersa.edu.br>

Submission of manuscript

k.skupniewicz@izoo.krakow.pl <k.skupniewicz@izoo.krakow.pl>
Para: Alexsandra Fernandes Pereira <alexandra.pereira@ufersa.edu.br>

22 de junho de 2020 09:39

Dear Author,

Thank you for your submission. Your manuscript entitled "Effects of incubation time and method of cell cycle synchronization on collared peccary skin-derived fibroblast cell lines" has been assigned the following number: 317/20 and has been sent for evaluation to the reviewers.

Sincerely,
Editorial Board of Annals of Animal Science
Katarzyna Skupniewicz

W dniu 2020-06-17 23:21, Alexsandra Fernandes Pereira napisał(a):

Dear Dr. Skupniewicz,

I would like to submit for evaluation in ANNALS OF ANIMAL SCIENCE the manuscript titled "EFFECTS OF INCUBATION TIME AND METHOD OF CELL CYCLE SYNCHRONIZATION ON COLLARED PECCARY SKIN-DERIVED FIBROBLAST CELL LINES".

Briefly, our manuscript proposes to develop another stage of cloning by somatic cell nuclear transfer (SCNT) in collared peccary, a wild mammal of ecological, economic and scientific importance and that has suffered population reduction. In the recent years, our group has been working to establish the use of the karyoplast (donor nuclei), which forms a database of somatic resources for the collared peccary (Borges et al., 2017; Borges et al., 2018; Queiroz Neta et al., 2018), in techniques such as cloning by SCNT. We aimed to achieve this goal by defining appropriate methodologies for the synchronization of these cells.

In the manuscript, we evaluated three different approaches for cell cycle synchronization in terms of their effects on synchronization in G0/G1 phases and on the viability of collared peccary skin fibroblasts:

(I) Serum starvation (SS) for 1 to 4 days,

(II) Contact inhibition (CI) for 1 to 3 days,

(III) Using cell cycle regulatory inhibitors (dimethyl sulfoxide, cycloheximide, cytochalasin B, or 6-dimethylaminopurine) for 1 and 2 days.

Several cell cycle synchronization protocols in G0/G1 stages have been proposed. However, the cellular response to these protocols has been variable in domestic and wild mammals, necessitating the evaluation of each protocol for a given cell type and species. Thus, we showed that CI for 3 days was the most efficient method for cell cycle synchronization in collared peccary fibroblasts. To our knowledge, this is the first work to elucidate nuclear reprogramming in cells of this species. Thus, we established the last step of the preparation for the use of these fibroblast as karyoplast for application in somatic cell nuclear transfer of collared peccary, with potential application to conservation of the species.

Moreover, all authors of manuscript agree with the content of the manuscript and submission in this journal. Additionally, none of the authors have conflict of interest. Finally, we sent to an English review company for conference to correct grammatical language, according to Certificate of Editing English. All material submitted

for consideration is original and is being submitted exclusively to
ANNALS OF ANIMAL SCIENCE.

Yours sincerely,

Alexsandra Fernandes Pereira

--

Profª. Dra. Alexsandra Fernandes Pereira
Bolsista de Produtividade do CNPq - Nível 2
Laboratório de Biotecnologia Animal - LBA
Programa de Pós-Graduação em Ciência Animal - PPGCA
[Texto das mensagens anteriores oculto]

--

Katarzyna Skupniewicz

- 1 **ANEXO G – RESUMOS PUBLICADOS NO RENORBIO 2017**
- 2 **I ENCONTRO DE BIOTECNOLOGIA NO NORDESTE**

NA0007 - INFLUÊNCIA DE DIFERENTES CRIOPROTETORES SOBRE A VIABILIDADE DE CÉLULAS SOMÁTICAS DE CATETOS (PECARI TAJACU LINNAEUS, 1758)

Autores: Gabriela Pereira De Oliveira Lira - 1º Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO (UFERSA)); Alexandre Rodrigues Silva - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO (UFERSA)); Moacir Franco De Oliveira - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO (UFERSA)); Luiza Bento De Queiroz Neta - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO (UFERSA)); Maria Valeria De Oliveira Santos - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO (UFERSA)); Lucas Emanuel Nascimento - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO (UFERSA)); Alana Azevedo Borges - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO (UFERSA)); Alessandra Fernandes Pereira1 - Orientador (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO (UFERSA))

Área: Agropecuária **Tipo:** Pesquisa **Nível:** Graduação **STA2:** Não

Resumo: A criopreservação de células somáticas é uma técnica interessante para a conservação de material biológico visando o uso na investigação básica, transferência nuclear (clonagem) e formação de bancos genéticos. Para todas essas aplicações, a escolha dos crioprotetores a serem empregados na criopreservação dessas células consiste na primeira etapa a ser estabelecida. Portanto, o objetivo do presente trabalho foi comparar a influência de diferentes crioprotetores intracelulares (dimetilsulfóxido, DMSO, e etilenoglicol, EG) e extracelular (sacarose, SAC) sobre a viabilidade de células somáticas derivadas de catetos, mamíferos silvestres com importância científica, ecológica e econômica. Para tanto, fragmentos teciduais da região auricular periférica de cinco catetos, mantidos no Centro de Multiplicação de Animais Silvestres (CEMAS/UFERSA), foram cultivados in vitro por 40 dias. Posteriormente, células somáticas recuperadas desses fragmentos foram submetidas à congelação lenta usando a solução de criopreservação constituída por meio essencial mínimo modificado por Dulbecco (DMEM) acrescido de 2,2 g/L de bicarbonato de sódio, 2% de solução de antibiótico-antimicótico, 10% de soro fetal bovino e seis combinações de crioprotetores a seguir: 10% de DMSO (DMSO), 10% de DMSO e 0,2 M de sacarose (DMSO-SUC), 10% de EG (EG), 10% EG e 0,2 M de sacarose (EG-SAC), 10% de DMSO e 10% de EG (DMSO-EG), 10% de DMSO, 10% de EG e 0,2 M de sacarose (DMSO-EG). Para todas as congelações, foi empregado o sistema de congelação Mr. Frosty (Mr. Frosty? Freezing Container) numa taxa de resfriamento de 1°C/min e concentração final de $2,0 \times 10^6$ células/mL. Além disso, células não criopreservadas foi considerada como grupo controle. Para a avaliação da viabilidade, células descongeladas e não criopreservadas foram submetidas ao ensaio de azul de tripan. Todas as análises foram realizadas em duplicatas. Os dados foram expressos como média $\pm$ desvio padrão e analisados usando ANOVA seguida de teste Tukey ($P < 0,05$). Após as análises, apenas células somáticas derivadas da congelação lenta usando DMSO-SAC ($69,8\% \pm 9,6$) mantiveram a viabilidade similar ao grupo não criopreservado ($85,7\% \pm 3,4$; $P > 0,05$), evidenciando um efeito positivo da presença de sacarose sobre a viabilidade celular. Além disso, embora os demais grupos tenham sido diferentes do controle foram similares entre si ($P > 0,05$): EG ($50,8\% \pm 19,5$), EG-SAC ($58,5\% \pm 7,1$), DMSO ($55,8\% \pm 10,1$), DMSO-EG ($58,7\% \pm 11,2$) e DMSO-EG-SAC ($47,9\% \pm 9,8$). Adicionalmente, exceto para o grupo DMSO-SAC, a presença de sacarose não influenciou na manutenção da viabilidade de células somáticas após a congelação lenta. Em conclusão, 10% de DMSO acrescido de 0,2 M de sacarose manteve a viabilidade de células somáticas derivadas de catetos após a descongelação, podendo essa combinação de crioprotetores ser utilizada na conservação de material biológico desta espécie.

Keywords: Palavras chaves: Mamíferos silvestres, criopreservação, cultivo in vitro, crioprotetores intracelulares, sacarose.

NA0005 - INFLUÊNCIA DO TEMPO DE MATURAÇÃO IN VITRO EM OÓCITOS DE CATETOS (PECARI TAJACU LINNEAUS, 1758)

Autores: Alexsandra Fernandes Pereira - 1º Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO); Alexandre Rodrigues Silva - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO); Moacir Franco De Oliveira - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO); Luiza Bento De Queiroz Neta - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO); Lhara Ricarliany Medeiros De Oliveira - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO); Gabriela Pereira De Oliveira Lira - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO); Maria Valeria De Oliveira Santos - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO); Alana Azevedo Borges - Co-Autor (UNIVERSIDADE FEDERAL RURAL DO SEMI-ARIDO)

Área: Agropecuária **Tipo:** Pesquisa **Nível:** STA2:

Resumo: A maturação in vitro (MIV) de oócitos consiste numa etapa essencial para biotecnologias reprodutivas, como a fecundação in vitro, injeção intracitoplasmática de espermatozoides e transferência nuclear de células somáticas. Todas essas biotécnicas podem ser utilizadas como ferramentas para a conservação de mamíferos silvestres, como os catetos. Esses animais se destacam pelo seu potencial zootécnico e econômico, sendo considerados como fonte para pecuária alternativa de pequenos produtores rurais. Nesse sentido, para alcançar o sucesso da MIV é necessário o estabelecimento do tempo de incubação desses oócitos. Portanto, o objetivo do presente trabalho foi avaliar a influência do tempo de MIV (24 h vs. 48 h) sobre o percentual de oócitos maturados de catetos. Para tanto, ovários recuperados de nove fêmeas mantidas do Centro de Multiplicação de Animais Silvestres (CEMAS/UFERSA) foram aspirados e fatiados para a obtenção de oócitos imaturos. Posteriormente, todos os oócitos foram classificados morfológicamente de acordo com a homogeneidade do citoplasma, número e compactação das camadas de células do cumulus em: Grau 1 (3 ou mais camadas de células do cumulus e citoplasma homogêneo); Grau 2 (1-2 camadas de células do cumulus e citoplasma homogêneo); Grau 3 (oócito parcialmente desnudo e ligeiramente heterogêneo); Grau 4 (oócito desnudo e heterogêneo). Os oócitos 1 e 2 foram selecionados e submetidos à MIV em TCM199 com 2,2 g/L de bicarbonato de sódio, 0,2 mM de piruvato de sódio, 1% de solução de antibiótico-antimicótico e suplementado com 10 µg/mL de FSH, 10% de soro fetal bovino e 100 µM de cisteamina a 38,5°C, 5% de CO₂ e umidade máxima por 24 ou 48 h de incubação. Após cada período, as células do cumulus foram removidas por pipetagens e as estruturas desnudadas foram verificadas quanto à presença do primeiro corpúsculo polar (1CP) sob estereomicroscópio. Assim, os oócitos com 1CP foram considerados maturados. Os dados foram analisados pelo teste exato de Fisher ($P < 0,05$) e expressos em média percentual $\pm$ erro padrão. Um total de quatro repetições (2-3 fêmeas por repetição) foi realizado, resultando em 59 oócitos imaturos e com uma taxa de recuperação de 83,1% (59/71) e 3,3 oócitos por ovário. Destes, 42 estruturas foram classificadas como viáveis (2,3 oócitos viáveis por ovário). Após diferentes períodos de incubação, valores de maturação superiores a 50% foram observados para ambos os grupos. Além disso, diferenças ($P < 0,05$) foram observadas entre 24 e 48 h para a presença do 1CP ($52,4 \pm 2,2$ (11/21) vs. $90,5 \pm 2,0$ (19/21)). Em conclusão, o tempo de 48 h foi mais adequado para a MIV de oócitos derivados de catetos, quanto à presença do 1CP. Tais resultados evidenciam que a MIV nesta espécie ocorre em período similar aos oócitos de suínos domésticos, espécie filogeneticamente próxima a esta espécie silvestre.

Keywords: Mamíferos silvestres, biotécnicas reprodutivas, maturação nuclear.

1 **ANEXO H – RESUMO PUBLICADO NO SEMIC 2018**
2 **XXIV SEMINÁRIO DE INICIAÇÃO CIENTÍFICA DA UFERSA**

Área temática: Ciências Agrárias

Influência de crioprotetores extracelulares sobre a criopreservação de células somáticas de catetos (*Pecari tajacu* Linnaeus, 1758)

Gabriela Pereira de Oliveira Lira, Alexsandra Fernandes Pereira, Alana Azevedo Borges, Matheus Barbosa do Nascimento

Resumo

A criopreservação de células somáticas consiste numa interessante ferramenta na conservação de catetos, visando a formação de bancos de recursos genéticos. Esses bancos quando associados à clonagem por transferência nuclear de células somáticas resultam em estratégias eficientes na manutenção da população desta espécie. Contudo, o sucesso desses bancos depende da combinação de crioprotetores utilizados na criopreservação. Embora o crioprotetor intracelular mais empregado seja o dimetilsulfóxido (DMSO) na congelação lenta de células somáticas de catetos, os agentes extracelulares ainda necessitam ser estudados, visando aumentar as taxas de viabilidade após a descongelação. Portanto, o objetivo foi avaliar a influência dos crioprotetores extracelulares, sacarose (SAC) e soro fetal bovino (SFB), sobre a criopreservação de células somáticas de catetos. Para tanto, fragmentos teciduais da pele da região auricular periférica de cinco catetos, pertencentes ao Centro de Multiplicação de Animais Silvestres (CEMAS/UFERSA), foram coletados e transportados ao laboratório em meio essencial mínimo modificado por Dulbecco (DMEM) suplementado com 10% de SFB e 2% de solução de antibióticos e antimicóticos, a 37°C, por até 1 h. Todos os procedimentos foram aprovados pelo Comitê de Ética de Uso de Animais (CEUA/UFERSA, no. 23091.001072/2015-92) e Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio, no. 48633-2). No laboratório, fragmentos (9,0 mm³) foram cultivados *in vitro* sob condições e atmosfera controladas (5% de CO₂ e 38,5°C). Posteriormente, células somáticas recuperadas desses fragmentos foram submetidas à congelação lenta utilizando DMEM acrescido 10% de DMSO suplementado com diferentes combinações de crioprotetores extracelulares: i) 0,2 M de SAC; ii) 10% de SFB; iii) 50% de SFB; iv) 0,2 M de SAC com 10% de SFB; v) 0,2 M de SAC com 50% de SFB. O sistema de congelação empregado foi Mr. Frosty numa taxa de resfriamento de 1°C/min e concentração final de 1,0 x 10⁴ células/mL. Após a descongelação e remoção dos crioprotetores, as células foram submetidas ao ensaio de viabilidade por corante azul de tripan. Todos os dados foram expressos como média ± erro padrão e analisados por ANOVA e Tukey (P < 0,05). Após cinco repetições (1 animal/1 repetição), foi obtida uma taxa média de viabilidade antes da congelação de 78,0% ± 12,9. Após a descongelação, as taxas de viabilidade obtidas em cada grupo foram: 0,2 M de sacarose (66,0% ± 6,3); 10% de SFB (87,8% ± 7,5); 50% de SFB (73,8% ± 10,9); 0,2 M de sacarose com 10% de SFB (67,1% ± 14,5); 0,2 M de sacarose com 50% de SFB (66,6% ± 9,8). Nenhuma diferença foi observada nas taxas de viabilidade avaliada por azul de tripan para células congeladas e não congeladas (P > 0,05). Além disso, nenhuma diferença foi observada entre os grupos criopreservados contendo diferentes crioprotetores extracelulares (P > 0,05). Portanto, células somáticas derivadas de catetos podem ser criopreservadas com sucesso independente do tipo de crioprotetor extracelular empregado.

Palavras-chave: Congelação lenta. Cultivo *in vitro*. Mamíferos silvestres. Sacarose. Soro fetal bovino.

Agência financiadora: Bolsista IC-PIVIC (2017-2018).

1 **ANEXO I – RESUMOS PUBLICADOS NO CONERA 2018**
2 **CONGRESSO NORTE-NORDESTE DE REPRODUÇÃO ANIMAL**

Produção *in vitro* de embriões de cateto (*Pecari tajacu* Linnaeus 1758) usando diferentes ativadores químicos

[*In vitro* production of collared peccary (*Pecari tajacu* Linnaeus 1758) embryos using different chemical activators]

Alana Azevedo Borges^{1*}, Maria Valéria de Oliveira Santos¹, Lucas Emanuel Nascimento¹, Gabriela Pereira de Oliveira Lira¹, Érika Almeida Praxedes¹, Moacir Franco de Oliveira², Alexandre Rodrigues Silva³, Alexsandra Fernandes Pereira¹

¹Laboratório de Biotecnologia Animal, Universidade Federal Rural do Semi-Árido (UFERSA), Mossoró, RN. ²Laboratório de Morfofisiologia Animal Aplicada, UFERSA, Mossoró, RN. ³Laboratório de Conservação de Germoplasma Animal, UFERSA, Mossoró, RN, *E-mail: alanaazevedob@gmail.com

A ativação artificial consiste numa etapa crucial para o sucesso da produção de embriões clones por transferência nuclear de células somáticas (TNCS). Nesse sentido, o estabelecimento de protocolos de ativação oocitária em catetos, mamíferos silvestres de importância ecológica, econômica e científica, representa um dos pontos fundamentais para o desenvolvimento da TNCS nesta espécie. Em mamíferos domésticos, o uso de ativadores químicos, como os ativadores primários de liberação inicial de cálcio [ionomicina] em combinação com ativadores secundários de inibição de proteína quinase [6-dimetilaminopurina (6-DMAP)] ou de síntese proteica [ciclohexamida (CHX)] tem sido empregado para a ativação oocitária com respostas variáveis entre essas associações. Portanto, o objetivo foi produzir embriões *in vitro* de catetos por ativação oocitária artificial usando a ionomicina em associação com diferentes ativadores secundários. Para tanto, oócitos imaturos derivados de doze fêmeas adultas estimuladas com 600 UI de PG600® (MSD Saúde Animal, São Paulo, Brasil) foram recuperados após eutanásia dos animais. Em seguida, apenas oócitos com mais de uma camada de células do *cumulus* e citoplasma homogêneo foram maturados *in vitro* por 44 h (38,5°C e 5% de CO₂) em meio TCM199 modificado. Decorrido o período de maturação, oócitos com o primeiro corpúsculo polar foram ativados artificialmente usando 5,0 µM de ionocimina por 4 min seguida de incubação por 3 h com 1,9 mM de 6-DMAP (grupo 6D) ou 10 µg/mL de CHX (grupo CHX). Os presumíveis zigotos foram cultivados *in vitro* por sete dias e avaliações foram realizadas em D3 e D7 quanto às taxas de clivagem e blastocisto, respectivamente. Além disso, para avaliação da qualidade embrionária, blastocistos foram marcados com Hoechst 33342 para contagem do número de blastômeros. Os dados foram descritos como média ± erro padrão, sendo as taxas de produção embrionária e número de blastômeros analisados pelo teste do chi-quadrado e ANOVA seguido de teste de Tukey, respectivamente (P < 0,05). Após três repetições (4 fêmeas por repetição), nenhuma diferença foi observada para as taxas de clivagem entre os grupos, sendo evidenciado valores superiores a 78% de embriões clivados (6D: 82,9% ± 2,4 e CHX: 78,4% ± 2,2). Além disso, nenhuma diferença foi observada para as taxas de blastocistos, tanto quando foram calculadas pelo número de total de estruturas ativadas (6D: 20,0% ± 0,3 e CHX: 5,4% ± 0,2), quanto pelo número total de embriões clivados (6D: 24,1% ± 0,3 e CHX: 6,9% ± 0,3). Finalmente, também nenhuma diferença foi observada quanto ao número total de blastômeros entre os grupos (6D: 172,0 ± 76,9 e CHX: 195,0 ± 0,0). Esses resultados indicam que ativadores químicos podem ser empregados com sucesso na produção de blastocistos de catetos, independente do mecanismo de ação do ativador secundário (6DMAP ou CHX). Em conclusão, a associação de ionomicina independente do tipo de ativador secundário pode ser empregada na ativação artificial de oócitos de catetos, visando seu uso na clonagem desta espécie.

Palavras-chave: ativação artificial, clonagem, mamíferos silvestres.

Keywords: artificial activation, cloning, wild mammals.

1 **ANEXO J – RESUMO PUBLICADO NO ISABR 2018**
2 **VII INTERNATIONAL SYMPOSIUM ON ANIMAL BIOLOGY OF**
3 **REPRODUCTION**



Evaluation of epidermal growth factor on *in vitro* maturation of collared peccaries' (*Pecari tajacu* Linnaeus, 1758) oocytes

A.A. Borges¹, M.V.O. Santos¹, G.P.O. Lira¹, L.E. Nascimento¹, E.A. Praxedes¹, A.R. Silva²,
M.F. Oliveira³, A.F. Pereira¹

¹Laboratory of Animal Biotechnology; ²Laboratory of Animal Germplasm Conservation, ³Laboratory of Applied Animal Morphophysiology, Federal Rural University of Semi-Arid, Mossoró, RN, Brazil.

Oocyte *in vitro* maturation (IVM) is the first step of the *in vitro* reproductive technologies that enables matured oocytes to be used for embryo production. In this sense, the establishment of culture environment, as medium composition, is essential for the success of the IVM. In some domestic mammals, the supplementation of IVM medium with 10 ng/mL of epidermal growth factor (EGF) increased the IVM rates, and we hypothesized that this effect could also be observed for collared peccary oocytes. Therefore, the aim was to evaluate the EGF on the IVM rates of collared peccary oocytes, wild mammals of great commercial and ecological interest. Thus, eight adult collared peccaries (two females per session) were ovarian stimulated with 600 IU of PG600® (400 IU eCG and 200 IU hCG, MSD Saúde Animal, São Paulo, Brazil). Four days after hormone administration, ovaries were recovered, and all visible follicles with a 3–6 mm diameter were aspirated for the oocyte recovery using a 21 G needle attached to a 5.0 mL syringe containing oocyte recovery medium. Oocytes were classified with a stereomicroscope and only oocytes with more than one layer of *cumulus* cells and homogeneous cytoplasm were used for IVM. Thus, oocytes were matured in TCM199 contained 20 µg/mL of FSH-LH, 10% of fetal bovine serum, 100 µM of cysteamine, 1% of antibiotic-antimycotic solution and in the absence (group without EGF) or presence of 10 ng/mL of EGF (group with EGF). Oocytes were divided randomly in both groups and matured *in vitro* for 44 h at 38.5°C in a humidified atmosphere with 5% CO₂. Immediately after the IVM, oocytes were evaluated for expansion of *cumulus* cells using a stereomicroscope. After, oocytes were denuded and assessed for the presence of first polar body using a stereomicroscope. Finally, to evaluate the nuclear stage, denuded oocytes were fixed in 4% paraformaldehyde in PBS for 30 min. Then, oocytes were stained with Hoechst 33342 (10 µg/mL) for 15 min and visualized with a fluorescent microscope for identification of nuclear status in second metaphase. All data were expressed as mean ± standard error and analysed by the chi-square test ($P < 0.05$). A total of 172 follicles were aspirated after four sessions of ovarian stimulation. The oocyte recovery rate was 69.8% ± 8.3 (120/172) with an average of 7.5 ± 2.4 oocytes per ovary, and 5.3 ± 1.7 viable oocytes per ovary. After the IVM, no difference ($P > 0.05$) was observed between oocytes matured in absence and presence of EGF for expansion of the *cumulus* cells (97.6% ± 1.2 vs. 100% ± 0.0), presence of first polar body (65.9% ± 1.2 vs. 70.5% ± 1.8) and nuclear status in second metaphase (62.5% ± 11.6 vs. 68.4% ± 4.9), respectively. Probably, as occurred in swine oocytes, domestic mammals phylogenetically closely to the collared peccaries, the presence of EGF during the IVM can act in the protein synthesis and this effect could be observed during the embryonic development. In conclusion, according to the meiotic potential observed in collared peccary oocytes, EGF has not improved the IVM rates in this species.

Financial support: MSD Saúde Animal, CNPq, and CAPES.

E-mail: alexsandra.pereira@ufersa.edu.br

1 **ANEXO L – RESUMO PUBLICADO NO CBRA 2019**
2 **CONGRESSO BRASILEIRO DE REPRODUÇÃO ANIMAL**

Efeito da privação de soro sobre a sincronização do ciclo celular de fibroblastos derivados da pele de catetos, *Pecari tajacu* (Linnaeus, 1758)

*Effect of serum starvation on cell cycle synchronization fibroblast derived from skin collared peccaries, *Pecari tajacu* (Linnaeus, 1758)*

**Alana Azevedo Borges¹, Maria Claudia dos Santos Luciano², Matheus Barbosa do Nascimento³,
Gabriela Pereira de Oliveira Lira¹, Fátima de Cássia Evangelista de Oliveira⁴,
Claudia do Ó Pessoa⁵, Alexsandra Fernandes Pereira^{6,*}**

¹Pós-graduandas pelo Programa de Pós-Graduação em Ciência Animal, Universidade Federal Rural do Semi-Árido (UFERSA), Mossoró, RN, Brasil; ²Doutoranda pela Rede Nordeste de Biotecnologia, Universidade Federal do Ceará (UFC), Fortaleza, CE, Brasil; ³Graduando do Curso de Biotecnologia, CCBS, UFERSA, Mossoró, RN, Brasil;

⁴Pós-Doutora do Laboratório de Oncologia Experimental no Núcleo de Pesquisa e Desenvolvimento de Medicamentos, UFC, Fortaleza, CE, Brasil; ⁵Docente do Departamento de Fisiologia e Farmacologia, UFC, Fortaleza, CE, Brasil; ⁶Docente do Curso de Biotecnologia, Centro de Ciências Biológicas e da Saúde (CCBS), UFERSA, Mossoró, RN, Brasil.

*E-mail: alexsandra.pereira@ufersa.edu.br

A clonagem por transferência nuclear de células somáticas consiste numa alternativa para a conservação de catetos, mamíferos silvestres de grande relevância por seu potencial econômico, científico e ecológico. Nesse contexto, a otimização das etapas envolvidas nessa biotécnica, como a sincronização dos carioplastos em G0/G1 do ciclo, são cruciais para o seu sucesso na espécie de interesse. Portanto, o objetivo foi avaliar o efeito da privação de soro fetal bovino (SFB) sobre a sincronização na fase G0/G1 de fibroblastos derivados de catetos. Para tanto, foram utilizados fibroblastos criopreservados da terceira passagem obtidos de fragmentos de pele de quatro catetos adultos. Após a descongelação, fibroblastos cultivados e com 70% de confluência tiveram o meio de cultivo com 10% de SFB substituído por 0,5% de SFB. Em seguida, células foram avaliadas de 24 h a 96 h de cultivo em privação de soro. Células em crescimento com 70% de confluência e não submetidas à supressão de SFB foram consideradas como grupo controle. Para análise do ciclo celular, células ao final do seu período de tratamento foram tripsinizadas, centrifugadas, fixadas em etanol e armazenadas a -4°C. Posteriormente, as células foram incubadas em solução composta por iodeto de propídio (20 µg/mL) e RNase (50 µg/mL) por 50 min. Subsequentemente, todas as células foram analisadas por citômetro de fluxo Guava EasyCyte Desktop (Guava Technologies). Para cada amostra, 15.000 eventos foram registrados, e histogramas gerados para avaliar o percentual de células em cada fase do ciclo celular (G0/G1, S, G2/M) usando o software MODFIT versão 5.0. Os dados foram expressos como média ± erro padrão e analisados pelo software GraphPad ($P < 0,05$). Assim, após cinco repetições (um animal/repetição), fibroblastos submetidos à privação de SFB por 96 h apresentaram um maior percentual de G0/G1 ($79,0\% \pm 1,6$), quando comparados aos fibroblastos não submetidos à sincronização ($68,1\% \pm 8,5$, $P < 0,05$). Além disso, nenhuma diferença foi observada entre os fibroblastos não sincronizados e sincronizados por 24 h ($78,7\% \pm 2,3$), 48 h ($78,1\% \pm 1,7$) e 72 h ($75,8\% \pm 2,9$). Adicionalmente, nenhuma diferença foi observada entre as demais fases (S, G2/M) em nenhum dos períodos avaliados ($P > 0,05$). Portanto, a privação de SFB por 96 h promoveu a sincronização de fibroblastos de catetos na fase G0/G1. Estes resultados são relevantes para o desenvolvimento da técnica de clonagem em catetos.

Palavras-chave: carioplasto, transferência nuclear de células somáticas, mamíferos silvestres.

Keywords: karyoplast, somatic cell nuclear transfer, wild mammals.

1	ANEXO M – RESUMOS PUBLICADOS NO RENORBIO 2019
2	II ENCONTRO DE BIOTECNOLOGIA NO NORDESTE

RENORBIO 2019 - II Encontro de Biotecnologia do Nordeste

Avaliação dos aspectos morfométricos de ovários oriundos de catetos após estimulação hormonal

Maria Bárbara Silva*, Alana Azevedo Borges, Maria Valéria de Oliveira Santos, Lucas Emanuel Nascimento, Gabriela Pereira de Oliveira Lira, Érika Almeida Praxedes, Moacir Franco de Oliveira, Alexandre Rodrigues Silva, Alexsandra Fernandes Pereira

Laboratório de Biotecnologia Animal, Universidade Federal Rural do Semi-Árido, Mossoró, RN, Brasil. *E-mail: mariabarbarasilva16@gmail.com

Suporte financeiro: CNPq, CAPES e MSD Saúde Animal.

Palavras-chave: mamíferos silvestres, desempenho reprodutivo, protocolos hormonais,

A produção *in vitro* de embriões está diretamente ligada à função ovariana, uma vez que os ovários possuem funções essenciais na produção de oócitos a ser empregados nas biotecnologias reprodutivas, como a fecundação *in vitro*. Nesse sentido, protocolos de estimulação ovariana são essenciais na recuperação de um maior quantitativo de oócitos. Contudo, a identificação dos aspectos morfométricos dos ovários é etapa importante para o aprimoramento desses protocolos. Portanto, o objetivo foi avaliar os aspectos morfométricos de ovários direitos e esquerdos de catetos, mamíferos silvestres de significativa importância econômica em virtude da qualidade de seus produtos (carne e couro). Para tanto, vinte fêmeas com idade de $25,1 \pm 2,5$ meses foram estimuladas com 600 UI de PG600® por via intramuscular (400 UI eCG e 200 UI hCG). Após quatro dias da administração hormonal, ovários foram recuperados e transportados ao laboratório em NaCl a 0,9% (37°C por 30 min). No laboratório, com o auxílio de um paquímetro digital, ovários direitos e esquerdos foram mensurados, quanto ao comprimento (mm), largura (mm) e espessura (mm). Além disso, usando uma balança digital, ovários foram mensurados quanto ao volume (mL) e peso (g). Todos os dados foram expressos como média $\pm$ erro padrão e analisados por ANOVA seguida por teste de Tukey ($P < 0,05$). Nenhuma diferença foi observada entre os ovários direitos e esquerdos, respectivamente, para comprimento ($21,2 \pm 0,9$ mm vs. $22,4 \pm 0,9$ mm), largura ($16,1 \pm 0,8$ mm vs. $16,9 \pm 0,8$ mm), espessura ($3,0 \pm 0,3$ mm vs. $3,8 \pm 0,4$ mm) e peso ($3,0 \pm 0,3$ g vs. $3,3 \pm 0,3$ g). Portanto, a estimulação hormonal ovariana com PG600® promoveu padrão similar morfométrico entre os ovários direitos e esquerdos de catetos.

RENORBIO 2019 - II Encontro de Biotecnologia do Nordeste

Efeito do fator de crescimento epidermal sobre a morfometria de oócitos maturados de catetos

Gabriela Pereira de Oliveira Lira*, Alana Azevedo Borges, Maria Valéria de Oliveira Santos, Lucas Emanuel Nascimento, Érika Almeida Praxedes, Moacir Franco de Oliveira, Alexandre Rodrigues Silva, Alexsandra Fernandes Pereira

Laboratório de Biotecnologia Animal, Universidade Federal Rural do Semi-Árido, Mossoró, RN, Brasil. *E-mail: gabrielaliraop@gmail.com

Suporte financeiro: CNPq, CAPES e MSD Saúde Animal.

Palavras-chave: mamíferos silvestres, maturação *in vitro*, produção *in vitro* de embriões.

A produção *in vitro* de embriões (PIVE) é o foco central da agropecuária, principalmente em virtude do interesse no aumento da produtividade por meio da multiplicação de indivíduos interessantes economicamente e de alta adaptação a cativeiros. Nesse cenário, incluem-se os catetos, os quais possuem uma interessante capacidade produtiva e potencial de mercado para seus produtos, especialmente em países da América Latina. Contudo, a eficiência da PIVE depende da bem-sucedida maturação *in vitro* (MIV) de oócitos. Portanto, o objetivo foi avaliar o efeito do fator de crescimento epidermal (EGF) sobre a morfometria de oócitos maturados de catetos. Para tanto, oócitos viáveis morfologicamente e derivados de oito fêmeas adultas e previamente submetidas à estimulação hormonal ovariana por quatro dias com 600 UI de PG600® (400 UI eCG e 200 UI hCG) foram maturados *in vitro*. Para a MIV, oócitos foram incubados por 44 h (38,5°C e 5% de CO₂) de acordo com os grupos: ausência ou presença de 10 ng/mL de EGF. Após a MIV, oócitos foram desnudos e avaliados morfometricamente, quanto diâmetro externo do oócito (DEO); espessura da zona pelúcida (EZO); diâmetro do ooplasma (DO); área do espaço perivitelino (AEP) e comprimento do espaço perivitelino (CEP). Tais parâmetros foram analisados por meio do software ImageJ, utilizando-se a barra de escala como escala arbitrária para análise de pixels. Os dados foram expressos como média $\pm$ erro padrão de quatro repetições (dois animais por repetição) e analisados a uma significância de $P < 0,05$. Nenhuma diferença foi observada entre os grupos maturados na ausência e presença de EGF, respectivamente, para DEO ($382,2 \pm 5,3 \mu\text{m}$ vs. $373,9 \pm 4,4 \mu\text{m}$), DO ($285,6 \pm 3,0$ vs. $284,3 \pm 4,2 \mu\text{m}$), AEP ($6284,0 \pm 390,9$ vs. $7312,6 \pm 823,1 \mu\text{m}^2$) e CEP ($13,8 \pm 0,8$ vs. $16,2 \pm 2,1 \mu\text{m}$). Contudo, quanto à EZO, oócitos maturados na presença de EGF apresentaram uma redução de sua espessura ($33,3 \pm 1,2 \mu\text{m}$), quando comparados aos oócitos maturados na ausência de EGF ($38,6 \pm 1,5 \mu\text{m}$). Esta diferença implica em uma característica positiva uma vez que a espessura da zona pelúcida consiste em um indicativo relevante para a eclosão e implantação de blastocistos derivados desses oócitos. Portanto, de acordo com os parâmetros morfométricos, sugere-se o uso do EGF como suplementação no meio de MIV de oócitos de catetos.

1 **ANEXO N – RESUMO PUBLICADO NO SEMIC 2019**
2 **XXV SEMINÁRIO DE INICIAÇÃO CIENTÍFICA DA UFERSA**
3

Área do conhecimento: Ciências Biológicas

EFEITO DO FATOR DE CRESCIMENTO EPIDERMAL SOBRE A MATURAÇÃO *in vitro* DE OÓCITOS DE CATETOS, *Pecari tajacu* LINNAEUS, 1758 (ARTIODACTYLA: TAYASSUIDAE)

Luanna Lorena Vieira Rodrigues; Alessandra Fernandes Pereira; Lucas Emanuel Nascimento; Alana Azevedo Borges; Maria Valéria de Oliveira Santos

A produção *in vitro* de embriões (PIVE) tem se estabelecido como uma ferramenta interessante para a conservação de mamíferos silvestres, como os catetos. Nesse contexto, o desenvolvimento de meios adequados para a maturação *in vitro* (MIV) consiste numa etapa fundamental para a eficiência da PIVE. Portanto, o objetivo foi avaliar o efeito do fator de crescimento epidermal (EGF) durante a MIV de complexos *cumulus*-oócito (CCOs) derivados de catetos. Inicialmente, todos os procedimentos foram aprovados pelo Comitê de Ética e Uso de Animais (CEUA/UFERSA, no. 23091.001072/2015-92) e Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio, no. 48633-2). Para tanto, oito fêmeas foram estimuladas hormonalmente com 600 UI de PG600® e, após a eutanásia dos animais no quarto dia de aplicação, os ovários foram removidos e transportados até o laboratório. Os folículos ovarianos com 2–6 mm de diâmetro foram aspirados e os CCOs apresentando mais de uma camada de células do *cumulus* e citoplasma homogêneo foram direcionados à MIV. Assim, CCOs foram maturados por 44 h (38,5°C e 5% de CO₂) em meio TCM199 modificado e na ausência e presença de 10 ng/mL de EGF. Após a MIV, oócitos foram avaliados quanto à expansão e viabilidade das células do *cumulus* por azul de tripan, maturação nuclear pela presença do primeiro corpúsculo polar (1CP) e níveis de espécies reativas de oxigênio (EROs) usando sonda fluorescente. Todos os dados foram expressos como média $\pm$ erro padrão e comparações foram realizadas usando o teste ANOVA seguido de teste de Tukey ($P < 0,05$). Após quatro repetições (dois animais/repetição), foram obtidos 16 ovários, resultando em 231 folículos aspirados e 120 CCOs recuperados, perfazendo uma taxa de recuperação de 49,1%. Assim, de acordo com a avaliação e classificação morfológica, 85 (70,8%) CCOs viáveis e 35 (29,2%) CCOs não viáveis foram recuperados, sendo empregado para a etapa de MIV apenas os CCOs viáveis. Após a MIV, os grupos avaliados mostraram altas taxas de expansão das células do *cumulus* (ausência de EGF: $98,8\% \pm 1,2$ e presença de EGF: $100\% \pm 0,0$) com apenas expansão nível III e IV identificadas ($P > 0,05$). Quanto à viabilidade das células do *cumulus*, nenhuma diferença foi observada entre os grupos na ausência ($79,4\% \pm 2,7$) e presença de EGF ($79,4\% \pm 2,7$). Além disso, nenhuma diferença foi observada entre os grupos de CCOs maturados na ausência e presença de EGF para as taxas de maturação nuclear ($73,1\% \pm 9,0$ vs. $76,0\% \pm 8,1$) e níveis de EROs ($0,3 \pm 0,1$ vs. $0,5 \pm 0,2$ unidades de fluorescência arbitrária). Em conclusão, o EGF não influencia as taxas de maturação nuclear, expansão e viabilidade das células do *cumulus* e níveis de EROs em CCOs de catetos. Provavelmente, o efeito dessa suplementação poderia ser melhor visualizado durante o desenvolvimento embrionário.

Palavras-chave: Mamíferos silvestres. Produção de embriões. Complexos *cumulus*-oócito.
Agência financiadora: Bolsista IC PIBIC.